

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :

C12N 15/12, C07K 14/52, C12N 15/10,
C07K 16/24, C12Q 1/68, C12N 15/62,
G01N 33/50, A01K 67/027

A1

(11) International Publication Number:

WO 98/33917

(43) International Publication Date:

6 August 1998 (06.08.98)

(21) International Application Number:

PCT/US98/01973

(22) International Filing Date:

2 February 1998 (02.02.98)

(30) Priority Data:

08/795,430

5 February 1997 (05.02.97)

US

(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications

US

08/795,430 (CIP)

Filed on

5 February 1997 (05.02.97)

US

PCT/FI96/00427 (CIP)

Filed on

1 August 1996 (01.08.96)

US

08/671,573 (CIP)

Filed on

28 June 1996 (28.06.96)

US

08/601,132 (CIP)

Filed on

14 February 1996 (14.02.96)

US

08/585,895 (CIP)

Filed on

12 January 1996 (12.01.96)

US

08/510,133 (CIP)

Filed on

1 August 1995 (01.08.95)

US

08/340,011 (CIP)

Filed on

14 November 1994 (14.11.94)

(71) Applicants (for all designated States except US): THE LUDWIG INSTITUTE FOR CANCER RESEARCH [US/US]; 1345 Avenue of the Americas, New York, NY 10105 (US). HELSINKI UNIVERSITY LICENSING LTD. [FI/FI]; Viikinkaari 8 A, FIN-00170 Helsinki (FI).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ALITALO, Kari [FI/FI]; Nyyrikintie 4A, FIN-02100 Espoo (FI). JOUKOV, Vladimir [RU/US]; Apartment 1F, 51 Massachusetts Avenue, Boston, MA 02115 (US).

(74) Agents: GASS, David, A. et al.; Marshall, O'Toole, Gerstein Murray & Borun, 6300 Sears Tower, 233 South Wacker Drive, Chicago IL 60606-6402 (US).


(81) Designated States: AU, CA, CN, JP, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: VASCULAR ENDOTHELIAL GROWTH FACTOR C (VEGF-C) PROTEIN AND GENE, MUTANTS THEREOF, AND USES THEREOF



(57) Abstract

Provided are purified and isolated VEGF-C polypeptides capable of binding to at least one of KDR receptor tyrosine kinase (VEGFR-2) and Flt4 receptor tyrosine kinase (VEGFR-3); analogs of such peptides that have VEGF-C-like or VEGF-like biological activities or that are VEGF or VEGF-C inhibitors; polynucleotides encoding the polypeptides; vectors and host cells that embody the polynucleotides; pharmaceutical compositions and diagnostic reagents comprising the polypeptides; and methods of making and using the polypeptides.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

**VASCULAR ENDOTHELIAL GROWTH FACTOR C (VEGF-C)
PROTEIN AND GENE, MUTANTS THEREOF, AND USES THEREOF**

This application is a continuation-in-part of United States Patent Application Serial No. 08/795,430, filed February 5, 1997; and a continuation-in-part of
5 International Patent Application PCT/FI96/00427, filed August 01, 1996; and a continuation-in-part of United States Patent Application Serial No. 08/671,573, filed June 28, 1996; and a continuation-in-part of United States Patent Application Serial Number 08/601,132, filed February 14, 1996; and a continuation-in-part of United States Patent Application Serial Number 08/585,895, filed January 12, 1996; and a continuation-in-part
10 of United States Patent Application Serial Number 08/510,133, filed August 1, 1995; and a continuation-in-part of United States Patent Application Serial Number 08/340,011, filed November 14, 1994.

FIELD OF THE INVENTION

The present invention generally relates to the field of genetic engineering
15 and more particularly to growth factors for endothelial cells and growth factor genes.

BACKGROUND OF THE INVENTION

Developmental growth, the remodeling and regeneration of adult tissues, as well as solid tumor growth, can only occur when accompanied by blood vessel formation. Angioblasts and hematopoietic precursor cells differentiate from the mesoderm and form
20 the blood islands of the yolk sac and the primary vascular system of the embryo. The development of blood vessels from these early (*in situ*) differentiating endothelial cells is termed vasculogenesis. Major embryonic blood vessels are believed to arise via vasculogenesis, whereas the formation of the rest of the vascular tree is thought to occur as a result of vascular sprouting from pre-existing vessels, a process called angiogenesis,
25 Risau *et al.*, *Devel. Biol.*, 125:441-450 (1988).

Endothelial cells give rise to several types of functionally and morphologically distinct vessels. When organs differentiate and begin to perform their specific functions, the phenotypic heterogeneity of endothelial cells increases. Upon angiogenic stimulation, endothelial cells may re-enter the cell cycle, migrate, withdraw
30 from the cell cycle and subsequently differentiate again to form new vessels that are

functionally adapted to their tissue environment. Endothelial cells undergoing angiogenesis degrade the underlying basement membrane and migrate, forming capillary sprouts that project into the perivascular stroma. Ausprunk *et al.*, *Microvasc. Rev.*, 14:51-65 (1977). Angiogenesis during tissue development and regeneration depends on the tightly controlled processes of endothelial cell proliferation, migration, differentiation, and survival. Dysfunction of the endothelial cell regulatory system is a key feature of many diseases. Most significantly, tumor growth and metastasis have been shown to be angiogenesis dependent. Folkman *et al.*, *J. Biol. Chem.*, 267:10931-10934 (1992).

Key signals regulating cell growth and differentiation are mediated by polypeptide growth factors and their transmembrane receptors, many of which are tyrosine kinases. Autophosphorylated peptides within the tyrosine kinase insert and carboxyl-terminal sequences of activated receptors are commonly recognized by kinase substrates involved in signal transduction for the readjustment of gene expression in responding cells. Several families of receptor tyrosine kinases have been characterized. Van der Geer *et al.*, *Ann. Rev. Cell Biol.*, 10:251-337 (1994). The major growth factors and receptors transducing angiogenic stimuli are schematically shown in Fig. 1.

Fibroblast growth factors are also known to be involved in the regulation of angiogenesis. They have been shown to be mitogenic and chemotactic for cultured endothelial cells. Fibroblast growth factors also stimulate the production of proteases, such as collagenases and plasminogen activators, and induce tube formation by endothelial cells. Saksela *et al.*, *Ann. Rev. Cell Biol.*, 4:93-126 (1988). There are two general classes of fibroblast growth factors, FGF-1 and FGF-2, both of which lack conventional signal peptides. Both types have an affinity for heparin, and FGF-2 is bound to heparin sulfate proteoglycans in the subendothelial extracellular matrix from which it may be released after injury. Heparin potentiates the stimulation of endothelial cell proliferation by angiogenic FGFs, both by protecting against denaturation and degradation and dimerizing the FGFs. Cultured endothelial cells express the FGF-1 receptor but no significant levels of other high-affinity fibroblast growth factor receptors.

Among other ligands for receptor tyrosine kinases, the platelet derived growth factor, PDGF-BB, has been shown to be weakly angiogenic in the chick chorioallantoic membrane. Risau *et al.*, *Growth Factors*, 7:261-266 (1992). Transforming growth factor α (TGF α) is an angiogenic factor secreted by several tumor cell types and

by macrophages. Hepatocyte growth factor (HGF), the ligand of the *c-met* proto-oncogene-encoded receptor, also is strongly angiogenic.

Recent evidence shows that there are endothelial cell specific growth factors and receptors that may be primarily responsible for the stimulation of endothelial cell growth, differentiation and certain differentiated functions. The best studied of these is vascular endothelial growth factor (VEGF), a member of the PDGF family. Vascular endothelial growth factor is a dimeric glycoprotein of disulfide-linked 23 kD subunits. Other reported effects of VEGF include the mobilization of intracellular calcium, the induction of plasminogen activator and plasminogen activator inhibitor-1 synthesis, stimulation of hexose transport in endothelial cells, and promotion of monocyte migration *in vitro*. Four VEGF isoforms, encoded by distinct mRNA splice variants, appear to be equally capable of stimulating mitogenesis in endothelial cells. However, each isoform has a different affinity for cell surface proteoglycans, which behave as low affinity receptors for VEGF. The 121 and 165 amino acid isoforms of VEGF (VEGF₁₂₁ and VEGF₁₆₅) are secreted in a soluble form, whereas the isoforms of 189 and 206 amino acid residues remain cell surface-associated and have a strong affinity for heparin. VEGF was originally purified from several sources on the basis of its mitogenic activity toward endothelial cells, and also by its ability to induce microvascular permeability, hence it is also called vascular permeability factor (VPF).

Two high affinity receptors for VEGF have been characterized: VEGFR-1/Flt-1 (fms-like tyrosine kinase-1) and VEGFR-2/KDR/Flk-1 (kinase insert domain containing receptor/fetal liver kinase-1). Those receptors are classified in the PDGF-receptor family, but they have seven rather than five immunoglobulin-like loops in their extracellular domain (see Fig. 1), and they possess a longer kinase insert than normally observed in this family. The expression of VEGF receptors occurs mainly in vascular endothelial cells, although some may be present on hematopoietic progenitor cells, monocytes, and melanoma cells. Only endothelial cells have been reported to proliferate in response to VEGF, and endothelial cells from different sources show different responses. Thus, the signals mediated through VEGFR-1 and VEGFR-2 appear to be cell type specific. The VEGF-related placenta growth factor (PlGF) was recently shown to bind to VEGFR-1 with high affinity. PlGF was able to enhance the growth factor activity of VEGF, but it did not stimulate endothelial cells on its own. Naturally occurring

VEGF/PlGF heterodimers were nearly as potent mitogens as VEGF homodimers for endothelial cells. Cao *et al.*, *J. Biol. Chem.*, 271:3154-62 (1996).

The Flt4 receptor tyrosine kinase (VEGFR-3) is closely related in structure to the products of the VEGFR-1 and VEGFR-2 genes. Despite this similarity, the mature
5 form of Flt4 differs from the VEGF receptors in that it is proteolytically cleaved in the extracellular domain into two disulfide-linked polypeptides. Pajusola *et al.*, *Cancer Res.*, 52:5738-5743 (1992). The 4.5 and 5.8 kb Flt4 mRNAs encode polypeptides which differ in their C-termini due to the use of alternative 3' exons. Isoforms of VEGF or PlGF do not show high affinity binding to Flt4 or induce its autophosphorylation.

10 Expression of Flt4 appears to be more restricted than the expression of VEGFR-1 or VEGFR-2. The expression of Flt4 first becomes detectable by *in situ* hybridization in the angioblasts of head mesenchyme, the cardinal vein, and extraembryonically in the allantois of 8.5 day p.c. mouse embryos. In 12.5 day p.c. embryos, the Flt4 signal is observed in developing venous and presumptive lymphatic
15 endothelia, but arterial endothelia appear negative. During later stages of development, Flt4 mRNA becomes restricted to developing lymphatic vessels. The lymphatic endothelia and some high endothelial venules express Flt4 mRNA in adult human tissues and increased expression occurs in lymphatic sinuses in metastatic lymph nodes and in lymphangioma. These results support the theory of the venous origin of lymphatic vessels.

20 Five endothelial cell specific receptor tyrosine kinases, Flt-1 (VEGFR-1), KDR/Flk-1 (VEGFR-2), Flt4 (VEGFR-3), Tie, and Tek/Tie-2 have so far been described, which possess the intrinsic tyrosine kinase activity essential for signal transduction. Targeted mutations inactivating Flt-1, Flk-1, Tie, and Tek in mouse embryos have indicated their essential and specific roles in vasculogenesis and angiogenesis at the
25 molecular level. VEGFR-1 and VEGFR-2 bind VEGF with high affinity (K_d 16 pM and 760 pM, respectively) and VEGFR-1 also binds the related placenta growth factor (PlGF; K_d about 200 pM). A ligand for Tek is reported in PCT patent publication WO 96/11269.

SUMMARY OF THE INVENTION

The present invention provides a ligand, designated VEGF-C, for the Flt4
30 receptor tyrosine kinase (VEGFR-3). Thus, the invention provides a purified and isolated polypeptide which is capable of binding to the Flt4 receptor tyrosine kinase. Preferably, an

- 5 -

Flt4 ligand of the invention is capable of stimulating tyrosine phosphorylation of Flt4 receptor tyrosine kinase in a host cell expressing the Flt4 receptor tyrosine kinase.

Preferred ligands of the invention are mammalian polypeptides. Highly preferred ligands are human polypeptides. As explained in detail below, dimers and multimers comprising 5 polypeptides of the invention linked to each other or to other polypeptides are specifically contemplated as aspects of the invention.

In one embodiment, an Flt4 ligand polypeptide has a molecular weight of approximately 23 kD as determined by SDS-PAGE under reducing conditions. For example, the invention includes a ligand composed of one or more polypeptides of 10 approximately 23 kD which is purifiable from conditioned media from a PC-3 prostatic adenocarcinoma cell line, the cell line having ATCC Acc. No. CRL 1435. Amino acid sequencing of this PC-3 cell-derived ligand polypeptide revealed that the ligand polypeptide comprises an amino terminal amino acid sequence set forth in SEQ ID NO: 5. The present invention also provides a new use for the PC-3 prostatic adenocarcinoma cell 15 line which produces an Flt4 ligand. In a preferred embodiment, the ligand may be purified and isolated directly from the PC-3 cell culture medium.

In a highly preferred embodiment, the ligand polypeptide comprises a fragment of the amino acid sequence shown in SEQ ID NO: 8 which binds with high affinity to the human Flt4 receptor tyrosine kinase. It will be understood that the term 20 "high affinity," in the context of a polypeptide ligand of a receptor tyrosine kinase, typically reflects a binding relationship characterized by sub-nanomolar dissociation constants (K_d), as reported herein for VEGF-C binding to VEGFR-2 and VEGFR-3, and reported elsewhere in the art for the binding of VEGF, PlGF, PDGF, and other factors to their receptors. Exemplary fragments include: a polypeptide comprising an amino acid 25 sequence set forth in SEQ ID NO: 8 from about residue 112 to about residue 213; a polypeptide comprising an amino acid sequence from about residue 104 to about residue 227 of SEQ ID NO: 8; and a polypeptide comprising an amino acid sequence from about residue 112 to about residue 227 of SEQ ID NO: 8. Other exemplary fragments include polypeptides comprising amino acid sequences of SEQ ID NO: 8 that span, approximately, 30 the following residues: 31-213, 31-227, 32-227, 103-217, 103-225, 104-213, 113-213, 103-227, 113-227, 131-211, 161-211, 103-225, 227-419, 228-419, 31-419, and 1-419, as described in greater detail below.

The present invention also provides one or more polypeptide precursors of an Flt4 ligand, wherein one such precursor (designated "prepro-VEGF-C") comprises the complete amino acid sequence (amino acid residues 1 to 419) shown in SEQ ID NO: 8.

Thus, the invention includes a purified and isolated polypeptide having the amino acid
5 sequence of residues 1 to 419 shown in SEQ ID NO: 8. Ligand precursors according to the invention, when expressed in an appropriate host cell, produce, via cleavage, a polypeptide which binds with high affinity to the Flt4 receptor tyrosine kinase. A putative 102 amino acid leader (prepro) peptide has been identified in the amino acid sequence shown in SEQ ID NO: 8. Thus, in a related aspect, the invention includes a purified and
10 isolated polypeptide having the amino acid sequence of residues 103-419 shown in SEQ ID NO: 8.

In one embodiment, an expressed Flt4 ligand polypeptide precursor is proteolytically cleaved upon expression to produce an approximately 23 kD Flt4 ligand polypeptide. Thus, an Flt4 ligand polypeptide is provided which is the cleavage product of
15 the precursor polypeptide shown in SEQ ID NO: 8 and which has a molecular weight of approximately 23 kD under reducing conditions.

Putative VEGF-C precursors/processing products consisting of polypeptides with molecular weights of about 29 and 32 kD also are considered aspects of the invention.

20 In another embodiment, an expressed Flt4 ligand polypeptide precursor is proteolytically cleaved upon expression to produce an approximately 21 kD VEGF-C polypeptide. Sequence analysis has indicated that an observed 21 kD form has an amino terminus approximately 9 amino acids downstream from the amino terminus of the 23 kD form, suggesting that alternative cleavage sites exist.

25 From the foregoing, it will be apparent that an aspect of the invention includes a fragment of the purified and isolated polypeptide having the amino acid sequence of residues 1 to 419 shown in SEQ ID NO: 8, the fragment being capable of binding with high affinity to Flt4 receptor tyrosine kinase. Preferred embodiments include fragments having an apparent molecular weight of approximately 21/23 kD and 29/32 kD
30 as assessed by SDS-PAGE under reducing conditions. More generally, the invention includes a purified and isolated polypeptide that is a VEGF-C of vertebrate origin, wherein the VEGF-C has a molecular weight of about 21-23 kD, as assessed by SDS-PAGE under

- 7 -

reducing conditions, and wherein the VEGF-C is capable of binding to Flt4 receptor tyrosine kinase (VEGFR-3). Vertebrate VEGF-C forms of about 30-32 kD that are capable of binding VEGFR-3 also are intended as an aspect of the invention.

Evidence suggests that the amino acids essential for retaining Flt4 ligand activity are contained within approximately amino acids 103/112-226/227 of SEQ ID NO: 8, and that a carboxy-terminal proteolytic cleavage to produce a mature, naturally-occurring Flt4 ligand occurs at the approximate position of amino acids 226-227 of SEQ ID NO: 8. Accordingly, a preferred Flt4 ligand comprises approximately amino acids 103-227 of SEQ ID NO: 8.

10 VEGF-C mutational analysis described herein indicates that a naturally occurring VEGF-C polypeptide spanning amino acids 103-227 of SEQ ID NO: 8, produced by a natural processing cleavage that defines the C-terminus, exists and is biologically active as an Flt4 ligand. A polypeptide fragment consisting of residues 104-213 of SEQ ID NO: 8 has been shown to retain VEGF-C biological activity. Additional
15 mutational analyses indicate that a polypeptide spanning only amino acids 113-213 of SEQ ID NO: 8 retains Flt4 ligand activity. Accordingly, preferred polypeptides comprise sequences spanning, approximately, amino acid residues 103-227, 104-213, or 113-213, of SEQ ID NO: 8.

Moreover, sequence comparisons of members of the VEGF family of
20 polypeptides provide an indication that still smaller fragments will retain biological activity, and such smaller fragments are intended as aspects of the invention. In particular, eight highly conserved cysteine residues of the VEGF family of polypeptides define a region from residue 131 to residue 211 of SEQ ID NO: 8 (see Figures 2, 5 & 10); therefore, a polypeptide spanning from about residue 131 to about residue 211 is expected to retain
25 VEGF-C biological activity. In fact, a polypeptide comprising approximately residues 161-211, which retains an evolutionarily-conserved RCXXCC motif, is postulated to retain VEGF-C activity, and therefore is intended as an aspect of the invention.

In addition to binding Flt4, VEGF-C polypeptides are shown herein to bind and activate KDR/flk-1 receptor tyrosine kinase (VEGFR-2). Thus, the invention includes
30 a purified and isolated polypeptide that is capable of binding to at least one of KDR receptor tyrosine kinase (VEGFR-2) and Flt4 receptor tyrosine kinase (VEGFR-3), the polypeptide comprising a portion of the amino acid sequence in SEQ ID NO: 8 effective to

permit such binding. In one preferred embodiment, the portion of the amino acid sequence in SEQ ID NO: 8 is a continuous portion having as its amino terminal residue an amino acid between residues 102 and 161 of SEQ ID NO: 8 and having as its carboxy terminal residue an amino acid between residues 210 and 228 of SEQ ID NO: 8. In a highly preferred embodiment, the portion has, as its amino terminal residue, an amino acid between residues 102 and 131 of SEQ ID NO: 8. In a very highly preferred embodiment, the portion of the amino acid sequence in SEQ ID NO: 8 is a continuous portion having as its amino terminal residue an amino acid between residues 102 and 114 of SEQ ID NO: 8 and having as its carboxy terminal residue an amino acid between residues 212 and 228 of SEQ ID NO: 8. Polypeptides of the invention which bind to and activate a receptor (e.g., VEGFR-2 or VEGFR-3) are useful for stimulating VEGF-C biological activities that are mediated through the receptor. Polypeptides of the invention which bind to but do not activate a receptor are useful for inhibiting VEGF-C activities mediated through that receptor.

15 The definition of polypeptides of the invention is intended to include within its scope variants thereof. The polypeptide variants contemplated include purified and isolated polypeptides having amino acid sequences that differ from the exact amino acid sequences of such polypeptides (e.g., VEGF-C, VEGF-C precursors and VEGF-C fragments) by conservative substitutions, as recognized by those of skill in the art, that are
20 compatible with the retention of at least one VEGF-C biological activity or VEGF-C-inhibitory activity of the polypeptide. The term "variants," when used to refer to polypeptides, also is intended to include polypeptides having amino acid additions, including but not limited to additions of a methionine and/or leader sequence to promote translation and/or secretion; additions of peptide sequences to facilitate purification (e.g.,
25 polyhistidine sequences and/or epitopes for antibody purification); and additions of polypeptide-encoding sequences to produce fusion proteins with VEGF-C. The term "variants" also is intended to include polypeptides having amino acid deletions at the amino terminus, the carboxy terminus, or internally of amino acids that are non-conserved amongst the human, mouse, and quail VEGF-C sequences taught herein, and that are
30 compatible with the retention of the VEGF-C or VEGF-C-inhibitory activity of the polypeptide to which the deletions have been made.

The term "variant" also is intended to include polypeptides having modifications to one or more amino acid residues that are compatible with retaining VEGF-C or VEGF-C inhibitory activity of the polypeptide. Such modifications include glycosylations (identical or different to glycosylations of native VEGF-C); and the addition
5 of other substituents (e.g., labels, compounds to increase serum half-life (e.g., polyethylene glycol), and the like.

Additional polypeptides of the invention include certain fragments that have been observed to result from the processing of prepro-VEGF-C into mature VEGF-C. For example, the invention includes a purified and isolated polypeptide having a molecular
10 weight of about 29 kD as assessed by SDS-PAGE under reducing conditions and having an amino acid sequence consisting essentially of a portion of SEQ ID NO: 8 having residue 228 of SEQ ID NO: 8 as its amino terminal amino acid residue; and a purified and isolated polypeptide having a molecular weight of about 15 kD as assessed by SDS-PAGE under reducing conditions and having an amino acid sequence consisting essentially of a portion
15 of SEQ ID NO: 8 having residue 32 of SEQ ID NO: 8 as its amino terminal amino acid residue. Such polypeptides are expected to modulate VEGF-C biological activity through their interactions with VEGF-C receptors and/or interactions with biologically active VEGF-C.

Some of the conserved cysteine residues in VEGF-C participate in
20 interchain disulfide bonding to make homo- and heterodimers of the various naturally occurring VEGF-C polypeptides. Beyond the preceding considerations, evidence exists that VEGF-C polypeptides lacking interchain disulfide bonds retain VEGF-C biological activity. Consequently, the materials and methods of the invention include all VEGF-C fragments that retain at least one biological activity of VEGF-C, regardless of the presence
25 or absence of interchain disulfide bonds. The invention also includes multimers (including dimers) comprising such fragments linked to each other or to other polypeptides. Fragment linkage may be by way of covalent bonding (e.g., disulfide bonding) or non-covalent bonding of polypeptide chains (e.g., hydrogen bonding, bonding due to stable or induced dipole-dipole interactions, bonding due to hydrophobic or hydrophilic interactions,
30 combinations of these bonding mechanisms, and the like). Thus, the invention includes a purified and isolated polypeptide multimer, wherein at least one monomer thereof is a polypeptide that is capable of binding to VEGFR-2 and/or VEGFR-3, the polypeptide

- 10 -

comprising a portion of the amino acid sequence in SEQ ID NO: 8 effective to permit such binding, and wherein the multimer itself is capable of binding to VEGFR-2 and/or VEGFR-3. In a preferred embodiment, the multimer has at least one VEGF-C biological activity as taught herein.

5 In one embodiment, at least one monomer of the multimer is a polypeptide from another member of the PDGF/VEGF family of proteins, e.g., a vascular endothelial growth factor (VEGF) polypeptide, a vascular endothelial growth factor B (VEGF-B) polypeptide, a platelet derived growth factor A (PDGF-A) polypeptide, a platelet derived growth factor B (PDGF-B) polypeptide, a *c-fos* induced growth factor (FIGF)
10 polypeptide, or a placenta growth factor (PIGF) polypeptide.

In a highly preferred embodiment, the multimer of the invention is a dimer of two monomer polypeptides. For example, the invention includes a dimer wherein each monomer thereof is capable of binding to at least one of VEGFR-2 and VEGFR-3 and has an amino acid sequence comprising a portion of SEQ ID NO: 8 effective to permit such
15 binding. Dimers having covalent attachments and dimers wherein the two monomers are free of covalent attachments to each other are contemplated.

In yet another aspect, the invention includes analogs of the polypeptides of the invention. The term "analog" refers to polypeptides having alterations involving one or more amino acid insertions, internal amino acid deletions, and/or non-conservative amino
20 acid substitutions (replacements). The definition of analog is intended to include within its scope variants of analog polypeptides embodying such alterations. The term "mutant," when used with respect to polypeptides herein, is intended to refer generically to VEGF-C variants, VEGF-C analogs, and variants of VEGF-C analogs. Preferred analogs possess at least 90% amino acid sequence similarity to the native peptide sequence from which the
25 analogs were derived. Highly preferred analogs possess 95%, 96%, 97%, 98%, 99%, or greater amino acid sequence similarity to the native peptide sequence.

For example, in one embodiment, the invention includes a polypeptide analog of a VEGF-C of vertebrate origin that is capable of binding to VEGFR-3 (e.g., an analog of a vertebrate VEGF-C of about 21-23 kD as assessed by SDS-PAGE under
30 reducing conditions), wherein an evolutionarily conserved cysteine residue in the VEGF-C has been deleted or replaced, and wherein the analog is capable of binding to VEGFR-3 and has reduced VEGFR-2 binding affinity relative to the wildtype VEGF-C. For analogs

- 11 -

according to this embodiment of the invention, the determination that a residue is "evolutionarily conserved" is made solely by reference to the alignment of human, mouse, and quail VEGF-C sequences provided herein and aligned to show similarity in Fig. 5.

The presence of the same residue in all three sequences indicates that the residue is evolutionarily conserved, notwithstanding the fact that VEGF-C from other species may lack the residue. In a preferred embodiment, the conserved cysteine residue corresponds to the cysteine at position 156 of SEQ ID NO: 8. "Correspondence to the cysteine at position 156" is readily determined from an analysis of the vertebrate VEGF-C sequence of interest, since the cysteine at position 156 of SEQ ID NO: 8 (human VEGF-C) falls within an evolutionarily conserved portion of VEGF-C (see Fig. 5, comparing human, mouse, and quail VEGF-C polypeptides). Alignment of human VEGF-C allelic variants, other mammalian VEGF-C polypeptides, and the like with the three VEGF-C forms in Fig. 5 will identify that cysteine which corresponds to the cysteine at position 156 of SEQ ID NO: 8, even if the allelic variant has greater or fewer than exactly 155 residues preceding the cysteine of interest.

In another embodiment, the invention includes a purified polypeptide that is an analog of human VEGF-C and that is capable of binding to at least one of Flt-1 receptor tyrosine kinase (VEGFR-1), KDR receptor tyrosine kinase (VEGFR-2), and Flt4 receptor tyrosine kinase (VEGFR-3).

Specifically contemplated is an analog of human VEGF-C that binds VEGFR-3 but has reduced VEGFR-2 binding affinity, as compared to the VEGFR-2 binding affinity of a wildtype human VEGF-C (e.g., as compared to the VEGFR-2 binding affinity of a human VEGF-C having an amino acid sequence consisting essentially of amino acids 103-227 of SEQ ID NO: 8). One such family of human VEGF-C analogs are VEGF-C Δ_{156} polypeptides. By "VEGF-C Δ_{156} polypeptide" is meant an analog wherein the cysteine at position 156 of SEQ ID NO: 8 has been deleted or replaced by another amino acid. A VEGF-C Δ_{156} polypeptide analog can be made from any VEGF-C polypeptide of the invention that comprises all of SEQ ID NO: 8 or a portion thereof that includes position 156 of SEQ ID NO: 8. Preferably, the VEGF-C Δ_{156} polypeptide analog comprises a portion of SEQ ID NO: 8 effective to permit binding to VEGFR-3.

For example, the invention includes a VEGF-C Δ_{156} polypeptide that binds VEGFR-3, has reduced VEGFR-2 binding affinity, and has an amino acid sequence

- 12 -

which includes amino acids 131 to 211 of SEQ ID NO: 8, wherein the cysteine residue at position 156 of SEQ ID NO: 8 has been deleted or replaced. In a preferred embodiment, the VEGF-C ΔC_{156} polypeptide comprises a continuous portion of SEQ ID NO: 8, the portion having as its amino terminal residue an amino acid between residues 102 and 114 of SEQ ID NO: 8, and having as its carboxy terminal residue an amino acid between residues 212 and 228 of SEQ ID NO: 8, wherein the cysteine residue at position 156 of SEQ ID NO: 8 has been deleted or replaced. In an embodiment exemplified herein, the cysteine residue at position 156 of SEQ ID NO: 8 has been replaced by a serine residue.

A second family of human VEGF-C analogs that bind VEGFR-3 but have reduced VEGFR-2 binding affinity are VEGF-C $\Delta R_{226}\Delta R_{227}$ polypeptides. By "VEGF-C $\Delta R_{226}\Delta R_{227}$ polypeptide" is meant an analog wherein the arginine residues at positions 226 and 227 of SEQ ID NO: 8 have been deleted or replaced by other amino acids, for the purpose of eliminating a proteolytic processing site of the carboxy terminal pro-peptide of VEGF-C. Preferably, the VEGF-C $\Delta R_{226}\Delta R_{227}$ polypeptide comprises a portion of SEQ ID NO: 8 effective to permit binding of VEGFR-3. For example, the invention includes a VEGF-C $\Delta R_{226}\Delta R_{227}$ polypeptide having an amino acid sequence comprising amino acids 112-419 of SEQ ID NO: 8, wherein the arginine residues at positions 226 and 227 of SEQ ID NO: 8 have been deleted or replaced. Specifically exemplified herein is a VEGF-C $\Delta R_{226}\Delta R_{227}$ polypeptide wherein the arginine residues at positions 226 and 227 of SEQ ID NO: 8 have been replaced by serine residues.

Another family of VEGF-C analogs of the invention are human VEGF-C^{basic} polypeptides. By "VEGF-C^{basic} polypeptide" is meant a VEGF-C analog wherein at least one amino acid having a basic side chain has been introduced into the VEGF-C coding sequence, to emulate one or more basic residues in VEGF (e.g., residues Arg₁₀₈, Lys₁₁₀, and His₁₁₂ in the VEGF₁₆₅ precursor shown in Fig. 2) that have been implicated in VEGF receptor binding. Preferably, two or three basic residues are introduced into VEGF-C. Based on the VEGF/VEGF-C polypeptide alignment provided herein, positions 187, 189, and 191 of SEQ ID NO: 8 are preferred positions to introduce basic residues. For example, the invention includes a VEGF-C^{basic} polypeptide that is capable of binding to at least one of VEGFR-1, VEGFR-2, and VEGFR-3, and that has an amino acid sequence comprising residues 131 to 211 of SEQ ID NO: 8, wherein the glutamic acid residue at position 187, the threonine residue at position 189, and the proline residue at position 191

- 13 -

of SEQ ID NO: 8 have been replaced by an arginine residue, a lysine residue, and a histidine residue, respectively.

In yet another aspect of the invention, VEGF-C structural information is employed to create useful analogs of VEGF. For example, mature VEGF-C contains an 5 unpaired cysteine (position 137 of SEQ ID NO: 8) and is able to form non-covalently bonded polypeptide dimers. In one embodiment, a VEGF analog is created wherein this unpaired cysteine residue from mature VEGF-C is introduced at an analogous position of VEGF (e.g., introduced in place of Leu₅₈ of the human VEGF165 precursor (Fig. 2, Genbank Acc. No. M32977). Such VEGF analogs are termed VEGF^{+cys} polypeptides.

10 Thus, the invention includes a human VEGF analog wherein a cysteine residue is introduced in the VEGF amino acid sequence at a position selected from residues 53 to 63 of the human VEGF165 precursor having the amino acid sequence set forth in SEQ ID NO: 56. At least four naturally occurring VEGF isoforms have been described, and VEGF^{+cys} polypeptide analogs of each isoform are contemplated. Most preferably, the 15 cysteine is introduced at a position in a VEGF isoform which corresponds to position 58 of the VEGF165 precursor having the amino acid sequence set forth in SEQ ID NO: 56.

The present invention also provides purified and isolated polynucleotides (*i.e.*, nucleic acids) encoding all of the polypeptides of the invention, including but not limited to cDNAs and genomic DNAs encoding VEGF-C precursors, VEGF-C, and 20 biologically active fragments thereof, and DNAs encoding VEGF-C variants and VEGF-C analogs. A preferred nucleic acid of the invention comprises a DNA encoding amino acid residues 1 to 419 of SEQ ID NO: 8 or one of the aforementioned fragments or analogs thereof. Due to the degeneracy of the genetic code, numerous such coding sequences are possible, each having in common the coding of the amino acid sequence shown in SEQ ID 25 NO: 8 or the fragment or analog thereof. Distinct polynucleotides encoding any polypeptide of the invention by virtue of the degeneracy of the genetic code are within the scope of the invention.

A preferred polynucleotide according to the invention comprises the human VEGF-C cDNA sequence set forth in SEQ ID NO: 7 from nucleotide 352 to 1611. Other 30 polynucleotides according to the invention encode a VEGF-C polypeptide from, *e.g.*, mammals other than humans, birds (*e.g.*, avian quails), and others. Still other

polynucleotides of the invention comprise a coding sequence for a VEGF-C fragment, and allelic variants of those DNAs encoding part or all of VEGF-C.

Still other polynucleotides of the invention comprise a coding sequence for a VEGF-C variant or a VEGF-C analog. Preferred variant-encoding and analog-encoding polynucleotides comprise the human, mouse, or quail VEGF-C cDNA sequences disclosed herein (e.g., nucleotides 352-1611 of SEQ ID NO: 7 or continuous portions thereof) wherein one or more codon substitutions, deletions, or insertions have been introduced to create the variant/analog-encoding polynucleotide. For example, a preferred polynucleotide encoding a VEGF-C ΔC_{156} polypeptide comprises all or a portion of SEQ ID NO: 7 wherein the cysteine codon at positions 817-819 has been replaced by a codon encoding a different amino acid (e.g., a serine-encoding TCC codon).

The invention further comprises polynucleotides that hybridize to the aforementioned polynucleotides under standard stringent hybridization conditions. Exemplary stringent hybridization conditions are as follows: hybridization at 42°C in 50% formamide, 5X SSC, 20 mM Na₂PO₄, pH 6.8; and washing in 0.2X SSC at 55°C. It is understood by those of skill in the art that variation in these conditions occurs based on the length and GC nucleotide content of the sequences to be hybridized. Formulas standard in the art are appropriate for determining appropriate hybridization conditions. See Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (Second ed., Cold Spring Harbor Laboratory Press, 1989) §§ 9.47-9.51. These polynucleotides, capable of hybridizing to polynucleotides encoding VEGF-C, VEGF-C fragments, or VEGF-C analogs, are useful as nucleic acid probes for identifying, purifying and isolating polynucleotides encoding other (non-human) mammalian forms of VEGF-C and human VEGF-C allelic variants. Additionally, these polynucleotides are useful in screening methods of the invention, as described below.

Preferred nucleic acids useful as probes of the invention comprise nucleic acid sequences of at least about 16 continuous nucleotides of SEQ ID NO: 7. More preferably, these nucleic acid probes would have at least about 20 continuous nucleotides found in SEQ ID NO: 7. In using these nucleic acids as probes, it is preferred that the nucleic acids specifically hybridize to a portion of the sequence set forth in SEQ ID NO: 7. Specific hybridization is herein defined as hybridization under standard stringent hybridization conditions. To identify and isolate other mammalian VEGF-C genes

- 15 -

specifically, nucleic acid probes preferably are selected such that they fail to hybridize to genes related to VEGF-C (*e.g.*, fail to hybridize to human VEGF or to human VEGF-B genes).

Thus, the invention comprehends polynucleotides comprising at least about
5 16 nucleotides wherein the polynucleotides are capable of specifically hybridizing to a gene encoding VEGF-C, *e.g.*, a human gene. The specificity of hybridization ensures that a polynucleotide of the invention is able to hybridize to a nucleic acid encoding a VEGF-C under hybridization conditions that do not support hybridization of the polynucleotide to nucleic acids encoding, *e.g.*, VEGF or VEGF-B. In one embodiment, polynucleotides of
10 at least about 16 nucleotides, and preferably at least about 20 nucleotides, are selected as continuous nucleotide sequences found in SEQ ID NO: 7 or the complement of the nucleotide sequence set forth in SEQ ID NO: 7.

In another embodiment, the invention includes polynucleotides having at least 90 percent (preferably at least 95 percent, and more preferably at least 97, 98, or 99
15 percent) nucleotide sequence identity with a nucleotide sequence encoding a polypeptide of the invention. In a highly preferred embodiment, the polynucleotides have at least 95 percent sequence identity with a nucleotide sequence encoding a human VEGF-C precursor (such as the VEGF-C precursor in SEQ ID NO: 8 and allelic variants thereof), human VEGF-C, or biologically active VEGF-C fragments.

20 Additional aspects of the invention include vectors which comprise nucleic acids of the invention; and host cells transformed or transfected with nucleic acids or vectors of the invention. Preferred vectors of the invention are expression vectors wherein nucleic acids of the invention are operatively connected to appropriate promoters and other control sequences that regulate transcription and/or subsequent translation, such that
25 appropriate prokaryotic or eukaryotic host cells transformed or transfected with the vectors are capable of expressing the polypeptide encoded thereby (*e.g.*, the VEGF-C, VEGF-C fragment, VEGF-C variant, or VEGF-C analog encoded thereby). A preferred vector of the invention is plasmid pFLT4-L, having ATCC accession no. 97231. Such vectors and host cells are useful for recombinantly producing polypeptides of the
30 invention, including VEGF-C, and fragments, variants, and analogs thereof.

In a related aspect of the invention, host cells such as procaryotic and eukaryotic cells, especially unicellular host cells, are modified to express polypeptides of

- 16 -

the invention. Host cells may be stably transformed or transfected with isolated DNAs of the invention in a manner allowing expression of polypeptides of the invention therein.

Thus, the invention further includes a method of making polypeptides of the invention. In a preferred method, a nucleic acid or vector of the invention is expressed in a host cell, and
5 a polypeptide of the invention is purified from the host cell or the host cell's growth medium.

Similarly, the invention includes a method of making a polypeptide capable of specifically binding to VEGFR-1, VEGFR-2 and/or VEGFR-3, comprising the steps of:

(a) transforming or transfecting a host cell with a nucleic acid of the invention; (b)
10 cultivating the host cell to express the nucleic acid; and (c) purifying a polypeptide capable of specifically binding to VEGFR-1, VEGFR-2, and/or VEGFR-3 from the host cell or from the host cell's growth media. The invention also includes purified and isolated polypeptides produced by methods of the invention. In one preferred embodiment, the invention includes a human VEGF-C polypeptide or biologically active fragment, variant,
15 or analog thereof that is substantially free of other human polypeptides.

Alternatively, host cells may be modified by activating an endogenous VEGF-C gene that is not normally expressed in the host cells or that is expressed at a lower rate than is desired. Such host cells are modified (*e.g.*, by homologous recombination) to express the VEGF-C by replacing, in whole or in part, the naturally-
20 occurring VEGF-C promoter with part or all of a heterologous promoter so that the host cells express VEGF-C. In such host cells, the heterologous promoter DNA is operatively linked to the VEGF-C coding sequences, *i.e.*, controls transcription of the VEGF-C coding sequences. See, for example, PCT International Publication No. WO 94/12650; PCT International Publication No. WO 92/20808; and PCT International Publication No.
25 WO 91/09955. The invention also contemplates that, in addition to heterologous promoter DNA, amplifiable marker DNA (*e.g.*, *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamyl phosphate synthase, aspartate transcarbamylase, and dihydro-orotase) and/or intron DNA may be recombined along with the heterologous promoter DNA into the host cells. If linked to the VEGF-C coding sequences,
30 amplification of the marker DNA by standard selection methods results in co-amplification of the VEGF-C coding sequences in such host cells. Thus, the invention includes, for example, a cell comprising a nucleic acid having a sequence encoding human VEGF-C and

- 17 -

further comprising a non-VEGF-C promoter sequence (i.e., a heterologous promoter sequence) or other non-VEGF-C control sequence that increases RNA transcription in the cell of the sequence encoding human VEGF-C.

The DNA sequence information provided by the present invention also
5 makes possible the development, by homologous recombination or "knockout" strategies [see, Capecchi, *Science*, 244: 1288-1292 (1989)], of rodents that fail to express functional VEGF-C or that express a VEGF-C fragment, variant, or analog. Such rodents are useful as models for studying the activities of VEGF-C and VEGF-C modulators *in vivo*.

In another aspect, the invention includes an antibody that specifically binds
10 to one or more polypeptides of the invention, and/or binds to polypeptide multimers of the invention. In the context of antibodies of the invention, the term "specifically binds" is intended to exclude antibodies that cross-react with now-identified, related growth factors, such as VEGF, VEGF-B, PDGF-A, PDGF-B, FIGF, and PlGF. However, due to the high level of amino acid similarity shared by VEGF-C polypeptides of different species, it will
15 be understood that antibodies that specifically bind to human VEGF-C polypeptides of the invention will, in many instances, also bind non-human (e.g., mouse, quail) VEGF-C polypeptides of the invention. Antibodies, both monoclonal and polyclonal, may be made against a polypeptide of the invention according to standard techniques in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual* (Cold Spring Harbor
20 Laboratory Press, Cold Spring Harbor, New York (1988)). Standard protein manipulation techniques and recombinant techniques also may be employed to generate humanized antibodies and antigen-binding antibody fragments and other chimeric antibody polypeptides, all of which are considered antibodies of the invention. The invention further includes hybridoma cells that produce antibodies of the invention or other cell types
25 that have been genetically engineered to express antibody polypeptides of the invention. Antibodies of the invention may be used in diagnostic applications to monitor angiogenesis, vascularization, lymphatic vessels and their disease states, wound healing, or certain tumor cells, hematopoietic, or leukemia cells. The antibodies also may be used to block the ligand from activating its receptors; to purify polypeptides of the invention; and
30 to assay fluids for the presence of polypeptides of the invention. The invention further includes immunological assays (including radio-immuno assays, enzyme linked

- 18 -

immunosorbent assays, sandwich assays and the like) which employ antibodies of the invention.

Ligands according to the invention may be labeled with a detectable label and used to identify their corresponding receptors *in situ*. Labeled Flt4 ligand and anti-
5 Flt4 ligand antibodies may be used as imaging agents in the detection of lymphatic vessels, high endothelial venules and their disease states, and Flt4 receptors expressed in histochemical tissue sections. The ligand or antibody may be covalently or non-covalently coupled to a suitable supermagnetic, paramagnetic, electron dense, echogenic, or radioactive agent for imaging. Other, non-radioactive labels, such as biotin and avidin,
10 may also be used.

A related aspect of the invention is a method for the detection of specific cells, *e.g.*, endothelial cells. These cells may be found *in vivo*, or in *ex vivo* biological tissue samples. The method of detection comprises the steps of contacting a biological tissue comprising, *e.g.*, endothelial cells, with a polypeptide according to the invention
15 which is capable of binding to VEGFR-2 and/or VEGFR-3, under conditions wherein the polypeptide binds to the cells, optionally washing the biological tissue, and detecting the polypeptide bound to the cells in the biological tissue, thereby detecting the cells. It will be apparent that certain polypeptides of the invention are useful for detecting and/or imaging cells that express both VEGFR-2 and VEGFR-3, whereas other polypeptides
20 (*e.g.*, VEGF-C ΔC_{156} polypeptides) are useful for imaging specifically those cells which express VEGFR-3.

The many biological activities described herein for VEGF-C (including but not limited to affecting growth and migration of vascular endothelial cells; promoting growth of lymphatic endothelial cells and lymphatic vessels; increasing vascular
25 permeability; and affecting myelopoiesis (*e.g.*, growth of neutrophilic granulocytes)) support numerous diagnostic and *in vitro* and *in vivo* clinical utilities for polypeptides and antibodies of the invention, for modulating (stimulating or inhibiting) these biological activities. Generally, VEGF-C and precursor, fragment, variant, and analog polypeptides that retain one or more VEGF-C biological activities are useful agonists for stimulating the
30 desired biological activity; whereas precursor, fragment, variant, and analog polypeptides that are capable of binding to VEGFR-2 and/or VEGFR-3 (either alone or as a homo- or hetero-dimer with other polypeptides) without stimulating receptor-mediated VEGF-C

- 19 -

activity (i.e., without activating the receptor) are useful as antagonists (inhibitors) of VEGF-C. Similarly, antibodies of the invention that bind biologically active VEGF-C forms and thereby interfere with VEGF-C-receptor interactions are useful as inhibitors of VEGF-C. Antisense oligonucleotides comprising a portion of the VEGF-C coding
5 sequence and/or its complement also are contemplated as inhibitors of the invention. Both biologically active polypeptides and inhibitor polypeptides of the invention have utilities in various imaging applications.

For example, the biological effects of VEGF-C on vascular endothelial cells indicate *in vivo* uses for polypeptides of the invention for stimulating angiogenesis (e.g.,
10 during wound healing, in tissue transplantation, in eye diseases, in the formation of collateral vessels around arterial stenoses and into injured tissues after infarction) and for inhibiting angiogenesis (e.g., to inhibit tumor growth and/or metastatic cancer). The biological effects on vascular endothelial cells indicate *in vitro* uses for biologically active forms of VEGF-C to promote the growth of (including proliferation of) cultured vascular
15 endothelial cells and precursors thereof.

The biological effects of VEGF-C on lymphatic endothelia indicate *in vivo* uses for polypeptides of the invention for stimulating lymphangiogenesis (e.g., to promote re-growth or permeability of lymphatic vessels in, for example, organ transplant patients; to mitigate the loss of axillary lymphatic vessels following surgical interventions in the
20 treatment of cancer (e.g., breast cancer); to treat aplasia of the lymphatic vessels or lymphatic obstructions) and for inhibiting it (e.g., to treat lymphangiomas). Additional *in vivo* uses for polypeptides of the invention include the treatment or prevention of inflammation, edema, elephantiasis, and Milroy's disease. The biological effects on lymphatic endothelial cells indicate *in vitro* uses for biologically active forms of VEGF-C
25 to promote the growth of cultured lymphatic endothelial cells and precursors thereof.

Thus, the invention includes a method of modulating (stimulating/increasing or inhibiting/decreasing) the growth of vertebrate endothelial cells or vertebrate endothelial precursor cells comprising contacting such endothelial cells or precursor cells with a polypeptide or antibody (or antigen-binding portion thereof) of the invention, in an amount
30 effective to modulate the growth of the endothelial or endothelial precursor cells.

Mammalian endothelial cells and their precursors are preferred. Human endothelial cells are highly preferred. In one embodiment, the endothelial cells are lymphatic endothelial

- 20 -

cells. In another embodiment, the cells are vascular endothelial cells. The method may be an *in vitro* method (e.g., for cultured endothelial cells) or an *in vivo* method. The *in vitro* growth modulation of CD34+ endothelial precursor cells [see, e.g., Asahara *et al.*, *Science*, 275:964-967 (1997)] isolated from peripheral blood, bone marrow, or cord blood is specifically contemplated. For *in vivo* methods, it is highly preferable to administer a pharmaceutical composition (comprising the polypeptide formulated in a pharmaceutically acceptable diluent, adjuvant, excipient, carrier, or the like) to the subject, in an amount effective to modulate the growth of lymphatic endothelial cells *in vivo*.

In one preferred embodiment, the endothelial cells are lymphatic endothelial cells, and the polypeptide is one that has reduced effect on the permeability of mammalian blood vessels compared to a wildtype VEGF-C polypeptide (e.g., compared with VEGF-C having an amino acid sequence set forth in SEQ ID NO: 8 from residue 103 to residue 227). VEGF-C ΔC_{156} polypeptides are contemplated for use in this embodiment.

In modulating the growth of endothelial cells *in vivo*, the invention contemplates the modulation of endothelial cell-related disorders. Endothelial cell disorders contemplated by the invention include, but are not limited to, physical loss of lymphatic vessels (e.g., surgical removal of axillary lymph tissue), lymphatic vessel occlusion (e.g., elephantiasis), and lymphangiomas. In a preferred embodiment, the subject, and endothelial cells, are human. The endothelial cells may be provided *in vitro* or *in vivo*, and they may be contained in a tissue graft. An effective amount of a polypeptide is defined herein as that amount of polypeptide empirically determined to be necessary to achieve a reproducible change in cell growth rate (as determined by microscopic or macroscopic visualization and estimation of cell doubling time, or nucleic acid synthesis assays), as would be understood by one of ordinary skill in the art.

Polypeptides of the invention may be used to stimulate lymphocyte production and maturation, and to promote or inhibit trafficking of leukocytes between tissues and lymphatic vessels or to affect migration in and out of the thymus.

The biological effects of VEGF-C on myelopoiesis indicate *in vivo* and *in vitro* uses for polypeptides of the invention for stimulating myelopoiesis (especially growth of neutrophilic granulocytes) or inhibiting it. Thus, the invention includes a method for modulating myelopoiesis in a mammalian subject comprising administering to a mammalian subject in need of modulation of myelopoiesis an amount of a polypeptide or antibody (or

- 21 -

antigen-binding portion thereof) of the invention that is effective to modulate myelopoiesis. In one embodiment, a mammalian subject suffering from granulocytopenia is selected, and the method comprises administering to the subject an amount of a polypeptide effective to stimulate myelopoiesis. In particular, a polypeptide of the invention is administered in an amount effective to increase the neutrophil count in blood of the subject. Preferred subjects are human subjects. An effective amount of a polypeptide is an amount of polypeptide empirically determined to be necessary to achieve a reproducible change in the production of neutrophilic granulocytes (as determined by microscopic or macroscopic visualization and estimation of cell doubling time, or nucleic acid synthesis assays), as would be understood by one of ordinary skill in the art.

In a related embodiment, the invention includes a method of increasing the number of neutrophils in the blood of a mammalian subject comprising the step of expressing in a cell in a subject in need of an increased number of blood neutrophils a DNA encoding a VEGF-C protein, the DNA operatively linked to a non-VEGF-C promoter or other non-VEGF-C control sequence that promotes expression of the DNA in the cell.

Similarly, the invention includes a method of modulating the growth of neutrophilic granulocytes *in vitro* or *in vivo* comprising the step of contacting mammalian stem cells with a polypeptide or antibody of the invention in an amount effective to modulate the growth of mammalian endothelial cells.

More generally, the invention includes a method for modulating the growth of CD34+ progenitor cells (especially hematopoietic progenitor cells and endothelial progenitor cells) *in vitro* or *in vivo* comprising the step of contacting mammalian CD34+ progenitor cells with a polypeptide or antibody of the invention in an amount effective to modulate the growth of mammalian endothelial cells. For *in vitro* methods, CD34+ progenitor cells isolated from cord blood or bone marrow are specifically contemplated.

It will be apparent from the Detailed Description below that *in vitro* and *in vivo* methods of the invention for stimulating the growth of CD34+ precursor cells also include methods wherein polypeptides of the invention are employed together (simultaneously or sequentially) with other polypeptide factors for the purpose of modulating hematopoiesis/myelopoiesis or endothelial cell proliferation. Such other factors include, but are not limited to colony stimulating factors ("CSFs," e.g.,

granulocyte-CSF (G-CSF), macrophage-CSF (M-CSF), and granulocyte-macrophage-CSF (GM-CSF)), interleukin-3 (IL-3, also called multi-colony stimulating factor), other interleukins, stem cell factor (SCF), other polypeptide factors, such as VEGF, and their analogs that have been described and are known in the art. See generally *The Cytokine Handbook, Second Ed.*, Angus Thomson (editor), Academic Press (1996); Callard and Gearing, *The Cytokine FactsBook*, Academic Press Inc. (1994); and Cowling and Dexter, *TIBTECH*, 10(10):349-357 (1992). The use of a polypeptide of the invention as a progenitor cell or myelopoietic cell growth factor or co-factor with one or more of the foregoing factors may potentiate previously unattainable myelopoietic effects and/or

10 potentiate previously attainable myelopoietic effects while using less of the foregoing factors than would be necessary in the absence of a polypeptide of the invention.

In addition to methods, the invention includes compositions comprising polypeptides of the invention in admixture with one or more of the factors identified in the previous paragraph. Preferred compositions further comprise a pharmaceutically

15 acceptable diluent, adjuvant, excipient, or carrier. The invention also includes kits comprising (a) at least one polypeptide of the invention packaged with (b) one or more of the foregoing polypeptides (e.g., in unit dosage form, but not in admixture with each other).

For methods which involve the *in vivo* administration of polypeptides or

20 antibodies of the invention, it is contemplated that the polypeptides or antibodies will be administered in any suitable manner using an appropriate pharmaceutically-acceptable vehicle, e.g., a pharmaceutically-acceptable diluent, adjuvant, excipient or carrier. Thus, the invention further includes compositions, e.g., pharmaceutical compositions, comprising one or more polypeptides or antibodies of the invention. By pharmaceutical composition

25 is meant a composition that may be administered to a mammalian host, e.g., orally, topically, parenterally (including subcutaneous injections, intravenous, intramuscular, intracisternal injection or infusion techniques), by inhalation spray, or rectally, in unit dosage formulations containing conventional non-toxic carriers, diluents (e.g., calcium carbonate, sodium carbonate, lactose, calcium phosphate, sodium phosphate, kaolin,

30 water), adjuvants, vehicles, and the like, including but not limited to flavoring agents, preserving agents; granulating and disintegrating agents; binding agents; time delay

- 23 -

materials; oils; suspending agents; dispersing or wetting agents; anti-oxidants; emulsifiers, etc.

The invention further provides a method of using a polypeptide of the invention for the manufacture of a medicament for use in any of the foregoing methods.

5 Similarly, the invention further provides a method of using a polypeptide of the invention for the manufacture of a medicament for the treatment of any of the foregoing indicated conditions and disease states. Such methods optionally involve the use of additional biologically active ingredients (e.g., VEGF, PlGF, G-CSF, etc.) for the manufacture of the medicament.

10 Effective amounts of polypeptides for the foregoing methods are empirically determined using standard *in vitro* and *in vivo* dose-response assays. In addition, experimental data provided herein provide guidance as to amounts of polypeptides of the invention that are effective for achieving a desired biological response. For example, the dissociation constants determined for one form of mature VEGF-C
15 ($K_D=135$ pM for VEGFR-3 and $K_D=410$ pM for VEGFR-2) provide an indication as to the concentration of VEGF-C necessary to achieve biological effects, because such dissociation constants represent concentrations at which half of the VEGF-C polypeptide is bound to the receptors through which VEGF-C biological effects are mediated. Results from *in vivo* Miles assays, wherein 0 - 8 picomoles of VEGF-C was injected intradermally,
20 provide an indication that picomole quantities of mature VEGF-C are sufficient to induce localized biological effects. *In vitro* analysis of ^3H -thymidine incorporation into bovine capillary endothelial cells treated with a mature VEGF-C form showed increasing VEGF-C effects on cell proliferation at concentrations of 10 - 1000 pM. Collectively, this data suggests that localized concentrations of 100 - 1000 pM of fully-processed VEGF-C have
25 VEGF-C biological activity *in vivo*. Effective concentrations of other polypeptides of the invention are generally expected to correlate with the dissociation constant of the polypeptides for the relevant receptors. Pharmacokinetic and pharmacological analyses reveals the preferred dosages, dosage formulations, and methods of administration to achieve the desired local or systemic concentration of a polypeptide of the invention.

30 Polypeptides of the invention also may be used to quantify future metastatic risk by assaying biopsy material for the presence of active receptors or ligands in a binding assay. Such a binding assay may involve the use of a detectably labeled polypeptide of the

invention or of an unlabeled polypeptide in conjunction with a labeled antibody, for example. Kits comprising such substances are included within the scope of the invention.

The present invention also provides methods for using the claimed nucleic acids (*i.e.*, polynucleotides) in screening for endothelial cell disorders. In a preferred
5 embodiment, the invention provides a method for screening an endothelial cell disorder in a mammalian subject comprising the steps of providing a sample of endothelial cell nucleic acids from the subject, contacting the sample of endothelial cell nucleic acids with a polynucleotide of the invention which is capable of hybridizing to a gene encoding VEGF-C (and preferably capable of hybridizing to VEGF-C mRNA), determining the level of
10 hybridization between the endothelial cell nucleic acids and the polynucleotide, and correlating the level of hybridization with a disorder. A preferred mammalian subject, and source of endothelial cell nucleic acids, is a human. The disorders contemplated by the method of screening with polynucleotides include, but are not limited to, vessel disorders such as the aforementioned lymphatic vessel disorders, and hypoxia.

15 Purified and isolated polynucleotides encoding other (non-human) VEGF-C forms also are aspects of the invention, as are the polypeptides encoded thereby, and antibodies that bind to non-human VEGF-C forms. Preferred non-human forms of VEGF-C are forms derived from other vertebrate species, including avian and mammalian species. Mammalian forms are highly preferred. Thus, the invention includes a purified and
20 isolated mammalian VEGF-C polypeptide, and also a purified and isolated polynucleotide encoding such a polypeptide.

In one embodiment, the invention includes a purified and isolated polypeptide having the amino acid sequence of residues 1 to 415 of SEQ ID NO: 11, which sequence corresponds to a putative mouse VEGF-C precursor. The putative mouse
25 VEGF-C precursor is believed to be processed into a mature mouse VEGF-C in a manner analogous to the processing of the human prepro-polypeptide. Thus, in a related aspect, the invention includes a purified and isolated polypeptide capable of binding with high affinity to an Flt4 receptor tyrosine kinase (*e.g.*, a human or mouse Flt-4 receptor tyrosine kinase), the polypeptide comprising a fragment of the purified and isolated polypeptide
30 having the amino acid sequence of residues 1 to 415 of SEQ ID NO: 11, the fragment being capable of binding with high affinity to the Flt4 receptor tyrosine kinase. The invention further includes multimers of the foregoing polypeptides and purified and

- 25 -

isolated nucleic acids encoding the foregoing polypeptides, such as a nucleic acid comprising all or a portion of the sequence shown in SEQ ID NO: 10.

In another embodiment, the invention includes a purified and isolated quail VEGF-C polypeptide, biologically active fragments and multimers thereof, and 5 polynucleotides encoding the foregoing polypeptides.

It is also contemplated that VEGF-C polypeptides from other species may be altered in the manner described herein with respect to human VEGF-C variants, in order to alter biological properties of the wildtype protein. For example, elimination of the cysteine at position 152 of SEQ ID NO: 11 or position 155 of SEQ ID NO: 13 is expected 10 to alter VEGFR-2 binding properties in the manner described below for human VEGF-C ΔC_{156} mutants.

In yet another embodiment, the invention includes a DNA comprising a VEGF-C promoter, that is capable of promoting expression of a VEGF-C gene or another operatively-linked, protein-encoding gene in native host cells, under conditions wherein 15 VEGF-C is expressed in such cells. Thus, the invention includes a purified nucleic acid comprising a VEGF-C promoter sequence. Genomic clone lambda 5 described herein comprises more than 5 kb of human genomic DNA upstream of the VEGF-C translation initiation codon, and contains promoter DNA of the invention. Approximately 2.4 kb of this upstream sequence is set forth in SEQ ID NO: 48. Thus, in one embodiment, the 20 invention includes a purified nucleic acid comprising a portion of SEQ ID NO: 48, wherein the portion is capable of promoting expression of a protein encoding gene operatively linked thereto under conditions wherein VEGF-C is expressed in native host cells. Similarly, the invention includes a chimeric nucleic acid comprising a VEGF-C promoter nucleic acid according to the invention operatively connected to a sequence encoding a 25 protein other than a human VEGF-C.

Additional aspects and embodiments of the invention will be apparent from the detailed description which follows.

BRIEF DESCRIPTION OF THE DRAWING

Figure 1 schematically depicts major endothelial cell receptor tyrosine 30 kinases and growth factors involved in vasculogenesis and angiogenesis. Major structural domains are depicted, including immunoglobulin-like domains (IGH), epidermal growth

- 26 -

factor homology domains (EGFH), fibronectin type III domains (FNIII), transmembrane (TM) and juxtamembrane (JM) domains, tyrosine kinase (TK1, TK2) domains, kinase insert domains (KI), and carboxy-terminal domains (CT).

Figure 2 shows a comparison of the deduced amino acid sequences of
 5 PDGF-A (SEQ ID NO: 53), PDGF-B (SEQ ID NO: 54), PIGF-1 (SEQ ID NO: 55),
 VEGF-B₁₆₇ (SEQ ID NO: 56), VEGF165 (SEQ ID NO: 57), and Flt4 ligand (VEGF-C,
 (SEQ ID NO: 8)).

Figure 3 schematically depicts the VEGF-C promoter-reporter constructs
 and their activities in transfected HeLa cells. A restriction map of a portion of a genomic
 10 clone that includes the VEGF-C initiation codon and about 6 kb of upstream sequence is
 depicted above the constructs. Constructs were made linking putative VEGF-C promoter
 to the Luciferase reporter gene in pGL3 vector (Promega) and introduced into HeLa cells
 by calcium phosphate-mediated transfection method. The Luciferase activity obtained was
 compared to the level using the promoterless pGL3basic construct to obtain a measure of
 15 relative promoter activity. Luciferase activity is expressed graphically as a ratio of activity
 of the constructs versus this control. Also shown are numerical ratios of Luciferase
 activity in experiments where the constructs were transfected into HeLa cells and cells
 were starved 24 hours followed by serum stimulation for four hours (Luciferase activity is
 expressed as a ratio of activity in serum-stimulated versus serum-starved cells).

20 Figure 4 graphically depicts the results of a competitive binding assay. The
 ability of VEGF165 (filled triangles: ▼), wildtype VEGF-C (filled circles: ●), and three
 VEGF-C mutants [VEGF-C R226,227S (open boxes: □); VEGF-C ΔNΔCHis (open
 circles: ○); and VEGF-C ΔNΔCHisC156S (open triangles: Δ)] to compete with ¹²⁵I-
 VEGF-CΔNΔCHis for binding to VEGFR-2 and VEGFR-3 is shown.

25 Figure 5 depicts the amino acid sequences of human (SEQ ID NO: 8),
 murine (SEQ ID NO: 11), and quail (SEQ ID NO: 13) VEGF-C polypeptides, aligned to
 show similarity. Residues conserved in all three species are depicted in bold.

Figures 6A-C depict electrophoretic fractionations of the various forms of
 recombinant VEGF-C produced by transfected 293 EBNA cells. Figure 6B depicts the
 30 electrophoretic fractionation, under non-reducing conditions, of polypeptides produced
 from mock (M) transfected cells, cells transfected with wild type (wt) VEGF-C cDNA,
 and cells transfected with a cDNA encoding the VEGF-C mutant VEGF-C-R102S. Each

of the bands identified in Figure 6B was excised and electrophoretically fractionated in a separate lane under reducing conditions. Fractionation of bands corresponding to wt VEGF-C are depicted in Figure 6A; fractionation of bands corresponding to the R102S mutant are depicted in Figure 6C.

5 Figures 7A-B depict the forms and sizes of wild type and mutant recombinant VEGF-Cs, as revealed by non-reducing gel electrophoresis. Figure 7A shows the VEGF-C forms secreted into the media; Figure 7B shows the VEGF-C forms retained by the cells. Mock (M) transfected cells served as a control.

 Figures 8A-B present a comparison of the pattern of immunoprecipitated, 10 labeled VEGF-C forms using antisera 882 and antisera 905. Adjacent lanes contain immunoprecipitates that were (lanes marked +) or were not (lanes marked -) subjected to reduction and alkylation.

 Fig. 9 is a schematic model of the proteolytic processing of VEGF-C. The regions of the VEGF-C polypeptide are depicted as follows: signal sequence = dark 15 shaded box; VEGF-homology domain = medium shaded box; N-terminal and C-terminal propeptides = dotted and open boxes, respectively. Conserved cysteine residues in the VEGF-homology domain are depicted with dots (for clarity, cysteine residues in the C-terminal propeptide are not marked). Putative sites of N-linked glycosylation are shown with Y symbols. Numbers indicate approximate molecular mass (kDa) of the 20 corresponding polypeptide as measured by SDS-PAGE in reducing conditions. Disulfide bonds are marked as -S-S-; non-covalent bonds are depicted as dotted lines. A question mark indicates the presence of a possible non-covalent bond. The proteolytic generation of a small fraction of disulfide-linked 21 kDa forms is not indicated in the figure. Several intermediate forms also are omitted to simplify the scheme. Particularly, only one 25 precursor polypeptide is cleaved initially. The figure is not intended to suggest that other intermediate forms, for example 21 kDa + 31 kDa, 31 kDa + 31 kDa + 29 kDa, do not exist.

 Figure 10 presents a comparison of the human and mouse VEGF-C amino acid sequences. The amino acid sequence of mouse VEGF-C is presented on the top line 30 and differences in the human sequence are marked below it. An arrow indicates the putative cleavage site for the signal peptidase; BR3P motifs, as well as a CR/SC motif, are

boxed; and conserved cysteine residues are marked in bold above the sequence. Arginine residue 158 is also marked in bold. The numbering refers to mouse VEGF-C residues.

Figures 11A and 11B depict the genomic structure of the human (11A) and murine (11B) VEGF-C genes. Sequences of exon-intron junctions are depicted together with exon and intron lengths. Intron sequences are depicted in lower case letters.

Nucleotides of the open reading frame observed in VEGF-C cDNAs are indicated as upper case letters in triplets (corresponding to the codons encoded at the junctions).

Figure 12 depicts the exon-intron organization of the human VEGF-C gene. Seven exons are depicted as open boxes, with exon size depicted in base pairs. Introns are depicted as lines, with intron size (base pairs) depicted above the lines. 5' and 3' untranslated sequences of a putative 2.4 kb mature mRNA are depicted as shaded boxes. The location of genomic clones used to characterize the VEGF-C gene are depicted below the map of the gene.

DETAILED DESCRIPTION OF THE INVENTION

Described herein is the isolation of a novel vascular endothelial growth factor and the cloning of a DNA encoding this novel growth factor from a cDNA library prepared from the human prostatic adenocarcinoma cell line PC-3. The isolated cDNA encodes a protein which is proteolytically processed and secreted to cell culture medium. The processing is described in detail below. The secreted protein, designated VEGF-C, binds to the extracellular domain and induces tyrosine autophosphorylation of both Flt4 (VEGFR-3) and KDR/flk-1 (VEGFR-2). In contrast, neither VEGF nor PlGF show high affinity binding to VEGFR-3 or induced its autophosphorylation. VEGF-C also stimulates the migration of endothelial cells in collagen gel and induces vascular permeability *in vivo*. In transgenic mice, VEGF-C induces proliferation of the lymphatic endothelium and causes an increase in neutrophilic granulocytes. Based on studies of VEGF-C variants and analogs and studies of VEGF precursors, it is anticipated that one or more VEGF-C precursors (the largest putative native VEGF-C precursor, excluding signal peptide, having the complete amino acid sequence from residue 32 to residue 419 of SEQ ID NO: 8) is capable of stimulating VEGFR-3.

In addition to providing a cDNA sequence encoding prepro-VEGF-C, the present application also provides significant guidance concerning portions of the VEGF-C

amino acid sequence which are necessary for biological activity and portions (of one or more amino acids) which, when altered, will modulate (up-regulate or inhibit) VEGF-C biological activities. Such alterations are readily achieved through recombinant DNA and protein techniques, such as site-directed mutagenesis of a VEGF-C encoding cDNA and
5 recombinant expression of the resultant modified cDNA. The skilled artisan also understands that, in recombinant production of proteins, additional sequence may be expressed along with a sequence encoding a polypeptide having a desired biological activity, while retaining a desired biological activity of the protein. For example, additional amino acids may be added at the amino terminus, at the carboxy-terminus, or as an
10 insertion into the polypeptide sequence. Similarly, deletion variants of a protein with a desired biological activity can be recombinantly expressed that lack certain residues of the endogenous/natural protein, while retaining a desired biological activity. Moreover, it is well-known that recombinant protein variants may be produced having conservative amino acid replacements (including but not limited to substitution of one or more amino acids for
15 other amino acids having similar chemical side-chains (acidic, basic, aliphatic, aliphatic hydroxyl, aromatic, amide, etc.)) which do not eliminate the desired biological activity of the protein. Accordingly, it is anticipated that such alterations of VEGF-C are VEGF-C equivalents within the scope of the invention.

As set forth in greater detail below, the putative prepro-VEGF-C has a
20 deduced molecular mass of 46,883; a putative prepro-VEGF-C processing intermediate has an observed molecular weight of about 32 kD; and mature VEGF-C isolated from conditioned media has a molecular weight of about 23 kD as assessed by SDS-PAGE under reducing conditions. A major part of the difference in the observed molecular mass of the purified and recombinant VEGF-C and the deduced molecular mass of the prepro-
25 VEGF-C encoded by the VEGF-C open reading frame (ORF) is attributable to proteolytic removal of sequences at the amino-terminal and carboxyl-terminal regions of the prepro-VEGF-C polypeptide. Extrapolation from studies of the structure of PDGF (Heldin *et al.*, *Growth Factors*, 8:245-52 (1993)) suggests that the region critical for receptor binding and activation by VEGF-C is contained within amino acids residues 104-213, which are
30 found in the secreted form of the VEGF-C protein (*i.e.*, the form lacking the putative prepro leader sequence and some carboxyterminal sequences). The 23 kD polypeptide binding VEGFR-3 corresponds to a VEGF-homologous domain of VEGF-C. After

biosynthesis, the nascent VEGF-C polypeptide may be glycosylated at three putative N-linked glycosylation sites identified in the deduced VEGF-C amino acid sequence.

Polypeptides containing modifications, such as N-linked glycosylations, are intended as aspects of the invention.

5 The carboxyl terminal amino acid sequences, which increase the length of the VEGF-C polypeptide in comparison with other ligands of this family, show a pattern of spacing of cysteine residues reminiscent of the Balbiani ring 3 protein (BR3P) sequence (Dignam *et al.*, *Gene*, 88:133-40 (1990); Paulsson *et al.*, *J. Mol. Biol.*, 211:331-49 (1990)). This novel C-terminal silk protein-like structural motif of VEGF-C may fold into
10 an independent domain, which is cleaved off after biosynthesis. Interestingly, at least one cysteine motif of the BR3P type is also found in the carboxyl terminus of VEGF. As explained in detail below, putative precursors and putative fully-processed VEGF-C were both detected in the cell culture media, suggesting cleavage by cellular proteases. The determination of amino-terminal and carboxy-terminal sequences of VEGF-C isolates was
15 performed to identify the proteolytic processing sites. Antibodies generated against different parts of the pro-VEGF-C molecule were used to determine the precursor-product relationship and ratio, their cellular distribution, and the kinetics of processing and secretion.

 VEGF-C has a conserved pattern of eight cysteine residues, which may
20 participate in the formation of intra- and interchain disulfide bonds, creating an antiparallel, dimeric, biologically active molecule, similar to PDGF. Mutational analysis of the cysteine residues involved in the interchain disulfide bridges has shown that, in contrast to PDGF, VEGF dimers need to be held together by these covalent interactions in order to maintain biological activity. Disulfide linking of the VEGF-C polypeptide chains was evident in the
25 analysis of VEGF-C in nonreducing conditions, although recombinant protein also contained "fully processed" ligand-active VEGF-C forms which lacked disulfide bonds between the polypeptides. (See Fig. 9.)

 VEGFR-3, which distinguishes between VEGF and VEGF-C, is closely related in structure to VEGFR-1 and VEGFR-2. Finnerty *et al.*, *Oncogene*, 8:2293-98
30 (1993); Galland *et al.*, *Oncogene*, 8:1233-40 (1993); Pajusola *et al.*, *Cancer Res.*, 52:5738-43 (1992). Besides VEGFR-3, VEGFR-2 tyrosine kinase also is activated in response to VEGF-C. VEGFR-2 mediated signals cause striking changes in the

morphology, actin reorganization and membrane ruffling of porcine aortic endothelial cells over-expressing this receptor. In these cells, VEGFR-2 also mediated ligand-induced chemotaxis and mitogenicity. Waltenberger *et al.*, *J. Biol. Chem.*, 269:26988-95 (1994). Similarly, the receptor chimera CSF-1R/VEGFR-3 was mitogenic when ectopically
5 expressed in NIH 3T3 fibroblastic cells, but not in porcine aortic endothelial cells (Pajusola *et al.*, 1994). Consistent with such results, the bovine capillary endothelial (BCE) cells, which express VEGFR-2 mRNA but very little or no VEGFR-1 or VEGFR-3 mRNAs, showed enhanced migration when stimulated with VEGF-C. Light microscopy of the BCE cell cultures in collagen gel also suggested that VEGF-C stimulated the proliferation
10 of these cells. The data thus indicate that the VEGF family of ligands and receptors show a great specificity in their signaling, which may be cell-type-dependent.

The expression pattern of the VEGFR-3 (Kaipainen *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 92:3566-70 (1995)) suggests that VEGF-C may function in the formation of the venous and lymphatic vascular systems during embryogenesis.
15 Constitutive expression of VEGF-C in adult tissues shown herein further suggests that this gene product also is involved in the maintenance of the differentiated functions of the lymphatic and certain venous endothelia where VEGFR-3 is expressed (Kaipainen *et al.*, 1995). Lymphatic capillaries do not have well-formed basal laminae and an interesting possibility exists that the silk-like BR3P motif is involved in producing a supramolecular
20 structure which could regulate the availability of VEGF-C in tissues. However, as shown here, VEGF-C also activates VEGFR-2, which is abundant in proliferating endothelial cells of vascular sprouts and branching vessels of embryonic tissues, but not so abundant in adult tissues. Millauer *et al.*, *Nature*, 367:576-78 (1993). These data have suggested that VEGFR-2 is a major regulator of vasculogenesis and angiogenesis. VEGF-C may thus
25 have a unique effect on lymphatic endothelium and a more redundant function, shared with VEGF, in angiogenesis and possibly in regulating the permeability of several types of endothelia. Because VEGF-C stimulates VEGFR-2 and promotes endothelial migration, VEGF-C may be useful as an inducer of angiogenesis of blood and lymphatic vessels in wound healing, in tissue transplantation, in eye diseases, and in the formation of collateral
30 vessels around arterial stenoses and into injured tissues after infarction.

Previously-identified growth factors that promote angiogenesis include the fibroblast growth factors, hepatocyte growth factor/scatter factor, PDGF and TGF- α .

- 32 -

(See e.g., Folkman, *Nature Med.*, 1:27-31 (1995); Friesel *et al.*, *FASEB J.*, 9:919-25 (1995); Mustonen *et al.*, *J. Cell. Biol.*, 129:895-98 (1995). However, VEGF has been the only growth factor relatively specific for endothelial cells. The newly identified factors VEGF-B [Olofsson *et al.*, *Proc. Natl. Acad. Sci.*, 93:2578-81 (1996)] and VEGF-C thus
5 increase our understanding of the complexity of the specific and redundant positive signals for endothelial cells involved in vasculogenesis, angiogenesis, permeability, and perhaps also other endothelial functions. Expression studies using Northern blotting show abundant VEGF-C expression in heart and skeletal muscle; other tissues, such as placenta, ovary, small intestine, thyroid gland, kidney, prostate, spleen, testis and large intestine also
10 express this gene. Whereas PlGF is predominantly expressed in the placenta, the expression patterns of VEGF, VEGF-B and VEGF-C overlap in many tissues, which suggests that members of the VEGF family may form heterodimers and interact to exert their physiological functions.

Targeted mutagenesis leading to inactivation of the VEGF receptor loci in
15 the mouse genome has shown that VEGFR-1 is necessary for the proper organization of endothelial cells forming the vascular endothelium, while VEGFR-2 is necessary for the generation of both endothelial and hematopoietic cells. This suggests that the four genes of the VEGF family can be targets for mutations leading to vascular malformations or cardiovascular diseases.

20 The following Examples illustrate preferred embodiments of the invention, wherein the isolation, characterization, and function of VEGF-C, VEGF-C variants and analogs, VEGF-C-encoding nucleic acids, and anti-VEGF-C antibodies according to the invention are shown.

Example 1

25 Production of pLTRFlt4l expression vector

The identification and isolation of two forms of Flt4 receptor tyrosine kinase (VEGFR-3) cDNA (Flt4 short form (Flt4s), Genbank Accession No. X68203, SEQ ID NO: 1; and Flt4 long form, (Flt4l), Genbank Accession Nos. X68203 and S66407, SEQ ID NO: 2) was reported in United States Patent Application Serial Number
30 08/340,011, filed November 14, 1994, incorporated by reference herein. An Flt4 expression vector designated pLTRFlt4l (encoding the long form of Flt4) was constructed

using the pLTRpoly expression vector reported in Mäkelä *et al.*, *Gene*, 118: 293-294 (1992) (Genbank accession number X60280, SEQ ID NO: 3) and the Flt4 cDNAs, in the manner described in commonly-owned PCT patent application PCT/FI96/00427, filed August 01, 1996, published as PCT publication No. WO 97/05250 on 13 February 1997, 5 and commonly-owned United States Patent Application Serial Nos. 08/671,573, filed June 28, 1996; 08/601,132, filed February 14, 1996; 08/585,895, filed January 12, 1996; and 08/510,133, filed August 1, 1995, all of which are incorporated by reference in their entirety.

EXAMPLE 2

10 Production and analysis of Flt4I transfected cells

NIH 3T3 cells (60 % confluent) were co-transfected with 5 micrograms of the pLTRFlt4I construct and 0.25 micrograms of the pSV2neo vector containing the neomycin phosphotransferase gene (Southern *et al.*, *J. Mol. Appl. Genet.*, 1:327 (1982)), using the DOTAP liposome-based transfection reagents (Boehringer-Mannheim, 15 Mannheim, Germany). One day after transfection, the cells were transferred into selection media containing 0.5 mg/ml geneticin (GIBCO, Grand Island, N.Y.). Colonies of geneticin-resistant cells were isolated and analyzed for expression of the Flt4 proteins. Cells were lysed in boiling lysis buffer containing 3.3% SDS and 125 mM Tris, pH 6.8. Protein concentrations of the samples were measured by the BCA method (Pierce, 20 Rockford, IL). About 50 micrograms of protein from each lysate were analyzed for the presence of Flt4 by 6% SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and immunoblotting using antisera against the carboxyl terminus of Flt4. Signals on Western blots were revealed using the ECL method (Amersham).

For production of anti-Flt4 antiserum, the Flt4 cDNA fragment encoding 25 the 40 carboxy-terminal amino acid residues of the Flt4 short form: NH₂-PMTPTTYKG SVDNQTDSGM VLASEEFEQI ESRHRQESGFR-COOH (SEQ ID NO: 4) was cloned as a 657 bp *EcoRI*-fragment into the pGEX-1λT bacterial expression vector (Pharmacia-LKB, Inc., Uppsala, Sweden) in frame with the glutathione-S-transferase coding region. The resultant GST-Flt4S fusion protein was produced in *E. coli* and purified by affinity 30 chromatography using a glutathione-Sepharose 4B column. The purified protein was lyophilized, dissolved in phosphate-buffered saline (PBS), mixed with Freund's adjuvant

- 34 -

and used for immunization of rabbits at bi-weekly intervals using methods standard in the art (Harlow *et al.*, *Antibodies: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1988)). Antisera were used, after the fourth booster immunization, for immunoprecipitation of Flt4 from transfected cells. Cell clones expressing Flt4 were also used for ligand stimulation analysis.

EXAMPLE 3

Construction of a Flt4 EC baculovirus vector and expression and purification of its product

Using the pVTBac plasmid described in Tessier *et al.*, *Gene* 98:177-183 (1991), and the Flt4 cDNAs described in Example 1, a baculovirus expression vector was constructed to facilitate expression of the extracellular domain of Flt4 (Flt4 EC), as described in commonly-owned PCT patent application PCT/FI96/00427, filed August 01, 1996, published as PCT publication No. WO 97/05250 on 13 February 1997, and commonly-owned United States Patent Application Serial Nos. 08/671,573, filed June 28, 1996; 08/601,132, filed February 14, 1996; 08/585,895, filed January 12, 1996; and 08/510,133, filed August 1, 1995, all of which are incorporated by reference herein. A nucleotide sequence encoding a 6xHis tag was operatively connected to the Flt4 EC coding sequence to facilitate purification.

The Flt4EC construct was transfected together with baculovirus genomic DNA into SF-9 cells by lipofection. Recombinant virus was purified, amplified and used for infection of High-Five cells (Invitrogen, San Diego, CA) using methods standard in the art. The Flt4 extracellular domain (Flt4EC) was purified from the culture medium of the infected High-Five cells using Ni-NTA affinity chromatography according to manufacturer's instructions (Qiagen) for binding and elution of the 6xHis tag encoded in the COOH-terminus of the recombinant Flt4 extracellular domain.

EXAMPLE 4

Isolation of an Flt4 Ligand from Conditioned Media

A human Flt4 ligand according to the invention was isolated from media conditioned by a PC-3 prostatic adenocarcinoma cell line (ATCC CRL 1435) in serum-free Ham's F-12 Nutrient mixture (GIBCO) (containing 7% fetal calf serum (FCS)). Cells were

- 35 -

reseeded and grown in this medium, which was subsequently changed to serum-free medium. The preparation of the conditioned media, and the identification of a component therein which stimulated Flt4 tyrosine phosphorylation, are described in detail in commonly-owned PCT patent application PCT/FI96/00427, filed August 01, 1996, and 5 commonly-owned United States Patent Application Serial Nos. 08/671,573, filed June 28, 1996; 08/601,132, filed February 14, 1996; 08/585,895, filed January 12, 1996; 08/510,133, filed August 1, 1995; and 08/340,011, filed November 14, 1994, all of which are incorporated by reference herein in their entirety. The ability of the conditioned medium to stimulate Flt4 phosphorylation was considerably increased when the PC-3 10 conditioned medium was concentrated four-fold using a Centricon-10 concentrator (Amicon). Pretreatment of the concentrated PC-3 conditioned medium with 50 microliters of Flt4 extracellular domain coupled to CNBr-activated sepharose CL-4B (Pharmacia; about 1mg of Flt4EC domain/ml sepharose resin) completely abolished Flt4 tyrosine phosphorylation. Similar pretreatment of the conditioned medium with unsubstituted 15 sepharose CL-4B did not affect stimulatory activity. Also, the flow through obtained after concentration, which contained proteins of less than 10,000 molecular weight, did not stimulate Flt4 phosphorylation.

In another experiment, a comparison of Flt4 autophosphorylation in transformed NIH 3T3 cells expressing LTRFlt4l was conducted, using unconditioned 20 medium, medium from PC-3 cells expressing the Flt4 ligand, or unconditioned medium containing either 50 ng/ml of VEGF165 or 50 ng/ml of PlGF-1. The cells were lysed, immunoprecipitated using anti-Flt4 antiserum and analyzed by Western blotting using anti-phosphotyrosine antibodies. Only the PC-3 conditioned medium expressing the Flt4 ligand (lane Flt-4L) stimulated Flt4 autophosphorylation.

25 These experiments showed that PC-3 cells produce a ligand which binds to the extracellular domain of Flt4 and activates this receptor.

EXAMPLE 5

Purification of the Flt4 Ligand

The ligand expressed by human PC-3 cells as characterized in Example 4 30 was purified and isolated using a recombinantly-produced Flt4 extracellular domain (Flt4EC) in affinity chromatography.

Two harvests of serum-free conditioned medium, comprising a total of 8 liters, were collected from 500 confluent 15 cm diameter culture dishes containing confluent layers of PC-3 cells. The conditioned medium was clarified by centrifugation at 10,000 x g and concentrated 80-fold using an Ultrasette Tangential Flow Device (Filtron, 5 Northborough, MA) with a 10 kD cutoff Omega Ultrafiltration membrane according to the manufacturer's instructions. Recombinant Flt4 extracellular domain was expressed in a recombinant baculovirus cell system and purified by affinity chromatography on Ni-agarose (Ni-NTA affinity column obtained from Qiagen). The purified extracellular domain was coupled to CNBr-activated Sepharose CL-4B at a concentration of 5 mg/ml 10 and used as an affinity matrix for ligand affinity chromatography.

Concentrated conditioned medium was incubated with 2 ml of the recombinant Flt4 extracellular domain-Sepharose affinity matrix in a rolling tube at room temperature for 3 hours. All subsequent purification steps were at +4 °C. The affinity matrix was then transferred to a column with an inner diameter of 15 mm and washed 15 successively with 100 ml of PBS and 50 ml of 10 mM Na-phosphate buffer (pH 6.8). Bound material was eluted step-wise with 100 mM glycine-HCl, successive 6 ml elutions having pHs of 4.0, 2.4, and 1.9. Several 2 ml fractions of the eluate were collected in tubes containing 0.5 ml 1 M Na-phosphate (pH 8.0). Fractions were mixed immediately and dialyzed in 1 mM Tris-HCl (pH 7.5). Aliquots of 75 µl each were analyzed for their 20 ability to stimulate tyrosine phosphorylation of Flt4. The ultrafiltrate, 100 µl aliquots of the concentrated conditioned medium before and after ligand affinity chromatography, as well as 15-fold concentrated fractions of material released from the Flt4 extracellular domain-Sepharose matrix during the washings were also analyzed for their ability to stimulate Flt4 tyrosine phosphorylation.

25 The concentrated conditioned medium induced prominent tyrosine phosphorylation of Flt4 in transfected NIH 3T3 cells over-expressing Flt4. This activity was not observed in conditioned medium taken after medium was exposed to the Flt4 Sepharose affinity matrix. The specifically-bound Flt4-stimulating material was retained on the affinity matrix after washing in PBS, 10 mM Na-phosphate buffer (pH 6.8), and at 30 pH 4.0. It was eluted in the first two 2 ml aliquots at pH 2.4. A further decrease of the pH of the elution buffer did not cause release of additional Flt4-stimulating material. No Flt4 phosphorylation was observed in a control wherein Flt4-expressing cells were treated

with unconditioned medium; similarly, no phosphorylation was observed following treatment of Flt4-expressing cells with the ultrafiltrate fraction of conditioned medium containing polypeptides of less than 10 kD molecular weight.

Small aliquots of the chromatographic fractions were concentrated in a SpeedVac concentrator (Savant, Farmingdale, N.Y.) and subjected to SDS-PAGE under reducing conditions with subsequent silver staining of the gel, a standard technique in the art. The major polypeptide, having a molecular weight of approximately 23 kD (reducing conditions), was detected in the fractions containing Flt4 stimulating activity. That polypeptide was not found in the other chromatographic fractions. On the other hand, besides these bands and a very faint band having a 32 kD mobility, all other components detected in the two active fractions were also distributed in the starting material and in small amounts in the other washing and eluting steps after their concentration. Similar results were obtained in three independent affinity purifications, indicating that the 23 kD polypeptide binds with high affinity to Flt4 and induces tyrosine phosphorylation of Flt4.

Fractions containing the 23 kD polypeptide were combined, dried in a SpeedVac concentrator and subjected to SDS-PAGE in a 12.5% gel. The proteins from the gel were then electroblotted to Immobilon-P (PVDF) transfer membrane (Millipore, Marlborough, MA) and visualized by staining of the blot with Coomassie Blue R-250. The region containing only the stained 23 kD band was cut from the blot and subjected to N-terminal amino acid sequence analysis in a Prosite Protein Sequencing System (Applied Biosystems, Foster City, CA). The data were analyzed using a 610A Data Analysis System (Applied Biosystems). Analysis revealed a single N-terminal sequence of NH₂-XEETIKFAAAHYNTEILK-COOH (SEQ ID NO: 5).

EXAMPLE 6

Construction of PC-3 cell cDNA library in a eukaryotic expression vector

Human poly(A)⁺ RNA was isolated from five 15 cm diameter dishes of confluent PC-3 cells by a single step method using oligo(dT) (Type III, Collaborative Biomedical Products, Becton-Dickinson Labware, Bedford, MA) cellulose affinity chromatography (Sambrook *et al.*, 1989). The yield was 70 micrograms. Six micrograms of the Poly(A)⁺ RNA were used to prepare an oligo(dT)-primed cDNA library in the

mammalian expression vector pcDNA I and the Librarian kit of Invitrogen according to the instructions included in the kit. The library was estimated to contain about 10^6 independent recombinants with an average insert size of approximately 1.8 kb.

EXAMPLES 7-9

5

Amplification of a cDNA encoding the Flt4 ligand amino terminus

The procedures used to isolate a cDNA encoding the Flt4 ligand are described in detail in commonly-owned PCT patent application PCT/FI96/00427, filed August 01, 1996, and commonly-owned United States Patent Application Serial Nos. 10 08/671,573, filed June 28, 1996; 08/601,132, filed February 14, 1996; 08/585,895, filed January 12, 1996; and 08/510,133, filed August 1, 1995, all of which are incorporated by reference herein. Initially, degenerate oligonucleotides were designed based on the N-terminal amino acid sequence of the isolated human Flt4 ligand (see Example 5) and were used as primers in a polymerase chain reaction (PCR) to amplify a partial cDNA encoding 15 the (fully-processed) Flt4 ligand amino terminus from the PC-3 cDNA library. The amplified cDNA fragment was cloned into a pCR II vector (Invitrogen) using the TA cloning kit (Invitrogen) and sequenced using the radioactive dideoxynucleotide sequencing method of Sanger. Six clones were analyzed and all six clones contained the sequence encoding the expected peptide (amino acid residues 104-120 of the Flt4 ligand precursor, 20 SEQ ID NO: 8). Nucleotide sequence spanning the region from the third nucleotide of codon 6 to the third nucleotide of codon 13 (the extension region between the PCR primers) was identical in all six clones and thus represented an amplified product from the unique sequence encoding part of the amino terminus of the Flt4 ligand.

Based on the unique nucleotide sequence encoding the N-terminus of the 25 isolated human Flt4 ligand, two pairs of nested primers were designed to amplify, in two nested PCR reactions, the complete 5'-end of the corresponding cDNAs from one microgram of DNA of the above-described PC-3 cDNA library. One major product of about 220 bp and three minor products of about 270 bp, 150 bp, and 100 bp were obtained.

30

The amplified fragment of approximately 220 bp was excised from an agarose gel, cloned into a pCRII vector using the TA cloning kit, and sequenced. Three

- 39 -

recombinant clones were analyzed and they contained the sequence 5'-
 TCACTATAGGGAGACCCAAGCTTGGTACCGAGCTCGGATCCACTAGTAACGGC
 CGCCAGTGTGGTGGGAATTCGACGAACTCATGACTGTACTCTACCCAGAATATT
GGAAAATGTACAAGTGTCTAGCTAAGGCAAGGAGGCTGGCAACATAACAGAGA
 5 ACAGGCCAACCTCAACTCAAGGACAGAAGAGACTATAAAATTCGCTGCAGCA
CACTACAAC- 3' (SEQ ID NO: 6). The beginning of the sequence represents the vector
 and the underlined sequence represents the amplified product of the 5'-end of the cDNA
 insert.

Based upon the amplified 5'-sequence of the clones encoding the amino
 10 terminus of the 23 kD human Flt4 ligand, two pairs of non-overlapping nested primers
 were designed to amplify the 3'-portion of the Flt4-ligand-encoding cDNA clones via PCR.
 Two DNA fragments were obtained, having sizes of 1350 bp and 570 bp. Those
 fragments were cloned into a pCRII vector and the inserts of the clones were sequenced.
 Both of these fragments were found to contain sequences encoding an amino acid
 15 sequence homologous to the VEGF sequence.

EXAMPLE 10

Screening the PC-3 cell cDNA library using the 5' PCR fragment of Flt4 ligand cDNA

A 153 bp fragment encoding the 5' end of the Flt4 ligand was labeled with
 20 [³²P]-dCTP using the Klenow fragment of *E. coli* DNA polymerase I (Boehringer
 Mannheim). That fragment was used as a probe for hybridization screening of the
 amplified PC-3 cell cDNA library.

Filter replicas of the library were hybridized with the radioactively labeled
 probe at 42°C for 20 hours in a solution containing 50% formamide, 5x SSPE, 5x
 25 Denhardt's solution, 0.1% SDS and 0.1 mg/ml denatured salmon sperm DNA. Filters
 were washed twice in 1x SSC, 0.1% SDS for 30 minutes at room temperature, then twice
 for 30 minutes at 65°C and exposed overnight.

On the basis of autoradiography, 10 positive recombinant bacterial colonies
 hybridizing with the probe were chosen from the library. Plasmid DNA was purified from
 30 these colonies and analyzed by *Eco*RI and *Not*I digestion and agarose gel electrophoresis
 followed by ethidium bromide staining. The ten plasmid clones were divided into three

groups on the basis of the presence of insert sizes of approximately 1.7, 1.9 and 2.1 kb, respectively. Inserts of plasmids from each group were sequenced using the T7 oligonucleotide as a primer and walking primers for subsequent sequencing reactions.

Sequence analysis showed that all clones contain the open reading frame 5 encoding the NH₂-terminal sequence of the 23 kD human Flt4 ligand. Dideoxy sequencing was continued using walking primers in the downstream direction. A complete human cDNA sequence and deduced amino acid sequence from a 2 kb clone is set forth in SEQ ID NOs: 7 and 8, respectively. A putative cleavage site of a "prepro" leader sequence is located between residues 102 and 103 of SEQ ID NO: 8. When compared 10 with sequences in the GenBank Database, the predicted protein product of this reading frame was found to include a region homologous with the predicted amino acid sequences of the PDGF/VEGF family of growth factors, as shown in Fig. 2.

Plasmid pFLT4-L, containing the 2.1 kb human cDNA clone in pcDNA1 vector, has been deposited with the American Type Culture Collection, 12301 Parklawn 15 Drive, Rockville, MD 20852 as accession number 97231.

EXAMPLE 11

Stimulation of Flt4 autophosphorylation by the protein product of the Flt4 ligand vector

The 2.1 kb human cDNA insert of plasmid pFlt4-L, which contains the 20 open reading frame encoding the sequence shown in SEQ ID NOs: 7 and 8; human prepro- VEGF-C, see below), was cut out from the pcDNA1 vector using *Hind*III and *Not*I restriction enzymes, isolated from a preparative agarose gel, and ligated to the corresponding sites in the pREP7 expression vector (Invitrogen). The pREP7 vector containing the pFlt4-L insert was transfected into 293-EBNA cells (Invitrogen) using the 25 calcium phosphate transfection method (Sambrook *et al.*, 1989). About 48 hours after transfection, the medium of the transfected cells was changed to DMEM medium lacking fetal calf serum and incubated for 36 hours. The conditioned medium was then collected, centrifuged at 5000 x g for 20 minutes, the supernatant was concentrated 5-fold using Centriprep 10 (Amicon) and used to stimulate NIH 3T3 cells expressing LTRFlt4l (the 30 Flt4 receptor), as in Example 4. The cells were lysed, immunoprecipitated using anti-Flt4 antiserum and analyzed by Western blotting using anti-phosphotyrosine antibodies.

- 41 -

The conditioned medium from two different dishes of the transfected cells stimulated Flt4 autophosphorylation in comparison with the medium from mock-transfected cells, which gave only background levels of phosphorylation of the Flt4 receptor. When the concentrated conditioned medium was pre-absorbed with 20
5 microliters of a slurry of Flt4EC domain coupled to Sepharose (see example 4), no phosphorylation was obtained, showing that the activity responsible for Flt4 autophosphorylation was indeed the Flt4 ligand. Thus, these results demonstrate that an expression vector having an approximately 2.1 kb insert and containing an open reading frame as shown in SEQ ID NO: 7 is expressed as a biologically active Flt4 ligand (VEGF-
10 C) in transfected cells. The sequence encoded by that open reading frame is shown in SEQ ID NO: 8.

The deduced molecular weight of a polypeptide consisting of the complete amino acid sequence in SEQ ID NO: 8 (residues 1 to 419) is 46,883. The deduced molecular weight of a polypeptide consisting of amino acid residues 103 to 419 of SEQ ID
15 NO: 8 is 35,881. The Flt4 ligand purified from PC-3 cultures had an observed molecular weight of about 23 kD as assessed by SDS-PAGE under reducing conditions. Thus, it appeared that the Flt4 ligand mRNA was translated into a precursor polypeptide, from which the mature ligand was derived by proteolytic cleavage. Also, the Flt4 ligand may be glycosylated at three putative N-linked glycosylation sites conforming to the consensus
20 which can be identified in the deduced Flt4 ligand amino acid sequence (N-residues underlined in Fig. 2).

The carboxyl terminal amino acid sequences, which increase the predicted molecular weight of the Flt4 ligand subunit in comparison with other ligands of this family, show a pattern of spacing of cysteine residues reminiscent of the Balbiani ring 3 protein
25 (BR3P) sequence (Dignam *et al.*, *Gene*, 88:133-140 (1990)). Such a sequence may encode an independently folded domain present in a Flt4 ligand precursor and it may be involved, for example, in the regulation of secretion, solubility, stability, cell surface localization or activity of the Flt4 ligand. Interestingly, at least one cysteine motif of the BR3P type is also found in the VEGF carboxy terminal amino acid sequences.

30 Thus, the Flt4 ligand mRNA appears first to be translated into a precursor from the mRNA corresponding to the cDNA insert of plasmid FLT4-L, from which the mature ligand is derived by proteolytic cleavage. To define the mature Flt4 ligand

polypeptide, one first expresses the cDNA clone (which is deposited in the pcDNAI expression vector) in cells, such as COS cells. One uses antibodies generated against encoded polypeptides, fragments thereof, or bacterial Flt4 fusion proteins, such as a GST-fusion protein, to raise antibodies against the VEGF-homologous domain and the amino-
5 and carboxyl-terminal propeptides of Flt4 ligand. One then follows the biosynthesis and processing of the Flt4 ligand in the transfected cells by pulse-chase analysis using radioactive cysteine for labeling of the cells, immunoprecipitation, and gel electrophoresis. Using antibodies against the three domains of the product encoded by the cDNA insert of plasmid FLT4-L, material for radioactive or nonradioactive amino-terminal sequence
10 analysis is isolated. The determination of the amino-terminal sequence of the mature VEGF-C polypeptide allows for identification of the amino-terminal proteolytic processing site. The determination of the amino-terminal sequence of the carboxyl-terminal propeptide will give the carboxyl-terminal processing site. This is confirmed by site-directed mutagenesis of the amino acid residues adjacent to the cleavage sites, which
15 would prevent the cleavage.

The Flt4 ligand is further characterizeable by progressive 3' deletions in the 3' coding sequences of the Flt4 ligand precursor clone, introducing a stop codon resulting in carboxy-terminal truncations of its protein product. The activities of such truncated forms are assayed by, for example, studying Flt4 autophosphorylation induced by the
20 truncated proteins when applied to cultures of cells, such as NIH 3T3 cells expressing LTRFlt4l. By extrapolation from studies of the structure of the related platelet derived growth factor (PDGF, Heldin *et al.*, *Growth Factors*, 8:245-252 (1993)) one determines that the region critical for receptor activation by the Flt4 ligand is contained within the first approximately 180 amino acid residues of the secreted VEGF-C protein lacking the
25 putative 102 amino acid prepro leader (SEQ ID NO: 8, residues 103-282), and apparently within the first approximately 120 amino acid residues (SEQ ID NO: 8, residues 103-223).

On the other hand, the difference between the molecular weights observed for the purified ligand and deduced from the open reading frame of the Flt4 ligand clone may be due to the fact that the soluble ligand was produced from an alternatively spliced
30 mRNA which would also be present in the PC-3 cells, from which the isolated ligand was derived. To isolate such alternative cDNA clones one uses cDNA fragments of the deposited clone and PCR primers made according to the sequence provided as well as

techniques standard in the art to isolate or amplify alternative cDNAs from the PC-3 cell cDNA library. One may also amplify using reverse transcription (RT)-PCR directly from the PC-3 mRNA using the primers provided in the sequence of the cDNA insert of plasmid FLT4-L. Alternative cDNA sequences are determined from the resulting cDNA clones.

5 One can also isolate genomic clones corresponding to the Flt4 ligand mRNA transcript from a human genomic DNA library using methods standard in the art and sequence such clones or their subcloned fragments to reveal the corresponding exons. Alternative exons can then be identified by a number of methods standard in the art, such as heteroduplex analysis of cDNA and genomic DNA, which are subsequently characterized.

10

EXAMPLE 12

Expression of the Gene Encoding VEGF-C in Human Tumor Cell Lines

Expression of transcripts corresponding to the Flt4 ligand (VEGF-C) was analyzed by hybridization of Northern blots containing isolated poly(A)⁺ RNA from HT-15 1080 and PC-3 human tumor cell lines. The probe was the radioactively labeled insert of the 2.1 kb cDNA clone (pFlt4-L/VEGF-C, specific activity 10^8 - 10^9 cpm/mg of DNA). The blot was hybridized overnight at 42°C using 50% formamide, 5x SSPE buffer, 2% SDS, 10 x Denhardt's solution, 100 mg/ml salmon sperm DNA and 1×10^6 cpm of the labeled probe/ml. The blot was washed at room temperature for 2 x 30 minutes in 2x SSC 20 containing 0.05% SDS, and then for 2 x 20 minutes at 52°C in 0.1x SSC containing 0.1% SDS. The blot was then exposed at -70°C for three days using intensifying screens and Kodak XAR film. Both cell lines expressed an Flt4 ligand mRNA of about 2.4 kb, as well as VEGF and VEGF-B mRNAs.

EXAMPLE 13

25

VEGF-C Chains Are Proteolytically Processed after Biosynthesis and Disulfide Linked

The predicted molecular mass of a secreted human VEGF-C polypeptide, as deduced from the VEGF-C open reading frame, is 46,883 kD, suggesting that VEGF-C mRNA may be first translated into a precursor, from which the observed ligands of 21/23 30 kD and 29/32 kD are derived by proteolytic cleavage.

This possibility was explored by metabolic labeling of 293 EBNA cells expressing VEGF-C. Initially, 293 EBNA cells were transfected with the VEGF-C cDNA construct. Expression products were labeled by the addition of 100 $\mu\text{Ci/ml}$ of Pro-mixTM L-[³⁵S] *in vitro* cell labeling mix ((containing ³⁵S-methionine and ³⁵S-cysteine) Amersham, 5 Buckinghamshire, England) to the culture medium devoid of cysteine and methionine. After two hours, the cell layers were washed twice with PBS and the medium was then replaced with DMEM-0.2% BSA. After 1, 3, 6, 12 and 24 hours of subsequent incubation, the culture medium was collected, clarified by centrifugation, and concentrated, and human VEGF-C was bound to 30 μl of a slurry of Flt4EC-Sepharose 10 overnight at +4°C, followed by three washes in PBS, two washes in 20 mM Tris-HCl (pH 7.5), alkylation, SDS-PAGE and autoradiography. Alkylation was carried out by treatment of the samples with 10 mM 1,4 Dithiothreitol (Boehringer-Mannheim, Mannheim, Germany) for one hour at 25°C, and subsequently with 30 mM iodoacetamide (Fluka, Buchs, Switzerland).

15 These experiments demonstrated that a putative precursor polypeptide of 32 kD apparent molecular mass was bound to the Flt4EC affinity matrix from the conditioned medium of metabolically labeled cells transfected with the human VEGF-C expression vector, but not from mock transfected cells. Increased amounts of a 23 kD receptor binding polypeptide accumulated in the culture medium of VEGF-C transfected 20 cells during a subsequent chase period of three hours, but not thereafter, suggesting that the 23 kD form is produced by proteolytic processing, which is incomplete, at least in the transiently transfected cells. Subsequent experiments showed that the 32kD VEGF-C form contains two components migrating in the absence of alkylation as polypeptides of 29 and 32 kD (Figs. 6-8).

25 In a related experiment, human VEGF-C isolated using Flt4EC-Sepharose after a 4 hour continuous metabolic labeling was analyzed by polyacrylamide gel electrophoresis in nonreducing conditions. Higher molecular mass forms were observed under nonreducing conditions, suggesting that the VEGF-C polypeptides can form disulfide-linked dimers and/or multimers. Gel photographs depicting these experimental 30 results are set forth in Figures 13A-B of PCT application PCT/FI96/00427 (publication WO 97/05250) and Figures 3A-B of U.S. Patent Application Serial No. 08/795,430, which are incorporated herein by reference.

Additional experiments have shown that higher molecular mass forms of VEGF-C (about 58 kD and about 43 kD) are observed under reducing conditions as well. (See below and Fig. 6A.)

EXAMPLE 14

5 Stimulation Of VEGFR-2 Autophosphorylation By VEGF-C

Conditioned medium (CM) from 293 EBNA cells transfected with the human VEGF-C vector also was used to stimulate porcine aortic endothelial (PAE) cells expressing VEGFR-2 (KDR). Pajusola *et al.*, *Oncogene*, 9:3545-55 (1994); Waltenberger *et al.*, *J. Biol. Chem.*, 269:26988-26995 (1994). The cells were lysed and
10 immunoprecipitated using VEGFR-2 - specific antiserum (Waltenberger *et al.*, 1994).

PAE-KDR cells (Waltenberger *et al.*, 1994) were grown in Ham's F12 medium-10% fetal calf serum (FCS). Confluent NIH 3T3-Flt4 cells or PAE-KDR cells were starved overnight in DMEM or Ham's F12 medium, respectively, supplemented with 0.2% bovine serum albumin (BSA), and then incubated for 5 minutes with the analyzed
15 media. Recombinant human VEGF (R&D Systems) and PDGF-BB, functional as stimulating agents, were used as controls. The cells were washed twice with ice-cold Tris-Buffered Saline (TBS) containing 100 mM sodium orthovanadate and lysed in RIPA buffer containing 1 mM phenylmethylsulfonyl fluoride (PMSF), 0.1 U/ml aprotinin and 1 mM sodium orthovanadate. The lysates were sonicated, clarified by centrifugation at
20 16,000 x g for 20 minutes and incubated for 3-6 hours on ice with 3-5 µl of antisera specific for Flt4 (Pajusola *et al.*, 1993), VEGFR-2 or PDGFR-β (Claesson-Welsh *et al.*, *J. Biol. Chem.*, 264:1742-1747 (1989); Waltenberger *et al.*, 1994). Immunoprecipitates were bound to protein A-Sepharose, washed three times with RIPA buffer containing 1mM PMSF, 1mM sodium orthovanadate, washed twice with 10 mM Tris-HCl (pH 7.4),
25 and subjected to SDS-PAGE using a 7% gel. Polypeptides were transferred to nitrocellulose by Western blotting and analyzed using PY20 phosphotyrosine-specific monoclonal antibodies (Transduction Laboratories) or receptor-specific antiserum and the ECL detection method (Amersham Corp.).

PAE cells expressing VEGFR-2 were treated with 10- or 2-fold
30 concentrated medium from mock-transfected 293-EBNA cells, or with 2-, 5- or 10-fold concentrated medium from 293-EBNA cell cultures expressing the recombinant VEGF-C.

VEGFR-2 was immunoprecipitated with specific antibodies and analyzed by SDS-PAGE and Western blotting using phosphotyrosine antibodies. For comparison, the treatments were also carried out with non-conditioned medium containing 50 ng/ml of purified recombinant VEGF. Additional cells were also treated with VEGF-C- or VEGF-5 containing media pretreated with Flt4EC.

The results of this experiment were as follows. A basal level of tyrosine phosphorylation of VEGFR-2 was detected in cells stimulated by CM from the mock-transfected cells. A further concentration of this medium resulted in only a slight enhancement of VEGFR-2 phosphorylation. CM containing recombinant VEGF-C 10 stimulated tyrosine autophosphorylation of VEGFR-2 and the intensity of the autophosphorylated polypeptide band was increased upon concentration of the VEGF-C CM. Furthermore, the stimulating effect was abolished after pretreatment of the medium with the Flt4EC affinity matrix. The maximal effect of VEGF-C in this assay was comparable to the effect of recombinant VEGF added to unconditioned medium at 15 concentration of 50 ng/ml. Pretreatment of the medium containing VEGF with Flt4EC did not abolish its stimulating effect on VEGFR-2. These results suggest that the VEGF-C expression vector encodes a ligand not only for Flt4 (VEGFR-3), but also for KDR/Flk-1 (VEGFR-2).

In order to further confirm that the stimulating effect of VEGF-C on 20 tyrosine phosphorylation of VEGFR-3 and VEGFR-2 was receptor-specific, we analyzed the effect of VEGF-C on tyrosine phosphorylation of PDGF receptor β (PDGFR- β) which is abundantly expressed on fibroblastic cells. PDGFR- β -expressing NIH 3T3 cells were treated with non-conditioned medium, 5-fold concentrated CM from mock-transfected or VEGF-C- transfected cells, or with non-conditioned medium containing 50 ng/ml of 25 recombinant human PDGF-BB. Medium containing VEGF-C was also pretreated with recombinant Flt4EC (lane 4). PDGFR- β was immunoprecipitated with specific antibodies and analyzed by SDS-PAGE and Western blotting using phosphotyrosine antibodies with subsequent stripping and reprobing of the membrane with antibodies specific for PDGFR- β . A weak tyrosine phosphorylation of PDGFR- β was detected upon stimulation 30 of Flt4-expressing NIH 3T3 cells with CM from the mock-transfected cells. A similar low level of PDGFR- β phosphorylation was observed when the cells were incubated with CM from the VEGF-C transfected cells, with or without prior treatment with Flt4EC. In

contrast, the addition of 50 ng/ml of PDGF-BB induced a prominent tyrosine autophosphorylation of PDGFR- β .

EXAMPLE 15

VEGF-C Stimulates Endothelial Cell Migration In Collagen Gel

5 Conditioned media (CM) from cell cultures transfected with the VEGF-C expression vector was placed in a well made in collagen gel and used to stimulate the migration of bovine capillary endothelial (BCE) cells in the three-dimensional collagen gel as follows.

10 BCE cells (Folkman *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 76:5217-5221 (1979)) were cultured as described in Pertovaara *et al.*, *J. Biol. Chem.*, 269:6271-74 (1994). The collagen gels were prepared by mixing type I collagen stock solution (5 mg/ml in 1 mM HCl) with an equal volume of 2x MEM and 2 volumes of MEM containing 10% newborn calf serum to give a final collagen concentration of 1.25 mg/ml.

15 The tissue culture plates (5 cm diameter) were coated with about 1 mm thick layer of the solution, which was allowed to polymerize at 37°C. BCE cells were seeded on top of this layer. For the migration assays, the cells were allowed to attach inside a plastic ring (1 cm diameter) placed on top of the first collagen layer. After 30 minutes, the ring was removed and unattached cells were rinsed away. A second layer of collagen and a layer of growth

20 medium (5% newborn calf serum (NCS)), solidified by 0.75% low melting point agar (FMC BioProducts, Rockland, ME), were added. A well (3 mm diameter) was punched through all the layers on both sides of the cell spot at a distance of 4 mm, and the sample or control media were pipetted daily into the wells. Photomicrographs of the cells migrating out from the spot edge were taken after six days through an Olympus CK 2

25 inverted microscope equipped with phase-contrast optics. The migrating cells were counted after nuclear staining with the fluorescent dye bisbenzimidazole (1 mg/ml, Hoechst 33258, Sigma).

 The number of cells migrating at different distances from the original area of attachment towards wells containing media conditioned by the non-transfected (control)

30 or transfected (mock; VEGF-C; VEGF) cells were determined 6 days after addition of the media. The number of cells migrating out from the original ring of attachment was

counted in five adjacent 0.5 mm x 0.5 mm squares using a microscope ocular lens grid and 10x magnification with a fluorescence microscope. Cells migrating further than 0.5 mm were counted in a similar way by moving the grid in 0.5 mm steps. The experiments were carried out twice with similar results. At each distance, VEGF-C-containing CM

5 stimulated cell migration more than medium conditioned by the non-transfected or mock-transfected cells but less than medium from cells transfected with a VEGF expression vector. Daily addition of 1 ng of FGF2 into the wells resulted in the migration of approximately twice the number of cells when compared to the stimulation by CM from VEGF-transfected cells.

10 In related experiments, a "recombinantly-matured" VEGF-C polypeptide (VEGF-C Δ N Δ CHis, described below) was shown to stimulate the incorporation of 3 H-thymidine into the DNA of BCE cells in a dose dependent manner (VEGF-C concentrations of 0, 10, 100, and 1000 pM tested). This data tends to confirm the observation, under light microscopy, that VEGF-C stimulates proliferation of these cells.

15

EXAMPLE 16

VEGF-C Is Expressed In Multiple Tissues

Northern blots containing 2 micrograms of isolated poly(A)⁺ RNA from multiple human tissues (blot from Clontech Laboratories, Inc., Palo Alto, CA) were probed with radioactively labeled insert of the 2.1 kb VEGF-C cDNA clone. Northern
20 blotting and hybridization analysis showed that the 2.4 kb RNA and smaller amounts of a 2.0 kb mRNA are expressed in multiple human tissues, most prominently in the heart, placenta, muscle, ovary and small intestine, and less prominently in prostate, colon, lung, pancreas, and spleen. Very little VEGF-C RNA was seen in the brain, liver, kidney, testis, or thymus and peripheral blood leukocytes (PBL) appeared negative. A similar analysis of
25 RNA from human fetal brain, lung, liver, and kidney tissues showed that VEGF-C is highly expressed in the kidney and lung and to a lesser degree in the liver, while essentially no expression is detected in the brain. Interestingly, VEGF expression correlates with VEGF-C expression in these tissues, whereas VEGF-B is highly expressed in all four fetal tissues analyzed.

EXAMPLE 17**The VEGF-C Gene Localizes To Chromosome 4q34**

A DNA panel of 24 interspecies somatic cell hybrids, which had retained one or two human chromosomes, was used for the chromosomal localization of the VEGF-C gene (Bios Laboratories, Inc., New Haven, CT). DNAs from human rodent somatic cell hybrids containing defined sets of human chromosomes were analyzed by Southern blotting and hybridization with a VEGF-C cDNA probe. Among 24 DNA samples on the hybrid panel, representing different human chromosomes, human-specific signals were observed only in hybrids which contained human chromosome 4. The results were confirmed by PCR of somatic cell hybrid DNAs using VEGF-C specific primers, where amplified bands were obtained only from DNAs containing human chromosome 4.

A genomic P1 plasmid for VEGF-C was isolated using specific primers and PCR and verified by Southern blotting and hybridization using a VEGF-C specific cDNA probe. The chromosomal localization of VEGF-C was further studied using metaphase FISH. Using the P1 probe for VEGF-C in FISH, a specific hybridization to the 4q34 chromosomal band was detected in 40 out of 44 metaphases. Double-fluorochrome hybridization using a cosmid probe specific for the aspartylglucosaminidase (AGA) gene showed that VEGF-C is located just proximal to the AGA gene previously mapped to the 4q34-35 chromosomal band.

Biotin-labeled VEGF-C P1 and digoxigenin-labeled AGA cosmid probes were hybridized simultaneously to metaphase chromosomes. This experiment demonstrated that the AGA gene is more telomerically located than the VEGF-C gene. The foregoing example demonstrates the utility of polynucleotides of the invention as chromosomal markers and for the presence or absence of the VEGF-C gene region in normal or diseased cells. The VEGF-C locus at 4q34 is a candidate target for mutations leading to vascular malformations or cardiovascular diseases.

EXAMPLE 18**Effect of glucose concentration and hypoxia on VEGF, VEGF-B and VEGF-C mRNA levels in C6 glioblastoma cells**

Confluent cultures of C6 cells (ATCC CCL 107) were grown on 10 cm diameter tissue culture plates containing 2.5 ml of DMEM and 5% fetal calf serum plus

antibiotics. The cultures were exposed for 16 hours to normoxia in a normal cell culture incubator containing 5% CO₂ or hypoxia by closing the culture plates in an airtight glass chamber and burning a piece of wood inside until the flame was extinguished due to lack of oxygen. Polyadenylated RNA was isolated (as in the other examples), and 8
5 micrograms of the RNA was electrophoresed and blot-hybridized with a mixture of the VEGF, VEGF-B and VEGF-C probes. The results show that hypoxia strongly induces VEGF mRNA expression, both in low and high glucose, but has no significant effect on the VEGF-B mRNA levels. The VEGF-C mRNA isolated from hypoxic cells runs slightly faster in gel electrophoresis and an extra band of faster mobility can be seen below the
10 upper mRNA band. This observation suggests that hypoxia affects VEGF-C RNA processing. One explanation for this observation is that VEGF-C mRNA splicing is altered, affecting the VEGF-C open reading frame and resulting in an alternative VEGF-C protein being produced by hypoxic cells. Such alternative forms of VEGF-C and VEGF-C-encoding polynucleotides are contemplated as an aspect of the invention. This data
15 indicates screening and diagnostic utilities for polynucleotides and polypeptides of the invention, such as methods whereby a biological sample is screened for the hypoxia-induced form of VEGF-C and/or VEGF-C mRNA. The data further suggests a therapeutic indication for antibodies and/or other inhibitors of the hypoxia-induced form of VEGF-C or the normal form of VEGF-C.

20

EXAMPLE 19

**Pulse-chase labeling and immunoprecipitation
of VEGF-C polypeptides from 293 EBNA cells
transfected with VEGF-C expression vector.**

The following VEGF-C branched amino-terminal peptide, designated
25 PAM126, was synthesized for production of anti-VEGF-C antiserum:

NH₂-E-E-T-I-K-F-A-A-A-H-Y-N-T-E-I-L-K-COOH (SEQ ID NO: 9).

In particular, PAM126 was synthesized as a branched polylysine structure K3PA4 having four peptide acid (PA) chains attached to two available lysine (K) residues. The synthesis was performed on a 433A Peptide Synthesizer (Applied Biosystems) using
30 Fmoc-chemistry and TentaGel S MAP RAM10 resin mix (RAPP Polymere GmbH, Tübingen, Germany), yielding both cleavable and resin-bound peptides. The cleavable

peptide was purified via reverse phase HPLC and was used together with the resin-bound peptide in immunizations. The correctness of the synthesis products were confirmed using mass-spectroscopy (Lasermatt).

The PAM126 peptide was dissolved in phosphate buffered saline (PBS),
5 mixed with Freund's adjuvant, and used for immunization of rabbits at bi-weekly intervals using methods standard in the art (Harlow and Lane, *Antibodies, a laboratory manual*, Cold Spring Harbor Laboratory Press (1988)). Antisera obtained after the fourth booster immunization was used for immunoprecipitation of VEGF-C in pulse-chase experiments, as described below.

10 For pulse-chase analysis, 293 EBNA cells transfected with a VEGF-C expression vector (i.e., the FLT4-L cDNA inserted into the pREP7 expression vector as described above) were incubated for 30 minutes in methionine-free, cysteine-free, serum-free DMEM culture medium at 37°C. The medium was then changed, and 200 μ Ci of Pro-mix™ (Amersham), was added. The cell layers were incubated in this labeling
15 medium for two hours, washed with PBS, and incubated for 0, 15, 30, 60, 90, 120, or 180 minutes in serum-free DMEM (chase). After the various chase periods, the medium was collected, the cells were again washed two times in PBS, and lysed in immunoprecipitation buffer. The VEGF-C polypeptides were analyzed from both the culture medium and from the cell lysates by immunoprecipitation, using the VEGF-C-specific antiserum raised
20 against the NH₂-terminal peptide (PAM126) of the 23 kD VEGF-C form.

Immunoprecipitated polypeptides were analyzed via SDS-PAGE followed by autoradiography.

The resultant autoradiograms demonstrated that immediately after a 2 hour labeling (chase time 0), the VEGF-C vector-transfected cells contained a radioactive
25 polypeptide band of about 58kD (originally estimated to be about 55 kD, and re-evaluated to be about 58 kD using different size standards), which was not observed in mock-transfected cells (M). Most of this ~58 kD precursor undergoes dimerization. This ~58 kD polypeptide band gradually diminished in intensity with increasing chase periods. A 32 kD polypeptide band also is observed in VEGF-C transfected cells (but not mock-
30 transfected cells). This 32 kD band disappears from cells with similar kinetics to that of the ~58 kD band. Additional analysis indicated that the 32 kD band was a doublet of 29 kD and 31-32 kD forms, held together by disulfide bonds. Simultaneously, increasing

amounts of 32 kD and subsequently 23 kD and 14-15 kD polypeptides appeared in the medium.

Collectively, the data from the pulse-chase experiments indicate that the ~58 kD intracellular polypeptide represents a pro-VEGF-C polypeptide, which is proteolytically cleaved either intracellularly or at the cell surface into the 29 kD and 31-32 kD polypeptides. The 29/31 kD form is secreted and simultaneously further processed by proteolysis into the 23 kD and 14-15 kD forms. In additional experiments, disulfide linked dimers of the 29 kD and 15 kD forms were observed. Without intending to be limited to a particular theory, it is believed that processing of the VEGF-C precursor occurs as removal of a signal sequence, removal of the COOH-terminal domain (BR3P), and removal of an amino terminal polypeptide, resulting in a VEGF-C polypeptide having the TEE... amino terminus.

At high resolution, the 23 kD polypeptide band appears as a closely-spaced polypeptide doublet, suggesting heterogeneity in cleavage or glycosylation.

15

EXAMPLE 20

Isolation of Mouse and Quail cDNA Clones Encoding VEGF-C

A. Murine VEGF-C

To clone a murine VEGF-C, approximately 1×10^6 bacteriophage lambda clones of a commercially-available 12 day mouse embryonal cDNA library (lambda EXlox library, Novagen, catalog number 69632-1) were screened with a radiolabeled fragment of human VEGF-C cDNA containing nucleotides 495 to 1661 of SEQ ID NO: 7. One positive clone was isolated.

A 1323 bp *EcoRI/HindIII* fragment of the insert of the isolated mouse cDNA clone was subcloned into the corresponding sites of the pBluescript SK+ vector (Stratagene) and sequenced. The cDNA sequence of this clone was homologous to the human VEGF-C sequence reported herein, except that about 710 bp of 5'-end sequence present in the human clone was not present in the mouse clone.

For further screening of mouse cDNA libraries, a *HindIII-BstXI* (*HindIII* site is from the pBluescript SK+ polylinker) fragment of 881 bp from the coding region of the mouse cDNA clone was radiolabeled and used as a probe to screen two additional mouse cDNA libraries. Two additional cDNA clones from an adult mouse heart ZAP II

cDNA library (Stratagene, catalog number 936306) were identified. Three additional clones also were isolated from a mouse heart 5'-stretch-plus cDNA library in λ gt11 (Clontech Laboratories, Inc., catalog number ML5002b). Of the latter three clones, one was found to contain an insert of about 1.9 kb. The insert of this cDNA clone was subcloned into *Eco*RI sites of pBluescript SK+ vector and both strands of this clone were completely sequenced, resulting in the nucleotide and deduced amino acid sequences shown in SEQ ID NOs: 10 and 11.

It is contemplated that the polypeptide corresponding to SEQ ID NO: 11 is processed into a mature mouse VEGF-C protein, in a manner analogous to the processing of the human VEGF-C prepropeptide. Putative cleavage sites for the mouse protein are identified using procedures outlined above for identification of cleavage sites for the human VEGF-C polypeptide.

The foregoing results demonstrate the utility of polynucleotides of the invention for identifying and isolating polynucleotides encoding other non-human mammalian VEGF-C proteins. Such identified and isolated polynucleotides, in turn, can be expressed (using procedures similar to those described in preceding examples) to produce recombinant polypeptides corresponding to non-human mammalian forms of VEGF-C.

B. Quail VEGF-C

The mouse and human VEGF-C sequences were used to design probes for isolating a quail VEGF-C cDNA from a quail cDNA library. A fragment of the human VEGF-C cDNA comprising nucleotides 495-1670 of SEQ ID NO: 7 was obtained by PCR amplification, cloned into the pCRII vector (Invitrogen) according to the manufacturer's instructions, and amplified. The insert was isolated by *Eco*RI digestion and preparative gel electrophoresis and then labeled using radioactive dCTP and random priming. A cDNA library made from quail embryos of stage E-4 in pcDNA-1 vector (Invitrogen) was then screened using this probe. About 200,000 colonies were plated and filter replicas were hybridized with the radioactive probe. Nine positive clones were identified and secondarily plated. Two of the nine clones hybridized in secondary screening. The purified clones (clones 1 and 14) had approximately 2.7 kb *Eco*RI inserts. Both clones were amplified and then sequenced using the T7 and SP6 primers (annealing to the vector). In addition, an internal *Sph*I restriction endonuclease cleavage site was identified about 1.9 kb from the

T7 primer side of the vector and used for subcloning 5'- and 3'- *Sph*I fragments, followed by sequencing from the *Sph*I end of the subclones. The sequences obtained were identical from both clones and showed a high degree of similarity to the human VEGF-C coding region. Subsequently, walking primers were made in both directions and double-stranded sequencing was completed for 1743 base pairs, including the full-length open reading frame.

The cDNA sequence obtained includes a long open reading frame and 5' untranslated region. The DNA and deduced amino acid sequences for the quail cDNA are set forth in SEQ ID NOs: 12 and 13, respectively. Studies performed with the putative quail VEGF-C cDNA have shown that its protein product is secreted from transfected cells and interacts with avian VEGFR-3 and VEGFR-2, further confirming the conclusion that the cDNA encodes a quail VEGF-C protein. The proteins secreted from 293-EBNA cells transfected with quail VEGF-C cDNA were analyzed in immunoprecipitation studies using the VEGF-C-specific polyclonal antisera generated against the PAM126 polypeptide (Example 19). A doublet band of about 30-32 kD, and a band of about 22-23 kD, were immunoprecipitated from the transfected cells but not from control cells. These immunoprecipitation studies thus provide a further indication that VEGF-C from non-human species is processed (from a prepro-VEGF-C form) in a manner analogous to the processing of human VEGF-C. As shown in Fig. 5, the human, murine, and avian (quail) VEGF-C precursor amino acid sequences share a significant degree of conservation. This high degree of homology between species permits the isolation of VEGF-C encoding sequences from other species, especially vertebrate species, and more particularly mammalian and avian species, using polynucleotides of the present invention as probes and using standard molecular biological techniques such as those described herein.

EXAMPLE 21

N-terminal peptide sequence analyses of recombinant VEGF-C

Cells (293 EBNA) transfected with VEGF-C cDNA (see Example 13) secrete several forms of recombinant VEGF-C (Fig. 6A, lane IP). In the absence of alkylation, the three major, proteolytically-processed forms of VEGF-C migrate in SDS-PAGE as proteins with apparent molecular masses of 32/29 kD (doublet), 21 kD and 15 kD. Two minor polypeptides exhibit approximate molecular masses of 63 and 52 kD,

respectively. One of these polypeptides is presumably a glycosylated and non-processed form; the other polypeptide is presumably glycosylated and partially processed. More precise size measurements (using SDS-PAGE under reducing conditions) revealed that the molecular masses of the VEGF-C forms that were initially estimated as 63, 52, 32, 23, and 14 kD (using SDS-PAGE under reducing conditions and a different set of size standards) are approximately 58, 43, 31, 29, 21, and 15 kD, respectively (the initial measurements in most cases falling within acceptable 10% error of the more precise measurements).

To determine sites of proteolytic cleavage of the VEGF-C precursor, an immunoaffinity column was used to purify VEGF-C polypeptides from the conditioned medium of 293 EBNA cells transfected with VEGF-C cDNA. To prepare the immunoaffinity column, a rabbit was immunized with a synthetic peptide corresponding to amino acids 104-120 of SEQ ID NO: 8: H₂N-EETIKFAAAHYNTEILK (see PAM126 in Example 19). The IgG fraction was isolated from the serum of the immunized rabbit using protein A Sepharose (Pharmacia). The isolated IgG fraction was covalently bound to CNBr-activated Sepharose CL-4B (Pharmacia) using standard techniques at a concentration of 5 mg IgG/ml of Sepharose. This immunoaffinity matrix was used to isolate processed VEGF-C from 1.2 liters of the conditioned medium (CM).

The purified material eluted from the column was analyzed by gel electrophoresis and Western blotting. Fractions containing VEGF-C polypeptides were combined, dialyzed against 10 mM Tris HCl, vacuum-dried, electrotransferred to Immobilon-P (polyvinylidene difluoride or PVDF) transfer membrane (Millipore, Marlborough, MA) and subjected to N-terminal amino acid sequence analysis.

The polypeptide band of 32 kD yielded two distinct sequences: NH₂-FESGLDLSDA... and NH₂-AVVMTQTPAS... (SEQ ID NO: 14), the former corresponding to the N-terminal part of VEGF-C after cleavage of the signal peptide, starting from amino acid 32 (SEQ ID NO: 8), and the latter corresponding to the kappa-chain of IgG, which was present in the purified material due to "leakage" of the affinity matrix during the elution procedure.

In order to obtain the N-terminal peptide sequence of the 29 kD form of VEGF-C, a construct (VEGF-C NHis) encoding a VEGF-C mutant was generated. In particular, the construct encoded a VEGF-C mutant that fused a 6xHis tag to the N-terminus of the secreted precursor (*i.e.*, between amino acids 31 and 33 in SEQ ID NO:

8). The phenylalanine at position 32 was removed to prevent possible cleavage of the tag sequence during secretion of VEGF-C. The VEGF-C NHis construct was cloned into pREP7 as a vector; the construction is described more fully in Example 28, below.

The calcium phosphate co-precipitation technique was used to transfect
5 VEGF-C NHis into 293 EBNA cells. Cells were incubated in DMEM/10% fetal calf serum in 15 cm cell culture dishes (a total of 25 plates). The following day, the cells were reseeded into fresh culture dishes (75 plates) containing the same medium and incubated for 48 hours. Cell layers were then washed once with PBS and DMEM medium lacking FCS was added. Cells were incubated in this medium for 48 hours and the medium was
10 collected, cleared by centrifugation at 5000 x g and concentrated 500X using an Ultrasette Tangential Flow Device (Filtron, Northborough, MA), as described in Example 5 above. VEGF-C NHis was purified from the concentrated conditioned medium using TALON™ Metal Affinity Resin (Clontech Laboratories, Inc.) and the manufacturer's protocol for native protein purification using imidazole-containing buffers. The protein was eluted with
15 a solution containing 20 mM Tris-HCl (pH 8.0), 100 mM NaCl, and 200 mM imidazole. The eluted fractions containing purified VEGF-C NHis were detected by immunoblotting with Antiserum 882 (antiserum from rabbit 882, immunized with the PAM-126 polypeptide). Fractions containing VEGF-C NHis were combined, dialyzed and vacuum-dried. Due to the presence of the 6xHis tag at the N-terminus of this form of VEGF-C,
20 the upper component of the major doublet of the VEGF-C NHis migrates slightly slower than the 32 kD form of wild type VEGF-C, thereby improving the separation of the VEGF-C NHis 32 kD mutant from the 29 kD band using SDS-PAGE. Approximately 15 µg of the purified VEGF-C were subjected to SDS-PAGE under reducing conditions, electrotransferred to Immobilon-P (PVDF) transfer membrane (Millipore, Inc.,
25 Marlborough, MA) and the band at 29 kD was subjected to N-terminal amino acid sequence analysis. This sequence analysis revealed an N-terminal sequence of H₂N-SLPAT . . . , corresponding to amino acids 228-232 of VEGF-C (SEQ ID NO: 8).

The polypeptide band of 21 kD yielded the sequence H₂N-AHYNTEILKS . . . , corresponding to an amino-terminus starting at amino acid 112 of SEQ ID NO: 8.
30 Thus, the proteolytic processing site which results in the 21 kD form of VEGF-C produced by transfected 293 EBNA cells apparently occurs nine amino acid residues

downstream of the cleavage site which results in the 23 kD form of VEGF-C secreted by PC-3 cells.

The N-terminus of the 15 kD form was identical to the N-terminus of the 32 kD form (NH₂-FESGLDLSDA...). The 15 kD form was not detected when 5 recombinant VEGF-C was produced by COS cells. This suggests that production of this form is cell lineage specific.

Example 22

Dimeric and monomeric forms of VEGF-C

The composition of VEGF-C dimers was analyzed as follows. Cells (293 10 EBNA cells), transfected with the pREP7 VEGF-C vector as described in Example 11, were metabolically labeled with Pro-mix L-[³⁵S] labeling mix (Amersham Corp.) to a final concentration of 100 µCi/ml.

In parallel, a VEGF-C mutant, designated "R102S", was prepared and analyzed. To prepare the DNA encoding VEGF-C-R102S, the arginine codon at position 15 102 of SEQ ID NO: 8 was replaced with a serine codon. This VEGF-C-R102S-encoding DNA, in a pREP7 vector, was transfected into 293 EBNA cells and expressed as described above. VEGF-C polypeptides were immunoprecipitated using antisera 882 (obtained by immunization of a rabbit with a polypeptide corresponding to residues 104-120 of SEQ ID NO: 8 (see previous Example)) and antisera 905 (obtained by immunization of a rabbit 20 with a polypeptide corresponding to a portion of the pro-VEGF-C leader: H₂N-ESGLDLSDAEPDAGEATAYASK (residues 33 to 54 of SEQ ID NO: 8).

The immunoprecipitates from each cell culture were subjected to SDS-PAGE under non-denaturing conditions (Fig. 6B). Bands 1-6 were cut out from the gel, 25 soaked for 30 minutes in 1x gel-loading buffer containing 200 mM β-mercaptoethanol, and individually subjected to SDS-PAGE under denaturing conditions (Figs. 6A and 6C, lanes 1-6).

As can be seen from Figures 6A-C, each high molecular weight form of VEGF-C (Fig. 6B, bands 1-4) consists of at least two monomers bound by disulfide bonds 30 (Compare Figs. 6A and 6C, lanes 1-4, in the reducing gels). The main component of bands 1-3 is the doublet of 32/29 kD, where both proteins are present in an equimolar

ratio. The main fraction of the 21 kD form is secreted as either a monomer or as a homodimer connected by means other than disulfide bonds (bands 6 and lanes 6 in Figs. 6A-C).

The R102S mutation creates an additional site for N-linked glycosylation in VEGF-C at the asparagine residue at position 100 in SEQ ID NO: 8. Glycosylation at this additional glycosylation site increases the apparent molecular weight of polypeptides containing the site, as confirmed in Figures 6A-C and Figures 7A-B. The additional glycosylation lowers the mobility of forms of VEGF-C-R102S that contain the additional glycosylation site, when compared to polypeptides of similar primary structure corresponding to VEGF-C. Figures 6A-C and Figures 7A-B reveal that the VEGF-C-R102S polypeptides corresponding to the 32 kD and 15 kD forms of wt VEGF-C exhibit increased apparent molecular weights, indicating that each of these polypeptides contains the newly introduced glycosylation site. In particular, the VEGF-C-R102S polypeptide corresponding to the 15 kD polypeptide from VEGF-C comigrates on a gel with the 21 kD form of the wild type (wt) VEGF-C, reflecting a shift on the gel to a position corresponding to a greater apparent molecular weight. (Compare lanes 4 in Figures 6A and 6C). The mobility of the 58 kD form of VEGF-C was slowed to 64 kD by the R102S mutation, indicating that this form contains the appropriate N-terminal peptide of VEGF-C. The mobilities of the 21, 29, and 43 kD forms were unaffected by the R102S mutation, suggesting that these polypeptides contain peptide sequences located C-terminally of R₁₀₂.

In a related experiment, another VEGF-C mutant, designated "R226,227S," was prepared and analyzed. To prepare a DNA encoding VEGF-C-R226,227S, the arginine codons at positions 226 and 227 of SEQ ID NO: 8 were replaced with serine codons by site-directed mutagenesis. The resultant DNA was transfected into 293 EBNA cells as described above and expressed and analyzed under the same conditions as described for VEGF-C and VEGF-C-R102S. In the conditioned medium from the cells expressing VEGF-C-R226,227S, no 32 kD form of VEGF-C was detected. These results indicate that a C-terminal cleavage site of wild-type VEGF-C is adjacent to residues 226 and 227 of SEQ ID NO: 8, and is destroyed by the mutation of the arginines to serines. Again, the mobility of the 29 kD component of the doublet was unchanged (Figures 7A-B).

Taken together, these data indicate that the major form of the processed VEGF-C is a heterodimer consisting of (1) a polypeptide of 32 kD containing amino acids 32-227 of the prepro-VEGF-C (amino acids 32 to 227 in SEQ ID NO: 8) attached by disulfide bonds to (2) a polypeptide of 29 kD beginning with amino acid 228 in SEQ ID NO: 8. These data are also supported by a comparison of the pattern of immunoprecipitated, labeled VEGF-C forms using antisera 882 and antisera 905.

When VEGF-C immunoprecipitation was carried out using conditioned medium, both antisera (882 and 905) recognized some or all of the three major processed forms of VEGF-C (32/29 kD, 21 kD and 15 kD). When the conditioned medium was reduced by incubation in the presence of 10 mM dithiothreitol for two hours at room temperature with subsequent alkylation by additional incubation with 25 mM iodoacetamide for 20 minutes at room temperature, neither antibody precipitated the 29 kD component, although antibody 882 still recognized polypeptides of 32 kD, 21 kD and 15 kD. In subsequent experiments it was observed that neither antibody was capable of immunoprecipitating the 43 kD form. These results are consistent with the nature of the oligopeptide antigen used to elicit the antibodies contained in antisera 882, an oligopeptide containing amino acid residues 104-120 of SEQ ID NO: 8. On the other hand, antisera 905 recognized only the 32 kD and 15 kD polypeptides, which include sequence of the oligopeptide (amino acids 33 to 54 of SEQ ID NO: 8) used for immunization to obtain antisera 905. Taking into account the mobility shift of the 32 kD and 15 kD forms, the immunoprecipitation results with the R102S mutant were similar (Figs. 8A-B). The specificity of antibody 905 is confirmed by the fact that it did not recognize a VEGF-C Δ N form wherein the N-terminal propeptide spanning residues 32-102 of the unprocessed polypeptide had been deleted (Fig. 8B).

The results of these experiments also demonstrate that the 21 kD polypeptide is found (1) in heterodimers with other molecular forms (see Figs. 6A-C and Figs. 7A-B), and (2) secreted as a monomer or a homodimer held by bonds other than disulfide bonds (Figs. 6A and 6B, lanes 6).

The experiments disclosed in this example demonstrate that several forms of VEGF-C exist. A variety of VEGF-C monomers were observed and these monomers can vary depending on the level and pattern of glycosylation. In addition, VEGF-C was observed as a multimer, for example a homodimer or a heterodimer. The processing of

VEGF-C is schematically presented in Fig. 9 (disulfide bonds not shown). All forms of VEGF-C are within the scope of the present invention.

Example 23

In situ Hybridization of Mouse Embryos

5 To analyze VEGF-C mRNA distribution in different cells and tissues, sections of 12.5 and 14.5-day post-coitus (p.c.) mouse embryos were prepared and analyzed via *in situ* hybridization using labeled VEGF-C probes. *In situ* hybridization of tissue sections was performed as described in Västrik *et al.*, *J. Cell Biol.*, 128:1197-1208 (1995). A mouse VEGF-C antisense RNA probe was generated from linearized

10 pBluescript II SK+ plasmid (Stratagene Inc., La Jolla, CA), containing a cDNA fragment corresponding to nucleotides 499-979 of a mouse VEGF-C cDNA (SEQ ID NO: 10). Radiolabeled RNA was synthesized using T7 polymerase and [³⁵S]-UTP (Amersham). Mouse VEGF-B antisense and sense RNA probes were synthesized in a similar manner from linearized pCRII plasmid containing the mouse VEGF-B cDNA insert as described

15 Olofsson *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 93:2576-2581 (1996). The high stringency wash was for 45 minutes at 65°C in a solution containing 30 mM dithiothreitol (DTT) and 4 x SSC. The slides were exposed for 28 days, developed and stained with hematoxylin. For comparison, similar sections were hybridized with a VEGFR-3 probe and the 12.5-day p.c. embryos were also probed for VEGF-B mRNA.

20 Darkfield and lightfield photomicrographs from these experiments are presented in commonly-owned PCT patent application PCT/FI96/00427, filed August 01, 1996, published as WO 97/05250, incorporated by reference herein. Observations from the photomicrographs are summarized below. In a 12.5 day p.c. embryo, a parasagittal section revealed that VEGF-C mRNA was particularly prominent in the mesenchyme

25 around the vessels surrounding the developing metanephros. In addition, hybridization signals were observed between the developing vertebrae, in the developing lung mesenchyme, in the neck region and developing forehead. The specificity of these signals was evident from the comparison with VEGF-B expression in an adjacent section, where the myocardium gave a very strong signal and lower levels of VEGF-B mRNA were

30 detected in several other tissues. Both genes appear to be expressed in between the

developing vertebrae, in the developing lung, and forehead. Hybridization of the VEGF-C sense probe showed no specific expression within these structures.

Studies also were conducted of the expression patterns of VEGF-C and VEGFR-3 in 12.5 day p.c. mouse embryos in the jugular region, where the developing dorsal aorta and cardinal vein are located. This is the area where the first lymphatic vessels sprout from venous sac-like structures according to the long-standing theory of Sabin, *Am. J. Anat.*, 9:43-91 (1909). An intense VEGF-C signal was detected in the mesenchyme surrounding the developing venous sacs which also were positive for VEGFR-3.

10 The mesenterium supplies the developing gut with blood and contains developing lymphatic vessels. The developing 14.5 day p.c. mesenterium is positive for VEGF-C mRNA, with particularly high expression in connective tissue surrounding certain vessels. The adjacent mesenterial VEGFR-3 signals that were observed originate from small capillaries of the mesenterium. Therefore, there appears to be a paracrine
15 relationship between the production of the mRNAs for VEGF-C and its receptor. This data indicates that VEGF-C is expressed in a variety of tissues. Moreover, the pattern of expression is consistent with a role for VEGF-C in venous and lymphatic vessel development. Further, the data reveals that VEGF-C is expressed in non-human animals.

Example 24

20 Analysis of VEGF, VEGF-B, and VEGF-C mRNA Expression in Fetal and Adult Tissues

A human fetal tissue Northern blot containing 2 μ g of polyadenylated RNAs from brain, lung, liver and kidney (Clontech Inc.) was hybridized with a pool of the following probes: a human full-length VEGF-C cDNA insert (Genbank Acc. No. X94216),
25 a human VEGF-B₁₆₇ cDNA fragment (nucleotides 1-382, Genbank Acc. No. U48800) obtained by PCR amplification; and a human VEGF 581 bp cDNA fragment covering base pairs 57-638 (Genbank Acc. No. X15997). Blots were washed under stringent conditions, using techniques standard in the art.

Mouse embryo multiple tissue Northern blot (Clontech Inc.) containing 2
30 μ g of polyadenylated RNAs from 7, 11, 15 and 17 day postcoital (p.c.) embryos was hybridized with mouse VEGF-C cDNA fragment (base pairs 499-656). A mouse adult

tissue Northern blot was hybridized with the probes for human VEGF, VEGF-B₁₆₇, VEGF-C and with a VEGFR-3 cDNA fragment (nucleotides 1-595; Genbank Acc. No. X68203).

In adult mouse tissues, both 2.4 kb and 2.0 kb mRNA signals were observed with the VEGF-C probe, at an approximately 4:1 ratio. The most conspicuous signals were obtained from lung and heart RNA, while kidney, liver, brain, and skeletal muscle had lower levels, and spleen and testis had barely visible levels. As in the human tissues, VEGF mRNA expression in adult mice was most abundant in lung and heart RNA, whereas the other samples showed less coordinate regulation with VEGF-C expression. Skeletal muscle and heart tissues gave the highest VEGF-B mRNA levels from adult mice, as previously reported Olofsson *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 93:2576-2581 (1996). Comparison with VEGFR-3 expression showed that the tissues where VEGF-C is expressed also contain mRNA for its cognate receptor tyrosine kinase, although in the adult liver VEGFR-3 mRNA was disproportionately abundant.

To provide a better insight into the regulation of the VEGF-C mRNA during embryonic development, polyadenylated RNA isolated from mouse embryos of various gestational ages (7, 11, 15, and 17 day p.c.) was hybridized with the mouse VEGF-C probe. These analyses showed that the amount of 2.4 kb VEGF-C mRNA is relatively constant throughout the gestational period.

20

Example 25

Regulation of mRNAs for VEGF family members by serum, interleukin-1 and dexamethasone in human fibroblasts in culture

Human IMR-90 fibroblasts were grown in DMEM medium containing 10% FCS and antibiotics. The cells were grown to 80% confluence, then starved for 48 hours in 0.5 % FCS in DMEM. Thereafter, the growth medium was changed to DMEM containing 5% FCS, with or without 10 ng/ml interleukin-1 (IL-1) and with or without 1 mM dexamethasone. The culture plates were incubated with these additions for the times indicated, and total cellular RNA was isolated using the TRIZOL kit (GIBCO-BRL). About 20 µg of total RNA from each sample was electrophoresed in 1.5% formaldehyde-agarose gels as described in Sambrook *et al.*, *supra* (1989). The gel was used for

Northern blotting and hybridization with radiolabeled insert DNA from the human VEGF clone (a 581 bp cDNA covering bps 57-638, Genbank Acc. No. 15997) and a human VEGF-B₁₆₇ cDNA fragment (nucleotides 1-382, Genbank Acc. No. U48800).

Subsequently, the Northern blots were probed with radiolabeled insert from the VEGF-C 5 cDNA plasmid. Primers were labeled using a standard technique involving enzymatic extension reactions of random primers, as would be understood by one of ordinary skill in the art.

The Northern blot analyses revealed that very low levels of VEGF-C and VEGF are expressed by the starved IMR-90 cells as well as cells after 1 hour of 10 stimulation. In contrast, abundant VEGF-B mRNA signal was visible under these conditions. After 4 hours of serum stimulation, there was a strong induction of VEGF-C and VEGF mRNAs, which were further increased in the IL-1 treated sample. The effect of IL-1 seemed to be abolished in the presence of dexamethasone. A similar pattern of enhancement was observed in the 8 hour sample, but a gradual down-regulation of all 15 signals was observed for both RNAs in the 24 hour and 48 hour samples. In contrast, VEGF-B mRNA levels remained constant and thus showed remarkable stability throughout the time period. The results are useful in guiding efforts to use VEGF-C and its fragments, its antagonists, and anti-VEGF-C antibodies in methods for treating a variety of disorders.

20

Example 26

Expression and analysis of recombinant murine VEGF-C

The mouse VEGF-C cDNA was expressed as a recombinant protein and the secreted protein was analyzed for its receptor binding properties. The binding of mouse VEGF-C to the human VEGFR-3 extracellular domain was studied by using media 25 from Bosc23 cells transfected with mouse VEGF-C cDNA in a retroviral expression vector.

The 1.8 kb mouse VEGF-C cDNA was cloned as an *EcoRI* fragment into the retroviral expression vector pBabe-puro containing the SV40 early promoter region [Morgenstern *et al.*, *Nucl. Acids Res.*, 18:3587-3595 (1990)], and transfected into the 30 Bosc23 packaging cell line [Pearet *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 90:8392-8396 (1994)] by the calcium-phosphate precipitation method. For comparison, Bosc23 cells

also were transfected with the previously-described human VEGF-C construct in the pREP7 expression vector. The transfected cells were cultured for 48 hours prior to metabolic labeling. Cells were changed into DMEM medium devoid of cysteine and methionine, and, after 45 minutes of preincubation and medium change, Pro-mixTM L-[³⁵S] 5 *in vitro* cell labeling mix (Amersham Corp.), in the same medium, was added to a final concentration of about 120 μ Ci/ml. After 6 hours of incubation, the culture medium was collected and clarified by centrifugation.

For immunoprecipitation, 1 ml aliquots of the media from metabolically-labeled Bosc23 cells transfected with empty vector or mouse or human recombinant 10 VEGF-C, respectively, were incubated overnight on ice with 2 μ l of rabbit polyclonal antiserum raised against an N-terminal 17 amino acid oligopeptide of mature human VEGF-C (H₂N-EETIKFAAAHYNTEILK) (SEQ ID NO: 8, residues 104-120). Thereafter, the samples were incubated with protein A sepharose for 40 minutes at 4°C with gentle agitation. The sepharose beads were then washed twice with 15 immunoprecipitation buffer and four times with 20 mM Tris-HCl, pH 7.4. Samples were boiled in Laemmli buffer and analyzed by 12.5% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE).

Immunoprecipitation of VEGF-C from media of transfected and metabolically-labeled cells revealed bands of approximately 30-32x10³ M_r (a doublet) and 20 22-23x10³ M_r in 12.5% SDS-PAGE. These bands were not detected in samples from nontransfected or mock-transfected cells. These results show that antibodies raised against human VEGF-C recognize the corresponding mouse ligand, and provide an indication that the proteolytic processing that occurs to produce murine VEGF-C is analogous to the processing that occurs to produce human VEGF-C.

25 For receptor binding experiments, 1 ml aliquots of media from metabolically-labeled Bosc23 cells were incubated with VEGFR-3 extracellular domain (see Example 3), covalently coupled to sepharose, for 4 hours at 4°C with gentle mixing. The sepharose beads were washed four times with ice-cold phosphate buffered saline (PBS), and the samples were analyzed by gel electrophoresis as described in Joukov *et al.*, 30 *EMBO J.*, 15:290-298 (1996).

Similar 30-32 x 10³ M_r doublet and 22-23 x 10³ M_r polypeptide bands were obtained in the receptor binding assay as compared to the immunoprecipitation assay.

- 65 -

Thus, mouse VEGF-C binds to human VEGFR-3. The slightly faster mobility of the mouse VEGF-C polypeptides that was observed may be caused by the four amino acid residue difference observed in sequence analysis (residues H88-E91, Fig. 10).

The capacity of mouse recombinant VEGF-C to induce VEGFR-3
5 autophosphorylation was also investigated. For the VEGFR-3 receptor stimulation experiments, subconfluent NIH 3T3-Flt4 cells, Pajusola *et al.*, *Oncogene*, 9:3545-3555 (1994), were starved overnight in serum-free medium containing 0.2% BSA. In general, the cells were stimulated with the conditioned medium from VEGF-C vector-transfected cells for 5 minutes, washed three times with cold PBS containing 200 μ M vanadate, and
10 lysed in RIPA buffer for immunoprecipitation analysis. The lysates were centrifuged for 25 minutes at 16000 x g and the resulting supernatants were incubated for 2 hours on ice with the specific antisera, followed by immunoprecipitation using protein A-sepharose and analysis in 7% SDS-PAGE. Polypeptides were transferred to nitrocellulose and analyzed by immunoblotting using anti-phosphotyrosine (Transduction Laboratories) and anti-
15 receptor antibodies, as described by Pajusola *et al.*, *Oncogene*, 9:3545-3555 (1994). Filter stripping was carried out at 50°C for 30 minutes in 100 mM 2-mercaptoethanol, 2% SDS, 62.5 mM Tris-HCl, pH 6.7, with occasional agitation. The results of the experiment demonstrated that culture medium containing mouse VEGF-C stimulates the autophosphorylation of VEGFR-3 to a similar extent as human baculoviral VEGF-C or the
20 tyrosyl phosphatase inhibitor pervanadate.

Mouse VEGF-C appeared to be a potent inducer of VEGFR-3 autophosphorylation, with the 195×10^3 M_r precursor and proteolytically-cleaved 125×10^3 M_r tyrosine kinase polypeptides of the receptor (Pajusola *et al.*, *Oncogene*, 9:3545-3555 (1994)), being phosphorylated.

25 VEGFR-2 stimulation was studied in subconfluent porcine aortic endothelial (PAE) cells expressing KDR (VEGFR-2) (PAE-VEGFR-2) [Waltenberger *et al.*, *J. Biol. Chem.*, 269:26988-26995 (1994)], which were starved overnight in serum-free medium containing 0.2% BSA. Stimulation was carried out and the lysates prepared as described above. For receptor immunoprecipitation, specific antiserum for VEGFR-2
30 [Waltenberger *et al.*, *J. Biol. Chem.*, 269:26988-26995 (1994)] was used. The immunoprecipitates were analyzed as described for VEGFR-3 in 7% SDS-PAGE followed by Western blotting with anti-phosphotyrosine antibodies, stripping of the filter, and re-

- 66 -

probing it with anti-VEGFR-2 antibodies (Santa Cruz). VEGFR-2 stimulation was first tried with unconcentrated medium from cells expressing recombinant VEGF-C, but immunoblotting analysis did not reveal any receptor autophosphorylation.

To further determine whether mouse recombinant VEGF-C can also induce VEGFR-2 autophosphorylation as observed for human VEGF-C, PAE cells expressing VEGFR-2 were stimulated with tenfold concentrated medium from cultures transfected with mouse VEGF-C expression vector and autophosphorylation was analyzed. For comparison, cells treated with tenfold concentrated medium containing human recombinant VEGF-C (Joukov *et al.*, (1996)), unconcentrated medium from human VEGF-C baculovirus infected insect cells, or pervanadate (a tyrosyl phosphatase inhibitor) were used. In response to human baculoviral VEGF-C as well as pervanadate treatment, VEGFR-2 was prominently phosphorylated, whereas human and mouse recombinant VEGF-C gave a weak and barely detectable enhancement of autophosphorylation, respectively. Media from cell cultures transfected with empty vector or VEGF-C cloned in the antisense orientation did not induce autophosphorylation of VEGFR-2. Therefore, mouse VEGF-C binds to VEGFR-3 and activates this receptor at a much lower concentration than needed for the activation of VEGFR-2. Nevertheless, the invention comprehends methods for using the materials of the invention to take advantage of the interaction of VEGF-C with VEGFR-2, in addition to the interaction between VEGF-C and VEGFR-3.

Example 27

VEGF-C E104-S213 fragment expressed in *Pichia* yeast stimulates autophosphorylation of Flt4 (VEGFR-3) and KDR (VEGFR-2)

A truncated form of human VEGF-C cDNA was constructed wherein (1) the sequence encoding residues of a putative mature VEGF-C amino terminus H₂N-E(104)ETIK (SEQ ID NO: 8, residues 104 et seq.) was fused in-frame to the yeast PHO1 signal sequence (Invitrogen *Pichia* Expression Kit, Catalog #K1710-01), and (2) a stop codon was introduced after amino acid 213 (H₂N- ...RCMS; *i.e.*, after codon 213 of SEQ ID NO: 7). The resultant truncated cDNA construct was then inserted into the *Pichia pastoris* expression vector pHIL-S1 (Invitrogen). For the cloning, an internal *Bgl*III site in

- 67 -

the VEGF-C coding sequence was mutated without change of the encoded polypeptide sequence.

This VEGF-C expression vector was then transfected into *Pichia* cells and positive clones were identified by screening for the expression of VEGF-C protein in the culture medium by Western blotting. One positive clone was grown in a 50 ml culture, and induced with methanol for various periods of time from 0 to 60 hours. About 10 μ l of medium was analyzed by gel electrophoresis, followed by Western blotting and detection with anti-VEGF-C antiserum, as described above. An approximately 24 kD polypeptide (band spreading was observed due to glycosylation) accumulated in the culture medium of cells transfected with the recombinant VEGF-C construct, but not in the medium of mock-transfected cells or cells transfected with the vector alone.

The medium containing the recombinant VEGF-C protein was concentrated by Centricon 30 kD cutoff ultrafiltration and used to stimulate NIH 3T3 cells expressing Flt4 (VEGFR-3) and porcine aortic endothelial (PAE) cells expressing KDR (VEGFR-2). The stimulated cells were lysed and immunoprecipitated using VEGFR-specific antisera and the immunoprecipitates were analyzed by Western blotting using anti-phosphotyrosine antibodies, chemiluminescence, and fluorography. As a positive control for maximal autophosphorylation of the VEGFRs, vanadate (VO_4) treatment of the cells for 10 minutes was used. Medium from *Pichia* cultures secreting the recombinant VEGF-C polypeptide induced autophosphorylation of both Flt4 polypeptides of 195 kD and 125 kD as well as the KDR polypeptide of about 200 kD. Vanadate, on the other hand, induces heavy tyrosyl phosphorylation of the receptor bands in addition to other bands probably coprecipitating with the receptors.

These results demonstrate that a VEGF-homologous domain of VEGF-C consisting of amino acid residues 104E - 213S (SEQ ID NO: 8, residues 104-213) can be recombinantly produced in yeast and is capable of stimulating the autophosphorylation of Flt4 (VEGFR-3) and KDR (VEGFR-2). Recombinant VEGF-C fragments such as the fragment described herein, which are capable of stimulating Flt4 or KDR autophosphorylation are intended as aspects of the invention; methods of using these fragments are also within the scope of the invention.

Example 28**Properties of the differentially processed forms of VEGF-C**

The following oligonucleotides were used to generate a set of VEGF-C variants and analogs:

- 5 5'- TCTCTTCTGTGCTTGAGTTGAG -3' (SEQ ID NO: 15), used to generate VEGF-C R102S (arginine mutated to serine at position 102 (SEQ ID NO: 8));
- 5'-TCTCTTCTGTCCCTGAGTTGAG -3' (SEQ ID NO: 16), used to generate VEGF-C R102G (arginine mutated to glycine at position 102 (SEQ ID NO: 8));
- 5'-TGTGCTGCAGCAAATTTTATAGTCTCTTCTGTGGCGGCGGC
- 10 GGCGGCGGGCGCCTCGCGAGGACC -3' (SEQ ID NO: 17), used to generate VEGF-C ΔN (deletion of N-terminal propeptide corresponding to amino acids 32-102 (SEQ ID NO: 8));
- 5'- CTGGCAGGGAAGTCTAATAATGGAATGAA - 3' (SEQ ID NO: 18), used to generate VEGF-C R226,227S (arginine codons mutated to serines at positions 226 and
- 15 227 (SEQ ID NO: 8));
- 5'-GGGCTCCGCGTCCGAGAGGTCGAGTCCGGACTCGTGATGGT
- GATGGTGATGGGCGGCGGCGGCGGGCGCCTCGCGAGGACC -3' (SEQ ID NO: 19), used to generate VEGF-C NHis (this construct encodes a polypeptide with a 6xHis tag fused to approximately the N-terminus of the secreted precursor, as described in
- 20 Example 21 (amino acid 33 of SEQ ID NO: 8)).

Some of the foregoing VEGF-C mutant constructs were further modified to obtain additional constructs. For example, VEGF-C R102G in pALTER (Promega) and oligonucleotide 5'-GTATTATAATGTCCTCCACCAAATTTTATAG -3' (SEQ ID NO: 20) were used to generate VEGF-C 4G, which encodes a polypeptide with four point

25 mutations: R102G, A110G, A111G, and A112G (alanines mutated to glycines at positions 110-112 (SEQ ID NO: 8). These four mutations are adjacent to predicted sites of cleavage of VEGF-C expressed in PC-3 and recombinantly expressed in 293 EBNA cells.

Another construct was created using VEGF-C ΔN and oligonucleotide 5'-

30 GTTCGCTGCCTGACACTGTGGTAGTGTTGCTGGC

 GGCCGCTAGTGATGGTGATGGTGATGAATAATGGAATGAACTTGTCTGTAAAC

 ATCCAG -3' (SEQ ID NO: 21) to generate VEGF-C ΔNΔCHis. This construct encodes

- 69 -

a polypeptide with a deleted N-terminal propeptide (amino acids 32-102); a deleted C-terminal propeptide (amino acids 226-419 of SEQ ID NO: 8); and an added 6xHis tag at the C-terminus (see SEQ ID NO: 59).

All constructs were further digested with *HindIII* and *NotI*, subcloned into
5 *HindIII/NotI* digested pREP7 vector, and used to transfect 293 EBNA cells. About 48 hours after transfection, the cells were either metabolically labelled with Pro-mix™ as described above, or starved in serum-free medium for 2 days. Media were then collected and used in subsequent experiments. Wild type (wt) VEGF-C, VEGF-C NHis and VEGF-C ΔNΔCHis were expressed to similar levels in 293 EBNA cells. At the same time,
10 expression of the VEGF-C 4G polypeptide was considerably lower, possibly due to the changed conformation and decreased stability of the translated product. However, all the above VEGF-C mutants were secreted from the cells.

The conditioned media from the transfected and starved cells were concentrated 5-fold and used to assess their ability to stimulate tyrosine phosphorylation of
15 Flt4 (VEGFR-3) expressed in NIH 3T3 cells and KDR (VEGFR-2) expressed in PAE cells. Wild type (wt) VEGF-C, as well as all three mutant polypeptides, stimulated tyrosine phosphorylation of VEGFR-3. The most prominent stimulation observed was by the short mature VEGF-C ΔNΔCHis. This mutant, as well as VEGF-C NHis, also stimulated tyrosine phosphorylation of VEGFR-2. Thus, despite the fact that a major
20 component of secreted recombinant VEGF-C is a dimer of 32/29 kD, the active part of VEGF-C responsible for its binding to VEGFR-3 and VEGFR-2 is localized between amino acids 102 and 226 (SEQ ID NO: 8) of the VEGF-C precursor. Analysis and comparison of binding properties and biological activities of these VEGF-C proteins and mutants, using assays such as those described herein, will provide data concerning the
25 significance of the observed major 32/29 kD and 21-23 kD VEGF-C processed forms. The data indicate that constructs encoding amino acid residues 103-225 of the VEGF-C precursor (SEQ ID NO: 8) generate a recombinant ligand that is functional for both VEGFR-3 and VEGFR-2.

The data from this and preceding examples demonstrate that numerous
30 fragments of the VEGF-C polypeptide retain biological activity. A naturally occurring VEGF-C polypeptide spanning amino acids 103-226 (or 103-227) of SEQ ID NO: 8, produced by a natural processing cleavage defining the C-terminus, has been shown to be

active. Example 27 demonstrates that a fragment with residues 104-213 of SEQ ID NO: 8 retains biological activity.

In addition, data from Example 21 demonstrates that a VEGF-C polypeptide having its amino terminus at position 112 of SEQ ID NO: 8 retains activity. 5 Additional experiments have shown that a fragment lacking residues 1-112 of SEQ ID NO: 8 retains biological activity.

In a related experiment, a stop codon was substituted for the lysine at position 214 of SEQ ID NO: 8 (SEQ ID NO: 7, nucleotides 991-993). The resulting recombinant polypeptide still was capable of inducing Flt4 autophosphorylation, indicating 10 that a polypeptide spanning amino acid residues 113-213 of SEQ ID NO: 8 is biologically active.

Sequence comparisons of members of the VEGF family of polypeptides provides an indication that still smaller fragments of the polypeptide depicted in SEQ ID NO: 8 will retain biological activity. In particular, eight highly conserved cysteine residues 15 of the VEGF family of polypeptides define a region from residues 131 - 211 of SEQ ID NO: 8 (see Figure 10) of evolutionary significance; therefore, a polypeptide spanning from about residue 131 to about residue 211 is expected to retain VEGF-C biological activity. In fact, a polypeptide which retains the conserved motif RCXXCC (e.g., a polypeptide comprising from about residue 161 to about residue 211 of SEQ ID NO: 8 is postulated to 20 retain VEGF-C biological activity. To maintain native conformation of these fragments, it may be preferred to retain about 1-2 additional amino acids at the carboxy-terminus and 1-2 or more amino acids at the amino terminus.

Beyond the preceding considerations, evidence exists that smaller fragments and/or fragment analogs which lack the conserved cysteines nonetheless will 25 retain VEGF-C biological activity. Consequently, the materials and methods of the invention include all VEGF-C fragments, variants, and analogs that retain at least one biological activity of VEGF-C, regardless of the presence or absence of members of the conserved set of cysteine residues.

Example 29**Expression of human VEGF-C under the human K14 keratin promoter in transgenic mice induces abundant growth of lymphatic vessels in the skin**

5 The Flt4 receptor tyrosine kinase is relatively specifically expressed in the endothelia of lymphatic vessels. Kaipainen *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 92: 3566-3570 (1995). Furthermore, the VEGF-C growth factor stimulates the Flt4 receptor, showing less activity towards the KDR receptor of blood vessels (Joukov *et al.*, *EMBO J.*, 15: 290-298 (1996); See Example 26).

10 Experiments were conducted in transgenic mice to analyze the specific effects of VEGF-C overexpression in tissues. The human K14 keratin promoter is active in the basal cells of stratified squamous epithelia (Vassar *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 86:1563-1567 (1989)) and was used as the expression control element in the recombinant VEGF-C transgene. The vector containing the K14 keratin promoter is
15 described in Vassar *et al.*, *Genes Dev.*, 5:714-727 (1991) and Nelson *et al.*, *J. Cell Biol.* 97:244-251 (1983).

 The recombinant VEGF-C transgene was constructed using the human full length VEGF-C cDNA (GenBank Acc. No. X94216). This sequence was excised from a pCI-neo vector (Promega) with *XhoI/NotI*, and the resulting 2027 base pair fragment
20 containing the open reading frame and stop codon (nucleotides 352-1611 of SEQ ID NO: 7) was isolated. The isolated fragment was then subjected to an end-filling reaction using the Klenow fragment of DNA polymerase. The blunt-ended fragment was then ligated to a similarly opened *BamHI* restriction site in the K14 vector. The resulting construct contained the *EcoRI* site derived from the polylinker of the pCI-neo vector. This *EcoRI*
25 site was removed using standard techniques (a Klenow-mediated fill-in reaction following partial digestion of the recombinant intermediate with *EcoRI*) to facilitate the subsequent excision of the DNA fragment to be injected into fertilized mouse oocytes. The resulting clone, designated K14-VEGF-C, is illustrated in Fig. 20 of commonly-owned PCT patent application PCT/FI96/00427, filed August 01, 1996, published as WO 97/05250.

30 The *EcoRI-HindIII* fragment from clone K14 VEGF-C containing the K14 promoter, VEGF-C cDNA, and K14 polyadenylation signal was isolated and injected into fertilized oocytes of the FVB-NIH mouse strain. The injected zygotes were transplanted

to oviducts of pseudopregnant C57BL/6 x DBA/2J hybrid mice. The resulting founder mice were analyzed for the presence of the transgene by polymerase chain reaction of tail DNA using the primers: 5'-CATGTACGAACCGCCAG-3' (SEQ ID NO: 22) and 5'-AATGACCAGAGAGAGGCGAG-3' (SEQ ID NO: 23). In addition, the tail DNAs were
5 subjected to *EcoRV* digestion and subsequent Southern analysis using the *EcoRI-HindIII* fragment injected into the mice. Out of 8 pups analyzed at 3 weeks of age, 2 were positive, having approximately 40-50 copies and 4-6 copies of the transgene in their respective genomes.

The mouse with the high copy number transgene was small, developed
10 more slowly than its litter mates and had difficulty eating (*i.e.*, suckling). Further examination showed a swollen, red snout and poor fur. Although fed with a special liquid diet, it suffered from edema of the upper respiratory and digestive tracts after feeding and had breathing difficulties. This mouse died eight weeks after birth and was immediately processed for histology, immunohistochemistry, and *in situ* hybridization.

15 Histological examination showed that in comparison to the skin of littermates, the dorsal dermis of K14-VEGF-C transgenic mice was atrophic and connective tissue was replaced by large lacunae devoid of red cells, but lined with a thin endothelial layer. These distended vessel-like structures resembled those seen in human lymphangiomas. The number of skin adnexal organs and hair follicles were reduced. In
20 the snout region, an increased number of vessels was also seen. Therefore, VEGF-C overexpression in the basal epidermis is capable of promoting the growth of extensive vessel structure in the underlying skin, including large vessel lacunae. The endothelial cells surrounding these lacunae contained abundant Flt4 mRNA in *in situ* hybridization (see Examples 23 and 30 for methodology). The vessel morphology indicates that VEGF-C
25 stimulates the growth of vessels having features of lymphatic vessels. The other K14-VEGF-C transgenic mouse had a similar skin histopathology.

Nineteen additional pups were analyzed at 3 weeks of age for the presence of the VEGF-C transgene, bring the number of analyzed pups to twenty-seven. A third transgene-positive pup was identified, having approximately 20 copies of the transgene in
30 its genome. The 20 copy mouse and the 4-6 copy mouse described above transmitted the gene to 6 out of 11 and 2 out of 40 pups, respectively. The physiology of these additional transgenic mice were further analyzed.

The adult transgenic mice were small and had slightly swollen eyelids and poorly developed fur. Histological examination showed that the epidermis was hyperplastic and the number of hair follicles was reduced; these effects were considered unspecific or secondary to other phenotypic changes. The dermis was atrophic (45% of the dermal thickness, compared to 65% in littermate controls) and its connective tissue was replaced by large dilated vessels devoid of red cells, but lined with a thin endothelial cell layer. Such abnormal vessels were confined to the dermis and resembled the dysfunctional, dilated spaces characteristic of hyperplastic lymphatic vessels. See Fossum, *et al.*, *J. Vet. Int. Med.*, 6: 283-293 (1992). Also, the ultrastructural features were reminiscent of lymphatic vessels, which differ from blood vessels by having overlapping endothelial junctions, anchoring filaments in the vessel wall, and a discontinuous or even partially absent basement membrane. See Leak, *Microvasc. Res.*, 2: 361-391 (1970). Furthermore, antibodies against collagen types IV, XVIII [Muragaki *et al.*, *Proc. Natl. Acad. Sci. USA*, 92: 8763-8776 (1995)] and laminin gave very weak or no staining of the vessels, while the basement membrane staining of other vessels was prominent. The endothelium was also characterized by positive staining with monoclonal antibodies against desmoplakins I and II (Progen), expressed in lymphatic, but not in vascular endothelial cells. See Schmelz *et al.*, *Differentiation*, 57: 97-117 (1994). Collectively, these findings strongly suggested that the abnormal vessels were of lymphatic origin.

In Northern hybridization studies, abundant VEGF-C mRNA was detected in the epidermis and hair follicles of the transgenic mice, while mRNAs encoding its receptors VEGFR-3 and VEGFR-2 as well as the Tie-1 endothelial receptor tyrosine kinase [Korhonen *et al.*, *Oncogene*, 9: 395-403 (1994)] were expressed in endothelial cells lining the abnormal vessels. In the skin of littermate control animals, VEGFR-3 could be detected only in the superficial subpapillary layer of lymphatic vessels, while VEGFR-2 was found in all endothelia, in agreement with earlier findings. See Millauer *et al.*, *Cell*, 72: 1-20 (1993); and Kaipainen *et al.*, *Proc. Natl. Acad. Sci. USA*, 92: 3566-3570 (1995).

The lymphatic endothelium has a great capacity to distend in order to adapt to its functional demands. To determine whether vessel dilation was due to endothelial distension or proliferation, *in vitro* proliferation assays were conducted. Specifically, to measure DNA synthesis, 3mm x 3mm skin biopsies from four transgenic and four control mice were incubated in D-MEM with 10 micrograms/ml BrdU for 6 hours at 37°C, fixed

- 74 -

in 70% ethanol for 12 hours, and embedded in paraffin. After a 30 minute treatment with 0.1% pepsin in 0.1 M HCl at room temperature to denature DNA, staining was performed using mouse monoclonal anti-BrdU antibodies (Amersham). It appeared that the VEGF-C-receptor interaction in the transgenic mice transduced a mitogenic signal, because, in contrast to littermate controls, the lymphatic endothelium of the skin from young K14-VEGF-C mice showed increased DNA synthesis as demonstrated by BrdU incorporation followed by staining with anti-BrdU antibodies. This data further confirms that VEGF-C acts as a true growth factor in mammalian tissues.

In related experiments, a similar VEGF transgene did not induce lymphatic proliferation, but caused enhanced density of hyperpermeable, tortuous blood microvessels instead.

Angiogenesis is a multistep process which includes endothelial proliferation, sprouting, and migration. See Folkman *et al.*, *J. Biol. Chem.*, 267: 10931-10934 (1992). To estimate the contribution of such processes to the transgenic phenotype, the morphology and function of the lymphatic vessels was analysed using fluorescent microlymphography using techniques known in the art. See Leu *et al.*, *Am. J. Physiol.*, 267: 1507-1513 (1994); and Swartz *et al.*, *Am. J. Physiol.*, 270: 324-329 (1996). Briefly, eight-week old mice were anesthetized and placed on a heating pad to maintain a 37°C temperature. A 30-gauge needle, connected to a catheter filled with a solution of FITC-dextran 2M (8 mg/ml in PBS), was injected intradermally into the tip of the tail. The solution was infused with a constant pressure of 50 cm water (averaging roughly 0.01 microliters per minute flow rate) until the extent of network filling remained constant (approximately 2 hours). Flow rate and fluorescence intensity were monitored continuously throughout the experiment. In these experiments, a typical honeycomb-like network with similar mesh sizes was observed in both control and transgenic mice, but the diameter of lymphatic vessels was about twice as large in the transgenic mice, as summarized in the table below. (The intravital fluorescence microscopy of blood vessels

- 75 -

was performed as has been described in the art. See Fukumura *et al.*, *Cancer Res.*, 55: 4824-4829 (1995).)

Structural parameters of lymphatic and blood vessel networks				
		transgenic	control	P-value**
		(n=4)	(n=5)	
5 lymphatic vessels*	diameter	142.3±26.2	68.2±21.7	.0143
	horizontal mesh size***	1003±87.1	960.8±93.1	.2207
	Vertical mesh size	507.3±58.9	488.8±59.9	.5403
		(n=3)	(n=6)	
blood vessels	median diameter	8.3±0.6	7.6±1.1	.1213
	vessel density, cm/cm ²	199.2±6.6	216.4±20.0	.3017

n=number of animals

* mean (μm)±SD

10 **Mann-Whitney test

***mesh size describes vessel density

Some dysfunction of the abnormal vessels was indicated by the fact that it took longer for the dextran to completely fill the abnormal vessels. Injection of FITC-dextran into the tail vein, followed by fluorescence microscopy of the ear, showed that the blood vascular morphology was unaltered and leukocyte rolling and adherence appeared normal in the transgenic mice. These results suggest that the endothelial proliferation induced by VEGF-C leads to hyperplasia of the superficial lymphatic network but does not induce the sprouting of new vessels.

These effects of VEGF-C overexpression are unexpectedly specific, especially since, as described in other examples, VEGF-C is also capable of binding to and activating VEGFR-2, which is the major mitogenic receptor of blood vessel endothelial cells. In culture, high concentrations of VEGF-C stimulate the growth and migration of bovine capillary endothelial cells which express VEGFR-2, but not significant amounts of VEGFR-3. In addition, VEGF-C induces vascular permeability in the Miles assay [Miles,

- 76 -

A. A., and Miles, E. M., *J. Physiol.*, 118:228-257 (1952); and Udaka, *et al.*, *Proc. Soc. Exp. Biol. Med.*, 133:1384-1387 (1970)], presumably via its effect on VEGFR-2. VEGF-C is less potent than VEGF in the Miles assay, 4- to 5-fold higher concentrations of VEGF-C Δ N Δ CHis being required to induce the same degree of permeability. *In vivo*, the specific effects of VEGF-C on lymphatic endothelial cells may reflect a requirement for the formation of VEGFR-3xVEGFR-2 heterodimers for endothelial cell proliferation at physiological concentrations of the growth factor. Such possible heterodimers may help to explain how three homologous VEGFs exert partially redundant, yet strikingly specific biological effects.

10 The foregoing *in vivo* data indicates utilities for both (i) VEGF-C polypeptides and polypeptide variants and analogs having VEGF-C biological activity, and (ii) anti-VEGF-C antibodies and VEGF-C antagonists that inhibit VEGF-C activity (*e.g.*, by binding VEGF-C or interfering with VEGF-C/receptor interactions. For example, the data indicates a therapeutic utility for VEGF-C polypeptides in patients wherein growth of lymphatic tissue may be desirable (*e.g.*, in patients following breast cancer or other surgery where lymphatic tissue has been removed and where lymphatic drainage has therefore been compromised, resulting in swelling; or in patients suffering from elephantiasis). The data indicates a therapeutic utility for anti-VEGF-C antibody substances and VEGF-C antagonists for conditions wherein growth-inhibition of lymphatic tissue may be desirable
15 (e.g., treatment of lymphangiomas). Accordingly, methods of administering VEGF-C and VEGF-C variants, analogs, and antagonists are contemplated as methods and materials of the invention.

Example 30

Expression of VEGF-C and Flt4 in the Developing Mouse

25 Embryos from a 16-day post-coitus pregnant mouse were prepared and fixed in 4% paraformaldehyde (PFA), embedded in paraffin, and sectioned at 6 μ m. The sections were placed on silanated microscope slides and treated with xylene, rehydrated, fixed for 20 minutes in 4% PFA, treated with proteinase K (7mg/ml; Merck, Darmstadt, Germany) for 5 minutes at room temperature, again fixed in 4% PFA and treated with
30 acetic anhydride, dehydrated in solutions with increasing ethanol concentrations, dried and used for *in situ* hybridization.

- 77 -

In situ hybridization of sections was performed as described (Västrik *et al.*, *J. Cell Biol.*, 128:1197-1208 (1995)). A mouse VEGF-C antisense RNA probe was generated from linearized pBluescript II SK+ plasmid (Stratagene Inc.), containing a fragment corresponding to nucleotides 499-979 of mouse VEGF-C cDNA, where the 5 noncoding region and the BR3P repeat were removed by Exonuclease III treatment. The fragment had been cloned into the *EcoRI* and *HindIII* sites of pBluescript II SK+. Radiolabeled RNA was synthesized using T7 RNA Polymerase and [³⁵S]-UTP (Amersham, Little Chalfont, UK). About two million cpm of the VEGF-C probe was applied per slide. After an overnight hybridization, the slides were washed first in 2x SSC and 20-30 mM DDT for 1 hour at 50°C. Treatment continued with a high stringency wash, 4x SSC and 20 mM DDT and 50% deionized formamide for 30 minutes at 65°C followed by RNase A treatment (20 µg/ml) for 30 minutes at 37°C. The high stringency wash was repeated for 45 minutes. Finally, the slides were dehydrated and dried for 30 minutes at room temperature. The slides were dipped into photography emulsion and exposed for 4 weeks. Slides were developed using Kodak D-16 developer, counterstained with hematoxylin and mounted with Permout (FisherChemical).

For *in situ* hybridizations of Flt4 sequences, a mouse Flt4 cDNA fragment covering bp 1-192 of the published sequence (Finnerty *et al.*, *Oncogene*, 8:2293-2298 (1993)) was used, and the above-described protocol was followed, with the following exceptions. Approximately one million cpm of the Flt4 probe were applied to each slide. The stringent washes following hybridization were performed in 1x SSC and 30 mM DTT for 105 minutes.

Darkfield and lightfield photomicrographs from these experiments are presented in commonly-owned PCT patent application PCT/FI96/00427, filed August 01, 1996, incorporated by reference herein. Observations from the photomicrographs are summarized below.

The most prominently Flt4-hybridizing structures appeared to correspond to the developing lymphatic and venous endothelium. A plexus-like endothelial vascular structure surrounding the developing nasopharyngeal mucous membrane was observed. The most prominent signal using the VEGF-C probe was obtained from the posterior part of the developing nasal conchae, which in higher magnification showed the epithelium surrounding loose connective tissue/forming cartilage. This structure gave a strong *in situ*

- 78 -

hybridization signal for VEGF-C. With the VEGF-C probe, more weakly hybridizing areas were observed around the snout, although this signal is much more homogeneous in appearance. Thus, the expression of VEGF-C is strikingly high in the developing nasal conchae.

5 The conchae are surrounded with a rich vascular plexus, important in nasal physiology as a source for the mucus produced by the epithelial cells and for warming inhaled air. It is suggested that VEGF-C is important in the formation of the conchal venous plexus at the mucous membranes, and that it may also regulate the permeability of the vessels needed for the secretion of nasal mucus. Possibly, VEGF-C and its derivatives,
10 and antagonists, could be used in the regulation of the turgor of the conchal tissue and mucous membranes and therefore the diameter of the upper respiratory tract, as well as the quantity and quality of mucus produced. These factors are of great clinical significance in inflammatory (including allergic) and infectious diseases of the upper respiratory tract. Accordingly, the invention contemplates the use of the materials of the invention, including
15 VEGF-C, Flt4, and their derivatives, in methods of diagnosing and treating inflammatory and infectious conditions affecting the upper respiratory tract, including nasal structures.

Example 31

Characterization of the exon-intron organization of the human VEGF-C gene

20 Two genomic DNA clones covering exons 1, 2, and 3 of the human VEGF-C gene were isolated from a human genomic DNA library using VEGF-C cDNA fragments as probes. In particular, a human genomic library in bacteriophage EMBL-3 lambda (Clontech) was screened using a PCR-generated fragment corresponding to nucleotides 629-746 of the human VEGF-C cDNA (SEQ ID NO: 7). One positive clone,
25 designated "lambda 3," was identified, and the insert was subcloned as a 14 kb *Xho*I fragment into the pBluescript II (pBSK II) vector (Stratagene). The genomic library also was screened with a labeled 130 bp *Not*I-*Sac*I fragment from the 5'-noncoding region of the VEGF-C cDNA (the *Not*I site is in the polylinker of the cloning vector; the *Sac*I site corresponds to nucleotides 92-97 of SEQ ID NO: 7). Two positive clones, designated
30 "lambda 5" and "lambda 8," were obtained. Restriction mapping analysis showed that

- 79 -

clone lambda 3 contains exons 2 and 3, while clone lambda 5 contains exon 1 and the putative promoter region.

Three genomic fragments containing exons 4, 5, 6 and 7 were subcloned from a genomic VEGF-C P1 plasmid clone. In particular, purified DNA from a genomic P1 plasmid clone 7660 (Paavonen *et al.*, *Circulation*, 93: 1079-1082 (1996)) was used. *EcoRI* fragments of the P1 insert DNA were ligated into pBSK II vector. Subclones of clone 7660 which contained human VEGF-C cDNA homologous sequences were identified by colony hybridization, using the full-length VEGF-C cDNA as a probe. Three different genomic fragments were identified and isolated, which contained the remaining 10 exons 4-7.

To determine the genomic organization, the clones were mapped using restriction endonuclease cleavage. Also, the coding regions and exon-intron junctions were partially sequenced. The result of this analysis is depicted in Figures 11A and 12. The sequences of all intron-exon boundaries (Fig. 11A, SEQ ID NOs: 24-35) conformed to the consensus splicing signals (Mount, *Nucl. Acids Res.*, 10: 459-472 (1982)). The length of the intron between exon 5 and 6 was determined directly by nucleotide sequencing and found to be 301 bp. The length of the intron between exons 2 and 3 was determined by restriction mapping and Southern hybridization and was found to be about 1.6 kb. Each of the other introns is over 10 kb in length.

A similar analysis was performed for the murine genomic VEGF-C gene. The sequences of murine VEGF-C intron-exon boundaries are depicted in Figure 11B and SEQ ID NOs: 36-47.

The restriction mapping and sequencing data indicated that the VEGF-C signal sequence and the first residues of the N-terminal propeptide are encoded by exon 1. The second exon encodes the carboxy-terminal portion of the N-terminal propeptide and the amino terminus of the VEGF homology domain. The most conserved sequences of the VEGF homology domain are distributed in exons 3 (containing 6 conserved cysteine residues) and 4 (containing 2 cys residues). The remaining exons encode cysteine-rich motifs of the type C-6X-C-10X-CRC (exons 5 and 7) and a fivefold repeated motif of type C-6X-B-3X-C-C-C, which is typical of a silk protein.

To further characterize the human VEGF-C gene promoter, the lambda 5 clone was further analyzed. Restriction mapping of this clone using a combination of

- 80 -

single- and double-digestions and Southern hybridizations indicated that it includes: (1) an approximately 6 kb region upstream of the putative initiator ATG codon, (2) exon 1, and (3) at least 5 kb of intron I of the VEGF-C gene.

A 3.7 kb *Xba* I fragment of clone lambda 5, containing exon 1 and 5' and 3' flanking sequences, was subcloned and further analyzed. As reported previously, a major VEGF-C mRNA band migrates at a position of about 2.4 kb. Calculating from the VEGF-C coding sequence of 1257 bp and a 391 bp 3' noncoding sequence plus a polyA sequence of about 50-200 bp, the mRNA start site was estimated to be about 550-700 bp upstream of the translation initiation codon.

10 RNase protection assays were employed to obtain a more precise localization of the mRNA start site. The results of these experiments indicated that the RNA start site in the human VEGF-C gene is located 539 bp upstream of the ATG translational initiation codon.

To further characterize the promoter of the human VEGF-C gene, a
15 genomic clone encompassing about 2.4 kb upstream of the translation initiation site was isolated, and the 5' noncoding cDNA sequence and putative promoter region were sequenced. The sequence obtained is set forth in SEQ ID NO: 48. (The beginning of the VEGF-C cDNA sequence set forth in SEQ ID NO: 7 corresponds to position 2632 of SEQ ID NO: 48; the translation initiation codon corresponds to positions 2983-2985 of
20 SEQ ID NO: 48.) Similar to what has been observed with the VEGF gene, the VEGF-C promoter is rich in G and C residues and lacks consensus TATA and CCAAT sequences. Instead, it has numerous putative binding sites (5'-GGGCGG-3' or 5'-CCGCCC-3') for Sp1, a ubiquitous nuclear protein that can initiate transcription of TATA-less genes. See Pugh and Tjian, *Genes and Dev.*, 5:105-119 (1991). In addition, sequences upstream of
25 the VEGF-C translation start site were found to contain frequent consensus binding sites for the AP-2 factor (5'-GCCN₃GCC-3') and binding sites for the AP-1 factor (5'-TKASTCA-3'). Binding sites for regulators of tissue-specific gene expression, like NFkB and GATA, are located in the distant part of VEGF-C promoter. This suggests that the cAMP-dependent protein kinase and protein kinase C, as activators of AP-2 transcription
30 factor [Curran and Franza, *Cell*, 55:395-397 (1988)], mediate VEGF-C transcriptional regulation.

- 81 -

The VEGF-C gene is abundantly expressed in adult human tissues, such as heart, placenta, ovary and small intestine, and is induced by a variety of factors. Indeed, several potential binding sites for regulators of tissue-specific gene expression, like NFkB (5'-GGGRNTYYC-3') and GATA, are located in the distal part of the VEGF-C promoter.

5 For example, NFkB is known to regulate the expression of tissue factor in endothelial cells. Also, transcription factors of the GATA family are thought to regulate cell-type specific gene expression.

Unlike VEGF, the VEGF-C gene does not contain a binding site for the hypoxia-inducible factor, HIF-1 (Levy *et al.*, *J. Biol. Chem.*, 270: 13333-13340 (1995)).

10 This finding suggests that if the VEGF-C mRNA is regulated by hypoxia, the mechanism would be based mainly on the regulation of mRNA stability. In this regard, numerous studies have shown that the major control point for the hypoxic induction of the VEGF gene is the regulation of the steady-state level of mRNA. See Levy *et al.*, *J. Biol. Chem.*, 271: 2746-2753 (1996). The relative rate of VEGF mRNA stability and decay is

15 considered to be determined by the presence of specific sequence motifs in its 3' untranslated region (UTR), which have been demonstrated to regulate mRNA stability. (Chen and Shyu, *Mol. Cell Biol.*, 14: 8471-8482 (1994)). The 3'-UTR of the VEGF-C gene also contains a putative motif of this type (TTATTT), at positions 1873-1878 of SEQ ID NO: 7.

20 To identify DNA elements important for basal expression of VEGF-C in transfected cells, a set of luciferase reporter plasmids containing serial 5' deletions through the promoter region was constructed. Restriction fragments of genomic DNA containing 5' portions of the first exon were cloned into the polylinker of the pGL3 reporter vector (Promega) and confirmed by sequencing. About 10 μ g of the individual constructs in

25 combination with 2 μ g of pSV2- β -galactosidase plasmid (used as a control of transfection efficiency) were transfected into HeLa cells using the calcium phosphate-mediated transfection method. Two days after transfection, the cells were harvested and subjected to the luciferase assay. The luciferase activity was normalized to that of the pGL3 control vector driven by SV40 promoter/enhancer.

30 As depicted in Fig. 3, the 5.5 kb *XhoI-RsrII* fragment of clone lambda 5 gave nearly 9-fold elevated activity when compared with a promoterless vector. Deletion of a 5' *XhoI-HindIII* fragment of 2 kb had no effect on the promoter activity. The activity

- 82 -

of the 1.16 kb *XbaI-RsrII* fragment was about twice that of the pGL3 basic vector, while the activity of the same fragment in the reverse orientation was at background level. Further deletion of the *XbaI-SacI* fragment caused an increase in the promoter activity, suggesting the presence of silencer elements in the region from -1057 to -199 (i.e., 199 to 5 1057 bp upstream from the transcription initiation site). The shortest fragment (*SacII-RsrII*) yielded only background activity, which was consistent with the fact that the mRNA initiation site was not present in this construct.

To determine whether further sequences in the first exon of human VEGF-C are important for basal expression, an *RsrII* fragment spanning nucleotides 214-495 (i.e., 10 214-495 bp downstream from the transcription initiation site) was subcloned in between of *XbaI-RsrII* fragment and the luciferase reporter gene. Indeed, the obtained construct showed an 50 % increase in activity when compared with the *XbaI-RsrII* construct.

The VEGF gene has been shown to be up-regulated by a number of stimuli including serum derived growth factors. To find out whether the VEGF-C gene also can 15 be stimulated by serum, RNA from serum-starved and serum-stimulated HT1080 cells was subjected to primer extension analysis, which demonstrated that VEGF-C mRNA is up-regulated by serum stimulation.

Additional serum stimulation experiments indicated that the serum stimulation leads to increased VEGF-C promoter activity. Cells were transfected as 20 described above and 24 h after transfection changed into medium containing 0.5% bovine serum albumin. Cells were then stimulated with 10 % fetal calf serum for 4 hours and analyzed. The *XbaI-RsrII* promoter construct derived from lambda 5 yielded a twofold increased activity upon serum stimulation, while the same fragment in the reverse orientation showed no response. All other promoter constructs also showed up- 25 regulation, ranging from 1.4 to 1.6 fold (Fig. 3).

Example 32

Identification of a VEGF-C splice variant

As reported in Example 16, a major 2.4 kb VEGF-C mRNA and smaller amounts of a 2.0 kb mRNA are observable. To clarify the origin of these RNAs, several 30 additional VEGF-C cDNAs were isolated and characterized. A human fibrosarcoma cDNA library from HT1080 cells in the lambda gt11 vector (Clontech, product

#HL1048b) was screened using a 153 bp human VEGF-C cDNA fragment as a probe as described in Example 10. See also Joukov *et al.*, *EMBO J.*, 15:290-298 (1996). Nine positive clones were picked and analyzed by PCR amplification using oligonucleotides 5'-CACGGCTTATGCAAGCAAAG-3' (SEQ ID NO: 49) and

5 5'-AACACAGTTTTCCATAATAG-3' (SEQ ID NO: 50) These oligonucleotides were selected to amplify the portion of the VEGF-C cDNA corresponding to nucleotides 495-1661 of SEQ ID NO: 7. PCR was performed using an annealing temperature of 55°C and 25 cycles.

The resultant PCR products were electrophoresed on agarose gels. Five clones out of the nine analyzed generated PCR fragments of the expected length of 1147 base pairs, whereas one was slightly shorter. The shorter fragment and one of the fragments of expected length were cloned into the pCRTMII vector (Invitrogen) and analyzed by sequencing. The sequence revealed that the shorter PCR fragment had a deletion of 153 base pairs, corresponding to nucleotides 904 to 1055 of SEQ ID NO: 7. These deleted bases correspond to exon 4 of the human and mouse VEGF-C genes, schematically depicted in Figs. 13A and 13B. Deletion of exon 4 results in a frameshift, which in turn results in a C-terminal truncation of the full-length VEGF-C precursor, with fifteen amino acid residues translated from exon 5 in a different frame than the frame used to express the full-length protein. Thus, the C-terminal amino acid sequence of the resulting truncated polypeptide would be --Leu (181)-Ser-Lys-Thr-Val-Ser-Gly-Ser-Glu-Gln-Asp-Leu-Pro-His-Glu-Leu-His-Val-Glu(199) (SEQ ID NO: 51). The polypeptide encoded by this splice variant would not contain the C-terminal cleavage site of the VEGF-C precursor. Thus, a putative alternatively spliced RNA form lacking conserved exon 4 was identified in HT-1080 fibrosarcoma cells and this form is predicted to encode a protein of 199 amino acid residues, which could be an antagonist of VEGF-C.

Example 33

VEGF-C is similarly processed in different cell cultures in vitro

To study whether VEGF-C is similarly processed in different cell types, 293 EBNA cells, COS-1 cells and HT-1080 cells were transfected with wild type human VEGF-C cDNA and labelled with Pro-Mix™ as described in Example 22. The conditioned media from the cultures were collected and subjected to immunoprecipitation using

- 84 -

antiserum 882 (described in Example 21, recognizing a peptide corresponding to amino acids 104-120 of SEQ ID NO: 8). The immunoprecipitated polypeptides were separated via SDS-PAGE, and detected via autoradiography. The major form of secreted recombinant VEGF-C observed from all cell lines tested is a 29/32 kD doublet. These two polypeptides are bound to each other by disulfide bonds, as described in Example 22. A less prominent band of approximately 21 kD also was detected in the culture media. Additionally, a non-processed VEGF-C precursor of 63 kDa was observed. This form was more prominent in the COS-1 cells, suggesting that proteolytic processing of VEGF-C in COS cells is less efficient than in 293 EBNA cells. Endogenous VEGF-C (in non-transfected cells) was not detectable under these experimental conditions in the HT-1080 cells, but was readily detected in the conditioned medium of the PC-3 cells. Analysis of the subunit polypeptide sizes and ratios in PC-3 cells and 293 EBNA cells revealed strikingly similar results: the most prominent form was a doublet of 29/32 kDa and a less prominent form the 21 kD polypeptide. The 21 kD form produced by 293 EBNA cells was not recognized by the 882 antibody in the Western blot, although it is recognized when the same antibody is used for immunoprecipitation (see data in previous examples). As reported in Example 21, cleavage of the 32 kD form in 293 EBNA cells occurs between amino acid residues 111 and 112 (SEQ ID NO: 8), downstream of the cleavage site in PC-3 cells (between residues 102 and 103). Therefore, the 21 kD form produced in 293 EBNA cells does not contain the complete N-terminal peptide used to generate antiserum 882. In a related experiment, PC-3 cells were cultured in serum-free medium for varying periods of time (1 - 8 days) prior to isolation of the conditioned medium. The conditioned medium was concentrated using a Centricon device (Amicon, Beverly, USA) and subjected to Western blotting analysis using antiserum 882. After one day of culturing, a prominent 32 kD band was detected. Increasing amounts of a 21-23 kD form were detected in the conditioned media from 4 day and 8 day cultures. The diffuse nature of this polypeptide band, which is simply called the 23 kD polypeptide in example 5 and several subsequent examples, is most likely due to a heterogenous and variable amount of glycosylation. These results indicate that, initially, the cells secrete a 32 kD polypeptide, which is further processed or cleaved in the medium to yield the 21-23 kD form. The microheterogeneity of this polypeptide band would then arise from the variable glycosylation degree and, from microheterogeneity of the processing cleavage sites, such

- 85 -

as obtained for the amino terminus in PC-3 and 293 EBNA cell cultures. The carboxyl terminal cleavage site could also vary, examples of possible cleavage sites would be between residues 225-226, 226-227 and 227-228 as well as between residues 216-217. Taken together, these data suggest the possibility that secreted cellular protease(s) are responsible for the generation of the 21-23 kD form of VEGF-C from the 32 kD polypeptide. Such proteases could be used in vitro to cleave VEGF-C precursor proteins in solution during the production of VEGF-C, or used in cell culture and in vivo to release biologically active VEGF-C.

Example 34

10

Differential binding of VEGF-C forms by the extracellular domains of VEGFR-3 and VEGFR-2

In two parallel experiments, 293 EBNA cells were transfected with a construct encoding recombinant wild type VEGF-C or a construct encoding VEGF-C Δ N Δ CHis (Example 28) and about 48 hours after transfection, metabolically labelled with Pro-MixTM as described in previous examples. The media were collected from mock-transfected and transfected cells and used for receptor binding analyses.

Receptor binding was carried out in binding buffer (PBS, 0.5% BSA, 0.02% Tween 20, 1 microgram/ml heparin) containing approximately 0.2 microgram of either (a) a fusion protein comprising a VEGFR-3 extracellular domain fused to an immunoglobulin sequence (VEGFR-3-Ig) or (b) a fusion protein comprising VEGFR-2 extracellular domain fused to an alkaline phosphatase sequence (VEGFR-2-AP; Cao *et al.*, *J. Biol. Chem.* 271:3154-62 (1996)). As a control, similar aliquots of the 293 EBNA conditioned media were mixed with 2 μ l of anti-VEGF-C antiserum (VEGF-C IP).

After incubation for 2 hours at room temperature, anti-VEGF-C antibodies and VEGFR-3-Ig protein were adsorbed to protein A-sepharose (PAS) and VEGFR-2-AP was immunoprecipitated using anti-AP monoclonal antibodies (Medix Biotech, Genzyme Diagnostics, San Carlos, CA, USA) and protein G-sepharose. Complexes containing VEGF-C bound to VEGFR-3-Ig or VEGFR-2-AP were washed three times in binding buffer, twice in 20 mM Tris-HCl (pH 7.4) and VEGF-C immunoprecipitates were washed three times in RIPA buffer and twice in 20 mM tris-HCl (pH 7.4) and analyzed via SDS-PAGE under reducing and nonreducing conditions. As a control, the same media were

- 86 -

precipitated with antiAP and protein G-sepharose (PGS) or with PAS to control for possible nonspecific adsorption.

These experiments revealed that VEGFR-3 bound to both the 32/29 kD and 21-23 kD forms of recombinant VEGF-C, whereas VEGFR-2 bound preferentially to the 21-23 kD component from the conditioned media. In addition, small amounts of 63 kD and 52 kD VEGF-C forms were observed binding with VEGFR-3. Further analysis under nonreducing conditions indicates that a great proportion of the 21-23 kD VEGF-C bound to either receptor does not contain interchain disulfide bonds. These findings reinforce the results that VEGF-C binds VEGFR-2. This data suggests a utility for recombinant forms of VEGF-C which are active towards VEGFR-3 only or which are active towards both VEGFR-3 and VEGFR-2. On the other hand, these results, together with the results in Example 28, do not eliminate the possibility that the 32/29 kD dimer binds VEGFR-3 but does not activate it. The failure of the 32/29 kD dimer to activate VEGFR-3 could explain the finding that conditioned medium from the N-His VEGF-C transfected cells induced a less prominent tyrosine phosphorylation of VEGFR-3 than medium from VEGF-C Δ N Δ CHis transfected cells, even though expression of the former polypeptide was much higher. Stable VEGF-C polypeptide mutants that bind to a VEGF-C receptor but fail to activate the receptor are useful as VEGF-C antagonists.

Example 35

20 **Discovery of VEGF-C analogs that selectively bind to and activate VEGFR-3, but not VEGFR-2**

To further identify the cysteine residues of VEGF-C that are critical for retaining VEGF-C biological activities, an additional VEGF-C mutant, designated VEGF-C Δ N Δ CHisC156S, was synthesized, in which the cysteine residue at position 156 of the 25 419 amino acid VEGF-C precursor (SEQ ID NO: 8; Genbank accession number X94216) was replaced with a serine residue.

The mutagenesis procedure was carried out using the construct of VEGF-C Δ N Δ CHis (see Example 28), cloned in the pALTER vector, and the *Altered sites II in vitro mutagenesis system* of Promega. An oligonucleotide 5'-
30 GACGGACACAGATGGAGGTTTAAAG-3' (SEQ ID NO: 52) was used to introduce the desired mutation in the cDNA encoding VEGF-C Δ N Δ CHis. The resulting mutated

- 87 -

VEGF-C cDNA fragment was subcloned into the *HindIII/NotI* sites of the pREP-7 vector (Invitrogen), and the final construct was re-sequenced to confirm the C156S mutation. The resultant clone has an open reading frame encoding amino acids 103-225 of SEQ ID NO: 8 (with a serine codon at position 156), and further encoding a 6xHis tag.

5 The wildtype VEGF-C cDNA and three VEGF-C mutant constructs (VEGF-C R226,227S, VEGF-C Δ N Δ CHis, and VEGF-C Δ N Δ CHisC156S) were used to transfect 293 EBNA cells, which were subcultured 16 hours after transfection. About 48 hours after transfection, the media were changed to DMEM/0.1% BSA, and incubation in this medium was continued for an additional 48 hours. The resultant conditioned media
10 were concentrated 30-fold using *Centriprep-10* (Amicon), and the amount of VEGF-C in the media was analyzed by Western blotting using the anti-VEGF-C antiserum 882 for immunodetection. Different amounts of the recombinant VEGF-C Δ N Δ CHis, purified from a yeast expression system, were analyzed in parallel as reference samples to measure and equalize the VEGF-C concentrations in the conditioned media. The conditioned
15 medium from mock-transfected cells was used to dilute the VEGF-C conditioned media to achieve equal concentrations.

 An aliquot of the transfected cells were metabolically labelled for 6 hours with 100 microcuries/ml of the PRO-MIXTM L-[³⁵S] *in vitro* cell labelling mix (Amersham). The conditioned media were collected, and binding of the radioactively
20 labelled VEGF-C proteins to the extracellular domains of VEGFR-3 and VEGFR-2 was analyzed using recombinantly produced VEGFR-3EC-Ig and VEGFR-2EC-Ig constructs (containing seven and three Ig loops of the extracellular domains of the respective receptors, fused to an immunoglobulin heavy chain constant region).

 All processed VEGF-C forms secreted to the culture medium bound to
25 VEGFR-3EC domain, with preferential binding of the 21 kDa form. When present at high concentrations, the VEGF-C forms of 58 kDa and 29/31 kDa bound to some extent non-specifically to protein A Sepharose.

 The VEGFR-2EC domain preferentially bound the mature 21 kDa form of wildtype VEGF-C and VEGF-C Δ N Δ CHis. Significantly, VEGF-C Δ N Δ CHisC156S failed
30 to bind the VEGFR2-EC.

 Next, the ability of the above-described VEGF-C polypeptides to compete with the ¹²⁵I-VEGF-C Δ N Δ CHis for binding to VEGFR-2 and VEGFR-3 was analyzed.

Scatchard analysis using VEGF-C Δ CANHis provided indications of the VEGF-C binding affinity for VEGFR-3 ($K_D=135$ pM) and VEGFR-2 ($K_D=410$ pM). Ten micrograms of the purified yeast VEGF-C Δ NACHis was labeled using 3 mCi of Iodine-125, carrier-free (Amersham), and an Iodo-Gen Iodination Reagent (Pierce), according to the standard protocol of Pierce. The resulting specific activity of the labeled VEGF-C Δ NACHis was 1.25×10^5 cpm/ng.

To study receptor binding, PAE/VEGFR-2 and PAE/VEGFR-3 cells were seeded into 24-well tissue culture plates (Nunc), which had been coated with 2% gelatin in PBS. The 125 I-VEGF-C Δ NACHis (2×10^5 cpm) and different amounts of media containing equal concentrations of the non-labeled VEGF-C (wildtype and mutants) were added to each plate in Ham's F12 medium, containing 25 mM HEPES (pH 8.0), 0.1% BSA, and 0.1% NaN_3 . The binding was allowed to proceed at room temperature for 90 minutes. The plates were then transferred onto ice and washed three times with ice-cold PBS containing 0.1% BSA. The cells were then lysed in 1 M NaOH, the lysates were collected, and the radioactivity was measured using a γ -counter. Binding in the presence of VEGF-C-containing conditioned medium was calculated as a percentage of binding observed in parallel control studies wherein equal volumes of medium from mock-transfected cells were used instead of VEGF-C conditioned media.

As shown in Fig. 4, left panel, all VEGF-C mutants displaced 125 I-VEGF-C Δ NACHis from VEGFR-3. The efficiency of displacement was as follows: VEGF-C Δ NACHisC156S > VEGF-C Δ NACHis > wildtype VEGF-C > VEGF-CR226,227S. These results indicate that enhanced binding to VEGFR-3 was obtained upon "recombinant maturation" of VEGF-C. Recombinant VEGF165 failed to displace VEGF-C from VEGFR-3.

VEGF, VEGF-C Δ NACHis, and wildtype VEGF-C all efficiently displaced labeled VEGF-C Δ NACHis from VEGFR-2, with VEGF-C Δ NACHis being more potent when compared to wildtype VEGF-C (Fig. 4, right panel). The non-processed VEGF-C R226,227S showed only weak competition of 125 I-VEGF-C Δ NACHis.

Surprisingly, VEGF-C Δ NACHisR156S failed to displace VEGF-C Δ NACHis from VEGFR-2, thus confirming the above described results obtained using a soluble extracellular domain of VEGFR-2.

- 89 -

The ability of the above mentioned VEGF-C forms to stimulate tyrosine phosphorylation of VEGFR-3 and VEGFR-2 was also investigated. Importantly, identical dilutions of the conditioned media were used for these experiments and for the competitive binding experiments described above. A Western blot analysis of the conditioned media
5 using anti-VEGF-C antiserum 882 was performed to confirm the approximately equal relative amounts of the factors present.

The stimulation of VEGFR-3 and VEGFR-2 autophosphorylation by the different VEGF-C forms in general correlated with their binding properties, as well as with the degree of "recombinant processing" of VEGF-C. The VEGF-C Δ N Δ CHisC156S
10 appeared to be at least as potent as VEGF-C Δ N Δ CHis in stimulating VEGFR-3 autophosphorylation. VEGF-C Δ N Δ CHis showed a higher potency when compared to wildtype VEGF-C in its ability to stimulate tyrosine autophosphorylation of both VEGFR-2 and VEGFR-3. The VEGF-CR226,227S conditioned medium possessed a considerably weaker effect on autophosphorylation of VEGFR-3, and almost no effect on VEGFR-2
15 autophosphorylation.

Stimulation of VEGFR-2 tyrosine phosphorylation by VEGF-C Δ N Δ CHisC156S did not differ from that of conditioned medium from the mock transfected cells, thus confirming the lack of VEGFR-2-binding and VEGFR-2-activating properties of this mutant.

20 The ability of VEGF-C Δ N Δ CHisC156S to alter vascular permeability *in vivo* was analyzed using the Miles assay (see Example 29). The recombinant VEGF-C forms assayed (Δ N Δ CHis, Δ N Δ CHisC156S) were produced by 293 cells, purified from conditioned media using Ni-NTA Superflow resin (QIAGEN) as previously described, and pretreated with 15 μ g/ml of anti-human VEGF neutralizing antibody (R&D systems) to
25 neutralize residual amounts of co-purified, endogenously produced VEGF. Eight picomoles of the various VEGF-C forms, as well as 2 pmol of recombinant human VEGF165 (R&D systems) and approximately 2 pmol of VEGF165 from the conditioned medium which were either non-treated or pretreated with the above mentioned VEGF-neutralizing antibody were injected subcutaneously to the back region of a guinea pig. The
30 area of injection was analyzed 20 minutes after injections. Both VEGF and VEGF-C Δ N Δ CHis caused increases in vascular permeability, whereas Δ N Δ CHisC156S did not affect vascular permeability. The neutralizing antibody completely blocked permeability

- 90 -

activity of VEGF but did not affect VEGF-C activity. Residual permeability activity observed for the VEGF-containing conditioned medium even after its treatment with VEGF neutralizing antibody was presumably caused by permeability factors other than VEGF that are produced by 293 cells.

5 In yet another assay, the ability of VEGF-C Δ N Δ CHis and VEGF-C Δ N Δ CHisC156S to stimulate migration of bovine capillary endothelial cells in a collagen gel was analyzed. The Δ N Δ CHis form dose-dependently stimulated migration, whereas the Δ N Δ CHisC156S form had no significant activity in the assay.

The Miles assay also was used to assay the ability of VEGF-C R226,227S
10 (8 pM, pretreated with anti-VEGF antibody) to induce vascular permeability. The results indicated that the ability of VEGF-C R226,227S to induce vascular permeability was much weaker when compared to wildtype and Δ N Δ CHis forms of VEGF-C. Collectively, this Miles assay data is consistent with the VEGFR-2 binding and autophosphorylation data described above, and indicates that VEGF-C effect on vascular permeability is mediated
15 via VEGFR-2.

Mitogenic signals from growth factor receptors are frequently relayed via the extracellular signal regulated kinases/mitogen activated protein kinases (ERK/MAPK) pathway into the nucleus. Purified recombinant VEGF-C Δ N Δ CHis and VEGF-C Δ N Δ C156S produced by a Pichia expression system were used to determine MAPK
20 pathway activation of cells expressing either VEGFR-2 or VEGFR-3. The growth factor treated cells were lysed, and activated MAPK was detected using Western blotting with antibodies against the phosphorylated forms of ERK1 and ERK2. At a concentration of 100 ng/ml, VEGF-C Δ N Δ CHis showed rapid activation of the ERK1 and ERK2 MAPK in both VEGFR-2- and VEGFR-3-expressing cells. In contrast, VEGF-C Δ N Δ C156S
25 activated ERK1 and ERK2 exclusively in the VEGFR-3-expressing cells. At the concentrations used, both VEGF-C Δ N Δ CHis and VEGF-C Δ N Δ C156S appeared to be equally potent in activating the MAPK through VEGFR-3. The amounts of total MAPK protein were confirmed to be similar in the treated and untreated cells, as shown by staining of the filter with p44/p42 MAPK antibodies made against a synthetic peptide of
30 rat p42.

The foregoing data indicates that proteolytic processing of VEGF-C results in an increase in its ability to bind and to activate VEGFR-3 and VEGFR-2. Non-

- 91 -

processed VEGF-C is a ligand and an activator of preferentially VEGFR-3, while the mature 21/23 kDa VEGF-C form is a high affinity ligand and an activator of both VEGFR-3 and VEGFR-2.

Moreover, replacement of the cysteine residue at position 156 (of prepro-
5 VEGF-C, SEQ ID NO: 8) creates a selective ligand and activator of VEGFR-3. This alteration inactivates the ability of processed VEGF-C to bind to VEGFR-2 and to activate VEGFR-2. Importantly, it is believed that the elimination of the cysteine at position 156 is the alteration responsible for this unexpected alteration in VEGF-C selectivity, and not the substitution of a serine per se. It is expected that replacement of the cysteine at position
10 156 with other amino acids, or the mere deletion of this cysteine, will also result in VEGF-C analogs having selective biological activity with respect to VEGFR-3. All such replacement and deletion analogs (collectively referred to as VEGF-C ΔC_{156} polypeptides) are contemplated as aspects of the present invention. Thus, "VEGF-C ΔC_{156} polypeptides" of the invention derived from human VEGF-C include polypeptides depicted in SEQ ID
15 NO: 58, fragments of those polypeptides (especially fragments having an amino terminus anywhere between residues 102 and 161 of SEQ ID NO: 58 and a carboxy-terminus anywhere between residues 210 and 228 of SEQ ID NO: 58). "VEGF-C ΔC_{156} polypeptides" of the invention also include the corresponding polypeptides derived from murine, quail, and other wildtype VEGF-C polypeptides.

20 VEGF-C polypeptides that have the C156S mutation (or functionally equivalent mutations at position 156) and that retain biological activity with respect to VEGFR-3, such as VEGF-C $\Delta N\Delta CHisC156S$, are useful in all of the same manners described above for wildtype VEGF-C proteins and biologically active fragments thereof where VEGFR-3 stimulation is desired. It is contemplated that most biologically active
25 VEGF-C fragments and processing variants, including but not limited to the biologically active fragments and variants identified in preceding examples, will retain VEGF-C biological activity (as mediated through VEGFR-3) when a ΔC_{156} mutation is introduced. All such biologically active VEGF-C ΔC_{156} polypeptides are intended as an aspect of the present invention.

30 Moreover, VEGF-C forms containing the C156S mutation or equivalent mutations can be used to distinguish those effects of VEGF-C mediated via VEGFR-3 and VEGFR-2 from those obtained via only VEGFR-3. The ability of such VEGF-C

- 92 -

polypeptides to selectively stimulate VEGFR-3 are also expected to be useful in clinical practice, it being understood that selectivity of a pharmaceutical is highly desirable in many clinical contexts. For example, the selectivity of VEGF-C ΔC_{156} polypeptides for VEGFR-3 binding suggests a utility for these peptides to modulate VEGF-C biological activities mediated through VEGFR-3, without significant concomitant modulation of blood vessel permeability or other VEGF-C activities that are modulated through VEGFR-2.

The data presented herein also indicates a utility for ΔC_{156} polypeptides that are capable of binding VEGFR-3, but that do not retain biological activity mediated through VEGFR-3. Specifically, such forms are believed to be capable of competing with wildtype VEGF-C for binding to VEGFR-3, and are therefore contemplated as molecules that inhibit VEGF-C-mediated stimulation of VEGFR-3. Because of the ΔC_{156} alteration, such polypeptides (especially covalent or noncovalent dimers of such polypeptides) are not expected to bind VEGFR-2. Thus, certain ΔC_{156} polypeptides and polypeptide dimers are expected to have utility as selective inhibitors of VEGF-C biological activity mediated through VEGFR-3 (i.e., without substantially altering VEGF-C mediated stimulation of VEGFR-2).

In another embodiment of the invention, heterodimers comprising a biologically active VEGF-C polypeptide in association with a ΔC_{156} polypeptide are contemplated. It is contemplated that such heterodimers can be formed *in vitro*, as described below in Example 37, or formed *in vivo* with endogenous VEGF-C following administration of a ΔC_{156} polypeptide. Such heterodimers are contemplated as modulators of VEGF-C mediated effects in cells where the biological effects of VEGF-C are mediated through VEGFR-2/VEGFR-3 heterodimers. VEGF-C ΔC_{156} polypeptides in homodimers or in heterodimers with wt VEGF-C might selectively inhibit the ability of the latter to induce VEGF-like effects, particularly to increase the vascular permeability.

Replacement of the second and/or the fourth of the eight conserved cysteine residues of VEGF abolishes VEGF dimer formation and VEGF biological activity. The analogous effect was investigated for VEGF-C, wherein the cysteines at positions 156 and 165 of SEQ ID NO: 8 correspond to the second and fourth conserved cysteines. No homodimers were obtained when VEGF-C $\Delta N\Delta C$ HisC156,165S (i.e., Cys₁₅₆ and Cys₁₆₅ both replaced with serine residues) or in VEGF-C $\Delta N\Delta C$ HisC165S were chemically crosslinked. On the other hand, about half of both crosslinked VEGF-C $\Delta N\Delta C$ His and

- 93 -

VEGF-CANACHisC156S migrated as dimers. This data indicates that VEGF-CANACHisC156S forms homodimers. Moreover, unlike VEGF-CANACHis, which forms preferentially non-covalently bound dimers, a fraction of VEGF-CANACHisC156S was disulfide bonded, as detected by SDS-PAGE in non-reducing conditions. In receptor binding studies (using procedures such as those described above), the C165S and C156,165S forms were both unable to bind VEGFR-3 or VEGFR-2. Collectively, these data suggest that homodimerization is required for VEGFR-3 activation by VEGF-C, and indicate that the inability of $\Delta\Delta$ C156S to activate VEGFR-2 and to induce VEGF-like effects is not due to an inability of this mutant to form homodimers.

10

Example 36**Utility for VEGF-C in promoting myelopoiesis**

The effects of VEGF-C on hematopoiesis were also analyzed. Specifically, leukocytes populations were analyzed in blood samples taken from the F1 transgenic mice described in Example 29, and from their non-transgenic littermates. Leukocyte population data from these mice and from non-transgenic FVB-NIH control mice (i.e., the strain used to generate the transgenic mice) are set forth in the tables below.

20

FVB/NIH MICE					
Cell Type	male 5.5 months	male 5.5 months	female 9.5 months	male 9.5 months	mean$\pm\sigma$
Lymphocytes	72.20%	82.17%	84.25%	74.25%	78.22 \pm 5.10
Neutrophils	23.00%	15.17%	14.25%	22.25%	18.67 \pm 3.98
Monocytes	0.65%	1.00%	0.25%	0.50%	0.60 \pm 0.27
Eosinophils	2.15%	1.70%	1.25%	3.00%	2.03 \pm 0.65
Basophils	0.00%	0.00%	0.00%	0.00%	0 \pm 0

VEGF-C TRANSGENIC MICE				
Cell Type	male 2 months	male 3.5 months	male 7 months	mean $\pm\sigma$
Lymphocytes	41.3%	41.50%	18.70%	33.83 \pm 10.70
Neutrophils	55.3%	53.80%	80.17%	63.09 \pm 12.09
Monocytes	2.16%	1.30%	0.67%	1.38 \pm 0.61
Eosinophils	1.17%	3.50%	.50%	1.72 \pm 1.29
Basophils	0.00%	0.00%	0.00%	0 \pm 0

VEGF-C NEGATIVE CONTROL MICE (NON-TRANSGENIC LITTERMATES OF VEGF-C TRANSGENIC MICE)					
Cell Type	male 2 months	male 2 months	male 3.5 months	male 7 months	mean $\pm\sigma$
Lymphocytes	89.00%	67.50%	91.00%	71.30%	79.7 \pm 10.41
Neutrophils	7.75%	23.00%	7.00%	23.75%	15.38 \pm 8.01
Monocytes	1.50%	0.50%	0.83%	0.75%	0.90 \pm 0.37
Eosinophils	1.50%	9.00%	0.67%	4.00%	3.79 \pm 3.25
Basophils	0.00%	0.00%	0.50%	0.50%	0.25 \pm 0.25

As the foregoing data indicates, the overexpression of VEGF-C in the skin of the transgenic mice correlates with a distinct alteration in leukocyte populations. Notably, the measured populations of neutrophils were markedly increased in the transgenic mice. One explanation for the marked increase in neutrophils is a myelopoietic activity attributable to VEGF-C. A VEGF-C influence on leukocyte trafficking in and out of tissues also may effect observed neutrophil populations. Fluorescence-activated cell sorting analysis, performed on isolated human bone marrow and umbilical cord blood CD34-positive hematopoietic cells, demonstrated that a fraction of these cells are positive for Flt4 (VEGFR-3). Thus, the VEGF-C effect on myelopoiesis may be exerted through this VEGFR-3-positive cell population and its receptors. In any case, the foregoing data indicates a use for VEGF-C polypeptides to increase granulocyte (and, in particular,

- 95 -

neutrophil) counts in human or non-human subjects, i.e., in order to assist the subject fight infectious diseases. The exploitation of the myelopoietic activity of VEGF-C polypeptides is contemplated both *in vitro* (i.e., in cell culture) and *in vivo*, as a sole myelopoietic agent and in combination with other effective agents (e.g., granulocyte colony stimulating factor 5 (G-CSF)).

Additional studies of the myelopoietic effect of VEGF-C, using VEGF-C mutants (e.g., VEGF-C ΔC_{156} polypeptides, VEGF-C $\Delta N\Delta CHis$, VEGF-C R226,227S) having altered VEGFR-2 binding affinities, will elucidate whether this effect is mediated through VEGFR-2, VEGFR-3, or both receptors, for example. The results of such 10 analysis will be useful in determining which VEGF-C mutants have utility as myelopoietic agents and which have utility as agents for inhibiting myelopoiesis.

Example 37

Generation of Heterodimers consisting of members of the VEGF family of growth factors

15 Both naturally-occurring and recombinantly-produced heterodimers of polypeptides of the PDGF/VEGF family of growth factors have been shown to exist in nature and possess mitogenic activities. See, e.g., Cao *et al.*, *J. Biol. Chem.*, 271:3154-62 (1996); and DiSalvo, *et al.*, *J. Biol. Chem.*, 270:7717-7723 (1995). Heterodimers comprising a VEGF-C polypeptide may be generated essentially as described In Cao *et al.* 20 (1996), using recombinantly produced VEGF-C polypeptides, such as the VEGF-C polypeptides described in the preceding examples. Briefly, a recombinantly produced VEGF-C polypeptide is mixed at an equimolar ratio with another recombinantly produced polypeptide of interest, such as a VEGF, VEGF-B, PlGF, PDGF α , PDGF β , or *c-fos* induced growth factor polypeptide. (See, e.g., Cao *et al.* (1990); Collins *et al.*, *Nature*, 25 316:748-750 (1985) (PDGF- β , GenBank Acc. No. X02811); Claesson-Welsh *et al.*, *Proc. Natl. Acad. Sci. USA*, 86(13):4917-4921 (1989) (PDGF- α , GenBank Acc. No. M22734); Claesson-Welsh *et al.*, *Mol. Cell. Biol.* 8:3476-3486 (1988) (PDGF- β , GenBank Acc. No. M21616); Olofsson *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 93:2576-2581 (1996) (VEGF-B, GenBank Acc. No. U48801); Maglione *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 88(20):9267- 30 9271 (1996) (PlGF, GenBank Acc. No. X54936); Heldin *et al.*, *Growth Factors*, 8:245-252 (1993); Folkman, *Nature Med.*, 1:27-31 (1995); Friesel *et al.*, *FASEB J.*, 9:919-25

(1995); Mustonen *et al.*, *J. Cell. Biol.*, 129:895-98 (1995); Orlandini, S., *Proc. Natl. Acad. Sci. USA*, 93(21):11675-11680 (1996); and others cited elsewhere herein. The mixed polypeptides are incubated in the presence of guanidine-HCl and DTT. The thiol groups are then protected with S-sulfonation, and the protein is dialyzed overnight, initially
5 against urea/glutathione-SH, glutathione-S-S-glutathione, and subsequently against 20 mM Tris-HCl.

In a preferred embodiment, a variety of differently processed VEGF-C forms and VEGF-C variants and analogs, such as the ones described in the preceding examples, are employed as the VEGF-C polypeptide used to generate such heterodimers.
10 Thereafter, the heterodimers are screened to determine their binding affinity with respect to receptors of the VEGF/PDGF family (especially VEGFR-1, VEGFR-2, and VEGFR-3), and their ability to stimulate the receptors (e.g., assaying for dimer-stimulated receptor phosphorylation in cells expressing the receptor of interest on their surface). The binding assays may be competitive binding assays such as those described herein and in the art. In
15 the initial binding assays, recombinantly produced proteins comprising the extracellular domains of receptors are employable, as described in preceding examples for VEGFR-2 and VEGFR-3. Heterodimers that bind and stimulate receptors are useful as recombinant growth factor polypeptides. Heterodimers that bind but do not stimulate receptors are useful as growth factor antagonists. Heterodimers that display agonistic or antagonistic
20 activities in the screening assays are further screened using, e.g., endothelial cell migration assays, vascular permeability assays, and *in vivo* assays. It will also be apparent from the preceding examples that dimers comprising two VEGF-C polypeptides (i.e., dimers of identical VEGF-C polypeptides as well as dimers of different VEGF-C polypeptides) are advantageously screened for agonistic and antagonistic activities using the same assays.

25 In one preferred embodiment, VEGF-C ΔC_{156} polypeptide is employed to make the dimers. It is anticipated that agonists and antagonists comprising a VEGF-C ΔC_{156} polypeptide will have increased specificity for stimulating and inhibiting VEGFR-3, without concomitant stimulation or inhibition of VEGFR-2.

In another preferred embodiment, VEGF-C polypeptides wherein the C-
30 terminal proteolytic cleavage site has been altered to reduce or eliminate C-terminal processing (e.g. VEGF-C R226,227S) is employed to make dimers for screening for inhibitory activity.

- 97 -

In yet another preferred embodiment, VEGF-C polypeptides comprising amino-terminal fragments (e.g., the VEGF-C 15 kD form described herein) of VEGF-C are employed to make dimers.

It is further contemplated that inactivation of only one polypeptide chain in a dimer could be enough to generate an inhibitory molecule, which is demonstrated e.g., by the generation of PDGF inhibitory mutant as reported in Vassbotn, *Mol. Cell. Biol.*, 13:4066-4076 (1993). Therefore, in one embodiment, inhibition is achieved by expression *in vivo* of a polynucleotide (e.g., a cDNA construct) encoding the heterodimerization partner which is unable to bind (or binds inefficiently) to the receptor, or by direct administration of that monomer in a pharmaceutical composition.

Example 38

Formation and Screening of Useful Recombinant VEGF/VEGF-C genes and polypeptides

Amino acid sequence comparison reveals that mature VEGF-C bears structural similarity to VEGF121 [Tischer *et al.*, *J. Biol. Chem.*, 266(18):11947-54 (1991)], with certain noteworthy structural differences. For example, mature VEGF-C contains an unpaired cysteine (position 137 of SEQ ID NO: 8) and is able to form non-covalently bonded polypeptide dimers. In one embodiment of the invention, a VEGF analog is created wherein the unpaired cysteine residue from mature VEGF-C is introduced at an analogous position of VEGF (e.g., introduced at Leu₃₈ of the human VEGF165 precursor (Fig. 2, Genbank Acc. No. M32977) to generate a VEGF^{+cys} mutant designated VEGF L58C). Such an alteration is introduced into the VEGF165 coding sequence using site-directed mutagenesis procedures known in the art, such as the procedures described above in preceding examples to generate various VEGF-C mutant forms. This VEGF^{+cys} mutant is recombinantly expressed and is screened (alone and as a heterodimer with other VEGF and VEGF-C forms) for VEGFR-2 and/or VEGFR-3 binding, stimulatory, and inhibitory activities, using *in vitro* and *in vivo* activity assays as described elsewhere herein. To form another VEGF analog of the invention, a VEGF^{+cys} mutant is altered to remove a conserved cysteine corresponding to cys₇₇ of the VEGF165 precursor. Elimination of this cysteine from the VEGF L58C would result in a VEGF analog resembling VEGF-CANΔCHisC156S. This VEGF analog is screened for its

- 98 -

VEGF-inhibitory activities with respect to VEGFR-2 and/or VEGFR-1 and for VEGF-C like stimulatory or inhibitory activities.

Another noteworthy structural difference between VEGF and VEGF-C is the absence in VEGF-C of several basic residues found in VEGF (e.g., residues Arg₁₀₈, 5 Lys₁₁₀ and His₁₁₂ in the VEGF₁₆₅ precursor shown in Fig. 2) that have been implicated in VEGF receptor binding. See Keyt *et al.*, *J. Biol. Chem.*, 271(10):5638-46 (1996). In another embodiment of the invention, codons for basic residues (lys, arg, his) are substituted into the VEGF-C coding sequence at one or more analogous positions by site-directed mutagenesis. For example, in a preferred embodiment, Glu₁₈₇, Thr₁₈₉, and Pro₁₉₁ 10 in VEGF-C (SEQ ID NO: 8) are replaced with Arg, Lys, and His residues, respectively. The resultant VEGF-C analogs (collectively termed "VEGF-C^{basic}" polypeptides) are recombinantly expressed and screened for VEGFR-1, VEGFR-2, and VEGFR-3 stimulatory and inhibitory activity. The foregoing VEGF and VEGF-C analogs that have VEGF-like activity, VEGF-C-like activity, or that act as inhibitors of VEGF or VEGF-C, 15 are contemplated as additional aspects of the invention. Polynucleotides encoding the analogs also are intended as aspects of the invention.

EXAMPLE 39

EFFECTS OF VEGF-C ON GROWTH AND DIFFERENTIATION OF HUMAN CD34+ PROGENITOR CELLS *IN VITRO*

20 Human CD34+ progenitor cells (HPC, 10×10^3) were isolated from bone marrow or cord blood mononuclear cells using the MACS CD34 Progenitor cell Isolation Kit (Miltenyi Biotec, Bergish Gladbach, Germany), according to the instructions of the manufacturer and cultured in RPMI 1640 medium supplemented with L-glutamine (2.5 mM), penicillin (125 IE/ml), streptomycin (125 µg/ml) and pooled 10 % umbilical cord 25 blood (CB) plasma at 37 °C in a humidified atmosphere in the presence of 5% CO₂ for seven days, with or without VEGF-C and with or without one of the combinations of growth factors described below. Each experiment was performed in triplicate. After seven days, total cell number was evaluated in each culture.

In a first set of experiments, VEGF-C was added, at concentrations ranging 30 from 10 ng/ml to 1 µg/ml, to the cultures of CB CD34+ HPCs. Cell numbers were evaluated at day 7 of culture. When added as a single factor, 100 ng/ml of VEGF-C was

- 99 -

found support the survival and proliferation of only a few CD34+ HPCs under serum-free conditions. With medium alone, most of the cells died within a culture period of 7 days. However, there were consistently more cells in the cultures provided with the VEGF-C.

A subsequent set of experiments investigated the co-stimulatory effect of 5 VEGF-C in cultures either supplemented with recombinant human stem cell factor (rhSCF, 20 ng/ml PreproTech, Rocky Hill, NY) alone or a combination of granulocyte macrophage colony stimulating factor (rhGM-CSF, 100 ng/ml, Sandoz, Basel, Switzerland) plus SCF. Addition of VEGF-C to SCF-supplemented cultures resulted in a slight co-stimulatory effect on cell growth of CD34+ cells, and this effect was already observable at a VEGF-C 10 concentration of 10 ng/ml. Addition of VEGF-C to GM-CSF- plus SCF-supplemented cultures clearly increased cell yields after 7 days of culture, with an optimum VEGF-C concentration of 100 ng/ml. Additional experiments were conducted to analyze the co-stimulatory effects of 100 ng/ml VEGF-C on total cell yields of serum-free cultures of CB CD34+ HPC cells supplemented with either GM-CSF alone, IL-3 (rhIL-3, 100 U/ml, 15 Behring AG, Marburg, Germany) alone; or a combination of GM-CSF plus IL-3. The results are shown below in the following table:

- 100 -

Total cell number ($E \times 10^3$) after a culture period of 7 days in RPMI + 10% CBPL, + specified growth factors with (+) or without (-) VEGF-C. (Cell number at day 0 = 10)			
Growth Factor(s)	experiment number	- VEGF-C	+ VEGF-C
GM-CSF	1	11	15
	2	10	17
	3	19	25
	mean \pm SE	13.3 \pm 2.8	19.0 \pm 3.1*
IL-3	1	113	130
	2	107	113
	3	200	433
	4	45	90
	mean \pm SE	116.2 \pm 31.9	191.5 \pm 80.9
GM-CSF + IL-3	1	150	160
	2	130	140
	3	140	155
	mean \pm SE	140.0 \pm 5.7	151.7 \pm 6.0*
GM-CSF + SCF	1	31	37
	2	60	227
	3	47	50
	mean \pm SE	46.0 \pm 8.3	104.7 \pm 61.3

*mean \pm SE; p=0.02

As depicted in the table, VEGF-C led to a consistent enhancement of cell growth when added as a supplement to each growth factor or combination of growth factors tested.

Effect of VEGF-C on granulomonocytic differentiation of CD34+ progenitors

Using cells from the (7 day) plasma-supplemented cultures described above, immunofluorescence triple stainings were performed to analyze the expression of the early granulomonocytic marker molecules lysozyme (LZ) and myeloperoxidase (MPO) as well as the lipopolysaccharide (LPS) receptor CD14. The table below depicts the percentages and numbers of cells expressing MPO and/or LZ:

Percentages and numbers of cells expressing the markers MPO and LZ after 7 days of culture with (+) or without (-) VEGF-C and specified growth factors						
factor	marker	exp. no.	percent of cells positive for cell marker		numbers of cells positive for cell marker ($E \times 10^{-3}$)	
			- VEGF-C	+ VEGF-C	- VEGF-C	+ VEGF-C
GM-CSF	MPO	1	57	69	6	11
		2	45	53	5	9
		3	18	24	10	13
		mean \pm SE	40.0 \pm 11.0	49.0 \pm 13*	7.0 \pm 1.5	11.0 \pm 1.5*
	LZ	1	54	70	6	11
		2	16	16	2	3
		3	15	23	9	13
		mean \pm SE	28.0 \pm 12.8	36.0 \pm 16.7	5.7 \pm 2.0	9.0 \pm 3.0
5 IL-3	MPO	1	20	28	23	36
		2	37	42	39	48
		3	5	9	10	35
		mean \pm SE	21.0 \pm 9.0	26.0 \pm 9.0	24.0 \pm 8.3	39.7 \pm 4.2
	LZ	1	15	22	17	29
		2	3	3	3	3
		3	3	5	6	22
		mean \pm SE	7.0 \pm 4.0	10.3 \pm 5.8	8.7 \pm 4.0	18.0 \pm 7.0
GM-CSF + IL-3	MPO	1	29	37	46	56
		2	38	40	49	56
		3	6	10	3	6
		mean \pm SE	24.0 \pm 9.0	39.3 \pm 16.6	32.7 \pm 14.8	39.3 \pm 16.6
	LZ	1	18	20	29	30
		2	2	3	3	3
		3	1	2	1	2
		mean \pm SE	7.0 \pm 5.5	8.3 \pm 5.8	11.0 \pm 9.0	12.0 \pm 9.0
10 GM-CSF + SCF	MPO	1	50	51	15	19
		2	16	21	10	48
		mean \pm SE	33.0 \pm 17.0	36.0 \pm 15.0	12.5 \pm 2.5	33.5 \pm 14.5
	LZ	1	15	15	5	6
		2	9	20	5	45
		mean \pm SE	12.0 \pm 3.0	18.0 \pm 2.0	5.0 \pm 0.0	25.5 \pm 19.5

Among the granulomonocytic markers tested, VEGF-C led to an increase in the proportion of LZ+ cells under all culture conditions. In comparison, LZ+CD14+ cells, which represent differentiated monocytic cells only very slightly increased upon addition of VEGF-C (data not shown). Co-stimulation of the cells with VEGF-C increased the expression of MPO, an early granulocytic marker molecule, only modestly, except in combination with both GM-CSF and IL-3, where the increase in the proportion of MPO+ cells was more pronounced.

VEGF-C exerts co-stimulatory effects in combination with M-CSF

In another series of experiments, CD34+ cells were cultured in medium supplemented with 50 ng/ml M-CSF, with or without 100 ng/ml VEGF-C, for seven days. Culture of CD34+ cells in the presence of M-CSF leads to the generation of CD14+ monocytes within 7 days. After seven days, the cultures were analyzed to determine the percentages of CD14+ cells and mean fluorescence intensity. The results are summarized in the table below:

Percentages of CD14 ⁺ cells and mean fluorescence intensity (MFI) of cells cultured with M-CSF in the absence or in the presence of VEGF-C				
	M-CSF alone		M-CSF + VEGF-C	
exp no	% CD14 ⁺	MFI	% CD14 ⁺	MFI
1	37	20	47	40
2	42	44	54	74
3	32	6	36	7
mean±SE	36.8±2.9	23.3±11.1	45.7±5.2	40.3±19.3

As shown in the table, addition of VEGF-C to these cultures increased both the proportion of CD14+ cells (37% CD14+ cells vs. 46%) and the fluorescence intensity of CD14 expression (MFI 23.3 vs. 40.3). However, cell numbers did not increase upon addition of VEGF-C to M-CSF supplemented cultures. Thus, VEGF-C had a small effect on the differentiation of monocytic cells, but not on their growth.

In the foregoing experiments the presence of VEGF-C was associated with enhanced numbers of cells in cultures of cord blood CD34+ cells. Under all conditions tested (GM-CSF, IL-3, GM-CSF + IL-3; GM-CSF + SCF), co-culture with VEGF-C led

- 103 -

to an enhancement of proportions of myeloid cells. These results indicate an application for VEGF-C in the stimulation and/or differentiation of CD34+ progenitor cells *in vitro* or *in vivo*. Furthermore, the use of VEGF-C alone also slightly increased the number of surviving cells. The results thus indicate uses for compositions comprising VEGF-C
5 prepared in admixture with the aforementioned or other growth factors, such as VEGF-C, and unit dose formulations comprising VEGF-C packaged together with the aforementioned or other growth factors. Such compositions, unit dose formulations, and methods of their use are intended as further aspects of the present invention.

Deposit of Biological Materials: Plasmid FLT4-L has been deposited
10 with the American Type Culture Collection (ATCC), 12301 Parklawn Dr., Rockville MD 20952 (USA), pursuant to the provisions of the Budapest Treaty, and has been assigned a deposit date of 24 July 1995 and ATCC accession number 97231.

While the present invention has been described in terms of specific
embodiments, it is understood that variations and modifications will occur to those in the
15 art. Accordingly, only such limitations as appear in the appended claims should be placed on the invention.

- 104 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Ludwig Institute for Cancer Research
Helsinki University Licensing
Alitalo, Kari (U.S. only)
Joukov, Vladimir (U.S. only)
- (ii) TITLE OF INVENTION: Vascular Endothelial Growth Factor C (VEGF-C)
Protein and Gene, Mutants Thereof, and Uses Thereof
- (iii) NUMBER OF SEQUENCES: 59
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Marshall, O'Toole, Gerstein, Murray & Borun
 - (B) STREET: 6300 Sears Tower, 233 South Wacker Drive
 - (C) CITY: Chicago
 - (D) STATE: Illinois
 - (E) COUNTRY: United States of America
 - (F) ZIP: 60606-6402
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/795,430
 - (B) FILING DATE: 05-FEB-1997
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: PCT/FI96/00427
 - (B) FILING DATE: 01-AUG-1996
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/671,573
 - (B) FILING DATE: 28-JUN-1996
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/601,132
 - (B) FILING DATE: 14-FEB-1996
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/585,895
 - (B) FILING DATE: 12-JAN-1996
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/510,133
 - (B) FILING DATE: 01-AUG-1995

- 105 -

(vii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: 08/340,011
(B) FILING DATE: 14-NOV-1994

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Gass, David A.
(B) REGISTRATION NUMBER: 38,153
(C) REFERENCE/DOCKET NUMBER: 28967/34140

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 312/474-6300
(B) TELEFAX: 312/474-0448
(C) TELEX: 25-3856

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4416 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CCACGCGCAG CGGCCGGAGA TGCAGCGGGG CGCCGCGCTG TGCCTGCGAC TGTGGCTCTG	60
CCTGGGACTC CTGGACGGCC TGGTGAGTGG CTA CTCCATG ACCCCCCCGA CCTTGAACAT	120
CACGGAGGAG TCACACGTCA TCGACACCGG TGACAGCCTG TCCATCTCCT GCAGGGGACA	180
GCACCCCTC GAGTGGGCTT GGCCAGGAGC TCAGGAGGCG CCAGCCACCG GAGACAAGGA	240
CAGCGAGGAC ACGGGGGTGG TGCGAGACTG CGAGGGCACA GACGCCAGGC CCTACTGCAA	300
GGTGTGCTG CTGCACGAGG TACATGCCAA CGACACAGGC AGCTACGTCT GCTACTACAA	360
GTACATCAAG GCACGCATCG AGGGCACCAC GGCCGCCAGC TCCTACGTGT TCGTGAGAGA	420
CTTTGAGCAG CCATTCATCA ACAAGCCTGA CACGCTCTTG GTCAACAGGA AGGACGCCAT	480
GTGGGTGCCC TGTCTGGTGT CCATCCCCGG CCTCAATGTC ACGCTGCGCT CGCAAAGCTC	540
GGTGCTGTGG CCAGACGGGC AGGAGGTGGT GTGGGATGAC CGGCGGGGCA TGCTCGTGTC	600
CACGCCACTG CTGCACGATG CCCTGTACCT GCAGTGCGAG ACCACCTGGG GAGACCAGGA	660
CTTCCTTTCC AACCCTTCC TGGTGACAT CACAGGCAAC GAGCTCTATG ACATCCAGCT	720
GTTGCCCAGG AAGTCGCTGG AGCTGCTGGT AGGGGAGAAG CTGGTCCTGA ACTGCACCGT	780
GTGGGCTGAG TTAACTCAG GTGTCACCTT TGA CTGGGAC TACCCAGGGA AGCAGGCAGA	840

GCGGGGTAAG TGGGTGCCCC AGCGACGCTC CCAGCAGACC CACACAGAAC TCTCCAGCAT	900
CCTGACCATC CACAACGTCA GCCAGCACGA CCTGGGCTCG TATGTGTGCA AGGCCAACAA	960
CGGCATCCAG CGATTTCGGG AGAGCACCGA GGTCAATTGTG CATGAAAATC CCTTCATCAG	1020
CGTCGAGTGG CTCAAAGGAC CCATCCTGGA GGCCACGGCA GGAGACGAGC TGGTGAAGCT	1080
GCCCCGTGAAG CTGGCAGCGT ACCCCCCGCC CGAGTTCCAG TGGTACAAGG ATGGAAAGGC	1140
ACTGTCCGGG CGCCACAGTC CACATGCCCT GGTGCTCAAG GAGGTGACAG AGGCCAGCAC	1200
AGGCACCTAC ACCCTCGCCC TGTGGAATC CGCTGCTGGC CTGAGGCGCA ACATCAGCCT	1260
GGAGCTGGTG GTGAATGTGC CCCCCAGAT ACATGAGAAG GAGGCCTCCT CCCCCAGCAT	1320
CTACTCGCGT CACAGCCGCC AGGCCCTCAC CTGCACGGCC TACGGGGTGC CCCTGCCTCT	1380
CAGCATCCAG TGGCACTGGC GGCCCTGGAC ACCCTGCAAG ATGTTTGCCC AGCGTAGTCT	1440
CCGGCGGCGG CAGCAGCAAG ACCTCATGCC ACAGTGCCGT GACTGGAGGG CGGTGACCAC	1500
GCAGGATGCC GTGAACCCCA TCGAGAGCCT GGACACCTGG ACCGAGTTTG TGGAGGGAAA	1560
GAATAAGACT GTGAGCAAGC TGGTGATCCA GAATGCCAAC GTGTCTGCCA TGTACAAGTG	1620
TGTGGTCTCC AACAAGGTGG GCCAGGATGA GCGGCTCATC TACTTCTATG TGACCACCAT	1680
CCCCGACGGC TTCACCATCG AATCCAAGCC ATCCGAGGAG CTACTAGAGG GCCAGCCGGT	1740
GCTCCTGAGC TGCCAAGCCG ACAGCTACAA GTACGAGCAT CTGCGCTGGT ACCGCCTCAA	1800
CCTGTCCACG CTGCACGATG CGCACGGGAA CCCGCTTCTG CTCGACTGCA AGAACGTGCA	1860
TCTGTTTCGCC ACCCCTCTGG CCGCCAGCCT GGAGGAGGTG GCACCTGGGG CGCGCCACGC	1920
CACGCTCAGC CTGAGTATCC CCCGCTCGC GCCCGAGCAC GAGGGCCACT ATGTGTGCGA	1980
AGTGCAAGAC CGGCGCAGCC ATGACAAGCA CTGCCACAAG AAGTACCTGT CGGTGCAGGC	2040
CCTGGAAGCC CCTCGGCTCA CGCAGAACTT GACCGACCTC CTGGTGAACG TGAGCGACTC	2100
GCTGGAGATG CAGTGCTTGG TGGCCGGAGC GCACGCGCCC AGCATCGTGT GGTACAAAGA	2160
CGAGAGGCTG CTGGAGGAAA AGTCTGGAGT CGACTTGCGG GACTCCAACC AGAAGCTGAG	2220
CATCCAGCGC GTGCGCGAGG AGGATGCGGG ACGCTATCTG TGCAGCGTGT GCAACGCCAA	2280
GGGCTGCGTC AACTCCTCCG CCAGCGTGCC CGTGGAAGGC TCCGAGGATA AGGGCAGCAT	2340
GGAGATCGTG ATCCTTGTCG GTACCGGCGT CATCGCTGTC TTCTTCTGGG TCCTCCTCCT	2400
CCTCATCTTC TGTAACATGA GGAGGCCGGC CCACGCAGAC ATCAAGACGG GCTACCTGTC	2460
CATCATCATG GACCCCGGGG AGGTGCCTCT GGAGGAGCAA TGCGAATACC TGTCTTACGA	2520
TGCCAGCCAG TGGGAATTCC CCCGAGAGCG GCTGCACCTG GGGAGAGTGC TCGGCTACGG	2580

- 107 -

CGCCTTCGGG AAGGTGGTGG AAGCCTCCGC TTTCGGCATC CACAAGGGCA GCAGCTGTGA	2640
CACCGTGGCC GTGAAAATGC TGAAAGAGGG CGCCACGGCC AGCGAGCACC GCGCGCTGAT	2700
GTCGGAGCTC AAGATCCTCA TTCACATCGG CAACCACCTC AACGTGGTCA ACCTCCTCGG	2760
GGCGTGCACC AAGCCGCAGG GCCCCCTCAT GGTGATCGTG GAGTTCTGCA AGTACGGCAA	2820
CCTCTCCAAC TTCCTGCGCG CCAAGCGGGA CGCCTTCAGC CCCTGCGCGG AGAAGTCTCC	2880
CGAGCAGCGC GGACGCTTCC GCGCCATGGT GGAGCTCGCC AGGCTGGATC GGAGGCGGCC	2940
GGGGAGCAGC GACAGGGTCC TCTTCGCGCG GTTCTCGAAG ACCGAGGGCG GAGCGAGGCG	3000
GGCTTCTCCA GACCAAGAAG CTGAGGACCT GTGGCTGAGC CCGCTGACCA TGGAAGATCT	3060
TGTCTGCTAC AGCTTCCAGG TGGCCAGAGG GATGGAGTTC CTGGCTTCCC GAAAGTGCAT	3120
CCACAGAGAC CTGGCTGCTC GGAACATTCT GCTGTCGGAA AGCGACGTGG TGAAGATCTG	3180
TGACTTTGGC CTTGCCCCGG ACATCTACAA AGACCCTGAC TACGTCCGCA AGGGCAGTGC	3240
CCGGCTGCCC CTGAAGTGGA TGGCCCCTGA AAGCATCTTC GACAAGGTGT ACACCACGCA	3300
GAGTGACGTG TGGTCCTTTG GGGTGCTTCT CTGGGAGATC TTCTCTCTGG GGGCCTCCCC	3360
GTACCCTGGG GTGCAGATCA ATGAGGAGTT CTGCCAGCGG CTGAGAGACG GCACAAGGAT	3420
GAGGGCCCCG GAGCTGGCCA CTCCCGCCAT ACGCCGCATC ATGCTGAACT GCTGGTCCGG	3480
AGACCCCAAG GCGAGACCTG CATTCTCGGA GCTGGTGGAG ATCCTGGGGG ACCTGCTCCA	3540
GGGCAGGGGC CTGCAAGAGG AAGAGGAGGT CTGCATGGCC CCGCGCAGCT CTCAGAGCTC	3600
AGAAGAGGGC AGCTTCTCGC AGGTGTCCAC CATGGCCCTA CACATCGCCC AGGCTGACGC	3660
TGAGGACAGC CCGCCAAGCC TGCAGCGCCA CAGCCTGGCC GCCAGGTATT ACAACTGGGT	3720
GTCTTTTCCC GGGTGCCTGG CCAGAGGGGC TGAGACCCGT GGTTCCTCCA GGATGAAGAC	3780
ATTTGAGGAA TTCCCCATGA CCCCACGAC CTACAAAGGC TCTGTGGACA ACCAGACAGA	3840
CAGTGGGATG GTGCTGGCCT CGGAGGAGTT TGAGCAGATA GAGAGCAGGC ATAGACAAGA	3900
AAGCGGCTTC AGGTAGCTGA AGCAGAGAGA GAGAAGGCAG CACATCGTCCG CATTTTCTTC	3960
TCTGCACTTA TAAGAAAGAT CAAAGACTTT AAGACTTTTCG CTATTTCTTC TACTGCTATC	4020
TACTACAAAC TTCAAAGAGG AACCAGGAGG ACAAGAGGAG CATGAAAGTG GACAAGGAGT	4080
GTGACCACTG AAGCACCACA GGGAAGGGGT TAGGCCTCCG GATGACTGCG GGCAGGCCTG	4140
GATAATATCC AGCCTCCAC AAGAAGCTGG TGGAGCAGAG TGTTCCCTGA CTCCTCCAAG	4200
GAAAGGGAGA CGCCCTTTCA TGGTCTGCTG AGTAACAGGT GCNTTCCCAG ACACTGGCGT	4260
TACTGCTTGA CCAAAGAGCC CTCAAGCGGC CCTTATGCCA GCGTGACAGA GGGCTCACCT	4320

- 108 -

CTTGCCTTCT AGGTCAC TTC TCACACAATG TCCCTTCAGC ACCTGACCCT GTGCCCCGCA 4380
 GTTATTCCTT GGTAATATGA GTAATACATC AAAGAG 4416

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 216 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CAAGAAAGCG GCTTCAGCTG TAAAGGACCT GGCCAGAATG TGGCTGTGAC CAGGGCACAC 60
 CCTGACTCCC AAGGGAGGCG GCGGCGGCCT GAGCGGGGGG CCCGAGGAGG CCAGGTGTTT 120
 TACAACAGCG AGTATGGGGA GCTGTGCGGAG CCAAGCGAGG AGGACCACTG CTCCCCGTCT 180
 GCGCGCGTGA CTTTCTTCAC AGACAACAGC TACTAA 216

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4273 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

AAGCTTATCG ATTTTGAACC CGGGGGTACC GAATTCCTCG AGTCTAGAGG AGCATGCCTG 60
 CAGGTCGACC GGGCTCGATC CCCTCGCGAG TTGGTTCAGC TGCTGCCTGA GGCTGGACGA 120
 CCTCGCGGAG TTCTACCGGC AGTGCAAATC CGTCGGCATC CAGGAAACCA GCAGCGGCTA 180
 TCCGCGCATC CATGCCCCCG AACTGCAGGA GTGGGGAGGC ACGATGGCCG CTTTGGTCCC 240
 GGATCTTTGT GAAGGAACCT TACTTCTGTG GTGTGACATA ATTGGACAAA CTACCTACAG 300
 AGATTTAAAG CTCTAAGGTA AATATAAAAT TTTTAAGTGT ATAATGTGTT AAACACTAGA 360
 TTCTAATTGT TTGTGTATTT TAGATTCCAA CCTATGGAAC TGATGAATGG GAGCAGTGGT 420
 GGAATGCCTT TAATGAGGAA AACCTGTTTT GCTCAGAAGA AATGCCATCT AGTGATGATG 480

- 109 -

AGGCTACTGC TGACTCTCAA CATTCTACTC CTCCAAAAAA GAAGAGAAAG GTAGAAGACC	540
CCAAGGACTT TCCTTCAGAA TTGCTAAGTT TTTTGAGTCA TGCTGTGTTT AGTAATAGAA	600
CTCTTGCTTG CTTTGCTATT TACACCACAA AGGAAAAAGC TGCACTGCTA TACAAGAAAA	660
TTATGGAAAA ATATTCTGTA ACCTTTATAA GTAGGCATAA CAGTTATAAT CATAACATAC	720
TGTTTTTTCT TACTCCACAC AGGCATAGAG TGTCTGCTAT TAATAACTAT GCTCAAAAAT	780
TGTGTACCTT TAGCTTTTTA ATTTGTAAAG GGGTTAATAA GGAATATTTG ATGTATAGTG	840
CCTTGACTAG AGATCATAAT CAGCCATACC ACATTGTAG AGGTTTTACT TGCTTTAAAA	900
AACCTCCCAC ACCTCCCCCT GAACCTGAAA CATAAAATGA ATGCAATTGT TGTGTGTAAC	960
TTGTTTATTG CAGCTTATAA TGGTTACAAA TAAAGCAATA GCATCACAAA TTTCACAAAT	1020
AAAGCATTTT TTTCAGTGCA TTCTAGTTGT GGTGTGTTCA AACTCATCAA TGTATCTTAT	1080
CATGTCTGGA TCTGCCGGTC TCCCTATAGT GAGTCGTATT AATTTGATA AGCCAGGTTA	1140
ACCTGCATTA ATGAATCGGC CAACGCGCGG GGAGAGGCGG TTTGCGTATT GGGCGCTCTT	1200
CCGCTTCCTC GCTCACTGAC TCGCTGCGCT CGGTGCTTCG GCTGCGGCGA GCGGTATCAG	1260
CTCACTCAA GGCGGTAATA CGGTTATCCA CAGAATCAGG GGATAACGCA GGAAAGAACA	1320
TGTGAGCAA AGGCCAGCAA AAGGCCAGGA ACCGTAAAAA GGACGCGTTG CTGGCGTTTT	1380
TCCATAGGCT CCGCCCCCT GACGAGCATC AAAAAATCG ACGCTCAAGT CAGAGGTGGC	1440
GAAACCCGAC AGGACTATAA AGATACCAGG CGTTTCCCCC TGGAAGCTCC CTCGTGCGCT	1500
CTCCTGTTCC GACCCTGCCG CTTACCGGAT ACCTGTCCGC CTTTCTCCCT TCGGGAAGCG	1560
TGGCGCTTTC TCAATGCTCA CGCTGTAGGT ATCTCAGTTC GGTGTAGGTC GTTCGCTCCA	1620
AGCTGGGCTG TGTGCACGAA CCCCCGTTT AGCCCGACCG CTGCGCCTTA TCCGGTAACT	1680
ATCGTCTTGA GTCCAACCCG GTAAGACACG ACTTATCGCC ACTGGCAGCA GCCACTGGTA	1740
ACAGGATTAG CAGAGCGAGG TATGTAGGCG GTGCTACAGA GTTCTTGAAG TGGTGGCCTA	1800
ACTACGGCTA CACTAGAAGG ACAGTATTTG GTATCTGCGC TCTGCTGAAG CCAGTTACCT	1860
TCGGAAAAAG AGTTGGTAGC TCTTGATCCG GCAAACAAAC CACCGCTGGT AGCGGTGGTT	1920
TTTTTGTTG CAAGCAGCAG ATTACGCGCA GAAAAAAGG ATCTCAAGAA GATCCTTTGA	1980
TCTTTTCTAC GGGGTCTGAC GCTCAGTGA ACGAAACTC ACGTTAAGG ATTTTGGTCA	2040
TGAGATTATC AAAAAGGATC TTCACCTAGA TCCTTTTAAA TTAAAAATGA AGTTTTAAAT	2100
CAATCTAAAG TATATATGAG TAACTTGGT CTGACAGTTA CCAATGCTTA ATCAGTGAGG	2160
CACCTATCTC AGCGATCTGT CTATTTCTGT CATCCATAGT TGCCTGACTC CCCGTCGTGT	2220

- 110 -

AGATAACTAC GATACGGGAG GGCTTACCAT CTGGCCCCAG TGCTGCAATG ATACCGCGAG	2280
ACCCACGCTC ACCGGCTCCA GATTTATCAG CAATAAACCA GCCAGCCGGA AGGGCCGAGC	2340
GCAGAAGTGG TCCTGCAACT TTATCCGCCT CCATCCAGTC TATTAATTGT TGCCGGGAAG	2400
CTAGAGTAAG TAGTTCGCCA GTTAATAGTT TGCGCAACGT TGTTGCCATT GCTACAGGCA	2460
TCGTGGTGTC ACGCTCGTCG TTTGGTATGG CTTCAATTCAG CTCCGGTTCC CAACGATCAA	2520
GGCGAGTTAC ATGATCCCCC ATGTTGTGCA AAAAAGCGGT TAGCTCCTTC GGTCTCCGA	2580
TCGTTGTCAG AAGTAAGTTG GCCGCAGTGT TATCACTCAT GGTATGGCA GCACTGCATA	2640
ATTCTCTTAC TGTCATGCCA TCCGTAAGAT GCTTTTCTGT GACTGGTGAG TACTCAACCA	2700
AGTCATTCTG AGAATAGTGT ATGCGGCGAC CGAGTTGCTC TTGCCCGGCG TCAATACGGG	2760
ATAATACCGC GCCACATAGC AGAACTTTAA AAGTGCTCAT CATTGGAAAA CGTTCCTTCGG	2820
GGCGAAAACT CTCAAGGATC TTACCGCTGT TGAGATCCAG TTCGATGTAA CCCACTCGTG	2880
CACCCAACCTG ATCTTCAGCA TCTTTTACTT TCACCAGCGT TTCTGGGTGA GCAAAAACAG	2940
GAAGGCAAAA TGCCGCAAAA AAGGGAATAA GGGCGACACG GAAATGTTGA ATACTCATAC	3000
TCTTCCTTTT TCAATATTAT TGAAGCATTT ATCAGGGTTA TTGTCTCATG AGCGGATACA	3060
TATTTGAATG TATTTAGAAA AATAAACAAA TAGGGGTTCC GCGCACATTT CCCCAGAAAG	3120
TGCCACCTGA CGTCTAAGAA ACCATTATTA TCATGACATT AACCTATAAA AATAGGCGTA	3180
TCACGAGGCC CTTTCGTCTC GCGCGTTTCG GTGATGACGG TGAAAACCTC TGACACATGC	3240
AGTCCCCGGA GACGGTCACA GCTTGCTGTG AAGCGGATGC CGGGAGCAGA CAAGCCCGTC	3300
AGGGCGCGTC AGCGGGTGTT GCGGGGTGTC GGGGCTGGCT TAACTATGCG GCATCAGAGC	3360
AGATTGTACT GAGAGTGCAC CATATGGACA TATTGTCGTT AGAACGCGGC TACAATTAAT	3420
ACATAACCTT ATGTATCATA CACATACGAT TTAGGTGACA CTATAGAACT CGAGCAGAGC	3480
TTCCAAATTG AGAGAGAGGC TTAATCAGAG ACAGAAACTG TTTGAGTCAA CTCAAGGATG	3540
GTTTGAGGGA CTGTTTAACA GATCCCCTTG GTTTACCACC TTGATATCTA CCATTATGGG	3600
ACCCCTCATT GTACTCCTAA TGATTTTGCT CTTCCGACCC TGCATTCTTA ATCGATTAGT	3660
CCAATTTGTT AAAGACAGGA TATCAGTGGT CCAGGCTCTA GTTTTGA CTC AACAAATATCA	3720
CCAGCTGAAG CCTATAGAGT ACGAGCCATA GATAAAATAA AAGATTTTAT TTAGTCTCCA	3780
GAAAAAGGGG GGAATGAAAG ACCCCACCTG TAGGTTTGCC AAGCTAGCTT AAGTAACGCC	3840
ATTTTGCAAG GCATGGAAAA ATACATAACT GAGAATAGAG AAGTTCAGAT CAAGGTCAGG	3900
AACAGATGGA ACAGCTGAAT ATGGGCCAAA CAGGATATCT GTGGTAAGCA GTTCCTGCCC	3960

- 111 -

CGGCTCAGGG CCAAGAACAG ATGGAACAGC TGAATATGGG CCAAACAGGA TATCTGTGGT 4020
 AAGCAGTTCC TGCCCCGGCT CAGGGCCAAG AACAGATGGT CCCCAGATGC GGTCCAGCCC 4080
 TCAGCAGTTT CTAGAGAACC ATCAGATGTT TCCAGGGTGC CCCAAGGACC TGAAATGACC 4140
 CTGTGCCTTA TTTGAACTAA CCAATCAGTT CGCTTCTCGC TTCTGTTCGC GCGCTTCTGC 4200
 TCCCCGAGCT CAATAAAAGA GCCCACAACC CCTCACTCGG GCGCCAGTC CTCCGATTGA 4260
 CTGAGTCGCC CGG 4273

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Pro Met Thr Pro Thr Thr Tyr Lys Gly Ser Val Asp Asn Gln Thr Asp
 1 5 10 15
 Ser Gly Met Val Leu Ala Ser Glu Glu Phe Glu Gln Ile Glu Ser Arg
 20 25 30
 His Arg Gln Glu Ser Gly Phe Arg
 35 40

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Xaa Glu Glu Thr Ile Lys Phe Ala Ala Ala His Tyr Asn Thr Glu Ile
 1 5 10 15
 Leu Lys

- 112 -

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 219 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TCACTATAGG GAGACCCAAG CTTGGTACCG AGCTCGGATC CACTAGTAAC GGCCGCCAGT	60
GTGGTGGAAT TCGACGAACT CATGACTGTA CTCTACCCAG AATATTGGAA AATGTACAAG	120
TGTCAGCTAA GGCAAGGAGG CTGGCAACAT AACAGAGAAC AGGCCAACCT CAACTCAAGG	180
ACAGAAGAGA CTATAAAATT CGCTGCAGCA CACTACAAC	219

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1997 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 352..1608

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CCCGCCCCGC CTCTCCAAAA AGCTACACCG ACGCGGACCG CGGCGGCGTC CTCCCTCGCC	60
CTCGCTTCAC CTCGCGGGCT CCGAATGCGG GGAGCTCGGA TGTCCGTTT CCTGTGAGGC	120
TTTACCTGA CACCCGCCGC CTTTCCCCGG CACTGGCTGG GAGGGCGCCC TGCAAAGTTG	180
GGAACGCGGA GCCCCGGACC CGCTCCCGCC GCCTCCGGCT CGCCAGGGG GGGTCGCCGG	240
GAGGAGCCCG GGGGAGAGGG ACCAGGAGGG GCCCGCGGCC TCGCAGGGGC GCCCGCGCCC	300
CCACCCCTGC CCCC GCCAGC GGACCGGTCC CCCACCCCCG GTCCTTCCAC C ATG CAC	357
	Met His

- 113 -

TTG CTG GGC TTC TTC TCT GTG GCG TGT TCT CTG CTC GCC GCT GCG CTG Leu Leu Gly Phe Phe Ser Val Ala Cys Ser Leu Leu Ala Ala Leu	405
5 10 15	
CTC CCG GGT CCT CGC GAG GCG CCC GCC GCC GCC GCC GCC TTC GAG TCC Leu Pro Gly Pro Arg Glu Ala Pro Ala Ala Ala Ala Ala Phe Glu Ser	453
20 25 30	
GGA CTC GAC CTC TCG GAC GCG GAG CCC GAC GCG GGC GAG GCC ACG GCT Gly Leu Asp Leu Ser Asp Ala Glu Pro Asp Ala Gly Glu Ala Thr Ala	501
35 40 45 50	
TAT GCA AGC AAA GAT CTG GAG GAG CAG TTA CGG TCT GTG TCC AGT GTA Tyr Ala Ser Lys Asp Leu Glu Glu Gln Leu Arg Ser Val Ser Val	549
55 60 65	
GAT GAA CTC ATG ACT GTA CTC TAC CCA GAA TAT TGG AAA ATG TAC AAG Asp Glu Leu Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys Met Tyr Lys	597
70 75 80	
TGT CAG CTA AGG AAA GGA GGC TGG CAA CAT AAC AGA GAA CAG GCC AAC Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln Ala Asn	645
85 90 95	
CTC AAC TCA AGG ACA GAA GAG ACT ATA AAA TTT GCT GCA GCA CAT TAT Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala His Tyr	693
100 105 110	
AAT ACA GAG ATC TTG AAA AGT ATT GAT AAT GAG TGG AGA AAG ACT CAA Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys Thr Gln	741
115 120 125 130	
TGC ATG CCA CGG GAG GTG TGT ATA GAT GTG GGG AAG GAG TTT GGA GTC Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe Gly Val	789
135 140 145	
GCG ACA AAC ACC TTC TTT AAA CCT CCA TGT GTG TCC GTC TAC AGA TGT Ala Thr Asn Thr Phe Phe Lys Pro Pro Cys Val Ser Val Tyr Arg Cys	837
150 155 160	
GGG GGT TGC TGC AAT AGT GAG GGG CTG CAG TGC ATG AAC ACC AGC ACG Gly Gly Cys Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr Ser Thr	885
165 170 175	
AGC TAC CTC AGC AAG ACG TTA TTT GAA ATT ACA GTG CCT CTC TCT CAA Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu Ser Gln	933
180 185 190	
GGC CCC AAA CCA GTA ACA ATC AGT TTT GCC AAT CAC ACT TCC TGC CGA Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser Cys Arg	981
195 200 205 210	
TGC ATG TCT AAA CTG GAT GTT TAC AGA CAA GTT CAT TCC ATT ATT AGA Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile Ile Arg	1029
215 220 225	

- 114 -

CGT TCC CTG CCA GCA ACA CTA CCA CAG TGT CAG GCA GCG AAC AAG ACC Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn Lys Thr 230 235 240	1077
TGC CCC ACC AAT TAC ATG TGG AAT AAT CAC ATC TGC AGA TGC CTG GCT Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys Leu Ala 245 250 255	1125
CAG GAA GAT TTT ATG TTT TCC TCG GAT GCT GGA GAT GAC TCA ACA GAT Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser Thr Asp 260 265 270	1173
GGA TTC CAT GAC ATC TGT GGA CCA AAC AAG GAG CTG GAT GAA GAG ACC Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu Glu Thr 275 280 285 290	1221
TGT CAG TGT GTC TGC AGA GCG GGG CTT CGG CCT GCC AGC TGT GGA CCC Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys Gly Pro 295 300 305	1269
CAC AAA GAA CTA GAC AGA AAC TCA TGC CAG TGT GTC TGT AAA AAC AAA His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys Asn Lys 310 315 320	1317
CTC TTC CCC AGC CAA TGT GGG GCC AAC CGA GAA TTT GAT GAA AAC ACA Leu Phe Pro Ser Gln Cys Gly Ala Asn Arg Glu Phe Asp Glu Asn Thr 325 330 335	1365
TGC CAG TGT GTA TGT AAA AGA ACC TGC CCC AGA AAT CAA CCC CTA AAT Cys Gln Cys Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro Leu Asn 340 345 350	1413
CCT GGA AAA TGT GCC TGT GAA TGT ACA GAA AGT CCA CAG AAA TGC TTG Pro Gly Lys Cys Ala Cys Glu Cys Thr Glu Ser Pro Gln Lys Cys Leu 355 360 365 370	1461
TTA AAA GGA AAG AAG TTC CAC CAC CAA ACA TGC AGC TGT TAC AGA CGG Leu Lys Gly Lys Lys Phe His His Gln Thr Cys Ser Cys Tyr Arg Arg 375 380 385	1509
CCA TGT ACG AAC CGC CAG AAG GCT TGT GAG CCA GGA TTT TCA TAT AGT Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro Gly Phe Ser Tyr Ser 390 395 400	1557
GAA GAA GTG TGT CGT TGT GTC CCT TCA TAT TGG AAA AGA CCA CAA ATG Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp Lys Arg Pro Gln Met 405 410 415	1605
AGC TAAGATTGTA CTGTTTTCCA GTTCATCGAT TTTCTATTAT GGAAACTGT Ser	1658
GTTGCCACAG TAGAACTGTC TGTGAACAGA GAGACCCTTG TGGGTCCATG CTAACAAAGA	1718
CAAAAGTCTG TCTTTCTGA ACCATGTGGA TAACTTTACA GAAATGGACT GGAGCTCATC	1778
TGCAAAAGGC CTCTTGTAAG GACTGGTTTT CTGCCAATGA CCAAACAGCC AAGATTTTCC	1838

- 115 -

TCTTGTGATT TCTTTAAAAG AATGACTATA TAATTTATTT CCACTAAAAA TATTGTTTCT 1898
 GCATTCATTT TTATAGCAAC AACCAATTGGT AAAACTCACT GTGATCAATA TTTTATATC 1958
 ATGCAAAAATA TGTTTAAAAT AAAATGAAAA TTGTATTAT 1997

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 419 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met His Leu Leu Gly Phe Phe Ser Val Ala Cys Ser Leu Leu Ala Ala
 1 5 10 15
 Ala Leu Leu Pro Gly Pro Arg Glu Ala Pro Ala Ala Ala Ala Phe
 20 25 30
 Glu Ser Gly Leu Asp Leu Ser Asp Ala Glu Pro Asp Ala Gly Glu Ala
 35 40 45
 Thr Ala Tyr Ala Ser Lys Asp Leu Glu Glu Gln Leu Arg Ser Val Ser
 50 55 60
 Ser Val Asp Glu Leu Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys Met
 65 70 75 80
 Tyr Lys Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln
 85 90 95
 Ala Asn Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala
 100 105 110
 His Tyr Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys
 115 120 125
 Thr Gln Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe
 130 135 140
 Gly Val Ala Thr Asn Thr Phe Phe Lys Pro Pro Cys Val Ser Val Tyr
 145 150 155 160
 Arg Cys Gly Gly Cys Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr
 165 170 175
 Ser Thr Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu
 180 185 190
 Ser Gln Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser
 195 200 205

- 116 -

Cys Arg Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile
 210 215 220
 Ile Arg Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn
 225 230 235 240
 Lys Thr Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys
 245 250 255
 Leu Ala Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser
 260 265 270
 Thr Asp Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu
 275 280 285
 Glu Thr Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys
 290 295 300
 Gly Pro His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys
 305 310 315 320
 Asn Lys Leu Phe Pro Ser Gln Cys Gly Ala Asn Arg Glu Phe Asp Glu
 325 330 335
 Asn Thr Cys Gln Cys Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro
 340 345 350
 Leu Asn Pro Gly Lys Cys Ala Cys Glu Cys Thr Glu Ser Pro Gln Lys
 355 360 365
 Cys Leu Leu Lys Gly Lys Lys Phe His His Gln Thr Cys Ser Cys Tyr
 370 375 380
 Arg Arg Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro Gly Phe Ser
 385 390 395 400
 Tyr Ser Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp Lys Arg Pro
 405 410 415
 Gln Met Ser

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 117 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Glu Glu Thr Ile Lys Phe Ala Ala Ala His Tyr Asn Thr Glu Ile Leu
1 5 10 15

Lys

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1836 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) **FEATURE:**

(A) NAME/KEY: CDS

(B) LOCATION: 168..1412

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GCGGCCGCGT	CGACGCAAAA	GTTGCGAGCC	GCCGAGTCCC	GGGAGACGCT	CGCCCCAGGGG		60
GGTCCCCGGG	AGGAAACCAC	GGGACAGGGA	CCAGGAGAGG	ACCTCAGCCT	CACGCCCCAG		120
CCTGCGCCAG	CCAACGGACC	GGCCTCCCTG	CTCCCGGTCC	ATCCACC	ATG CAC TTG Met His Leu 1		176
CTG TGC TTC TTG TCT CTG GCG TGT TCC CTG CTC GCC GCT GCG CTG ATC Leu Cys Phe Leu Ser Leu Ala Cys Ser Leu Leu Ala Ala Ala Leu Ile 5 10 15							224
CCC AGT CCG CGC GAG GCG CCC GCC ACC GTC GCC GCC TTC GAG TCG GGA Pro Ser Pro Arg Glu Ala Pro Ala Thr Val Ala Ala Phe Glu Ser Gly 20 25 30 35							272
CTG GGC TTC TCG GAA GCG GAG CCC GAC GGG GGC GAG GTC AAG GCT TTT Leu Gly Phe Ser Glu Ala Glu Pro Asp Gly Gly Glu Val Lys Ala Phe 40 45 50							320
GAA GGC AAA GAC CTG GAG GAG CAG TTG CGG TCT GTG TCC AGC GTA GAT Glu Gly Lys Asp Leu Glu Glu Gln Leu Arg Ser Val Ser Ser Val Asp 55 60 65							368
GAG CTG ATG TCT GTC CTG TAC CCA GAC TAC TGG AAA ATG TAC AAG TGC Glu Leu Met Ser Val Leu Tyr Pro Asp Tyr Trp Lys Met Tyr Lys Cys 70 75 80							416
CAG CTG CGG AAA GGC GGC TGG CAG CAG CCC ACC CTC AAT ACC AGG ACA Gln Leu Arg Lys Gly Gly Trp Gln Gln Pro Thr Leu Asn Thr Arg Thr 85 90 95							464

- 118 -

GGG GAC AGT GTA AAA TTT GCT GCT GCA CAT TAT AAC ACA GAG ATC CTG Gly Asp Ser Val Lys Phe Ala Ala Ala His Tyr Asn Thr Glu Ile Leu 100 105 110 115	512
AAA AGT ATT GAT AAT GAG TGG AGA AAG ACT CAA TGC ATG CCA CGT GAG Lys Ser Ile Asp Asn Glu Trp Arg Lys Thr Gln Cys Met Pro Arg Glu 120 125 130	560
GTG TGT ATA GAT GTG GGG AAG GAG TTT GGA GCA GCC ACA AAC ACC TTC Val Cys Ile Asp Val Gly Lys Glu Phe Gly Ala Ala Thr Asn Thr Phe 135 140 145	608
TTT AAA CCT CCA TGT GTG TCC GTC TAC AGA TGT GGG GGT TGC TGC AAC Phe Lys Pro Pro Cys Val Ser Val Tyr Arg Cys Gly Gly Cys Cys Asn 150 155 160	656
AGC GAG GGG CTG CAG TGC ATG AAC ACC AGC ACA GGT TAC CTC AGC AAG Ser Glu Gly Leu Gln Cys Met Asn Thr Ser Thr Gly Tyr Leu Ser Lys 165 170 175	704
ACG TTG TTT GAA ATT ACA GTG CCT CTC TCA CAA GGC CCC AAA CCA GTC Thr Leu Phe Glu Ile Thr Val Pro Leu Ser Gln Gly Pro Lys Pro Val 180 185 190 195	752
ACA ATC AGT TTT GCC AAT CAC ACT TCC TGC CGG TGC ATG TCT AAA CTG Thr Ile Ser Phe Ala Asn His Thr Ser Cys Arg Cys Met Ser Lys Leu 200 205 210	800
GAT GTT TAC AGA CAA GTT CAT TCA ATT ATT AGA CGT TCT CTG CCA GCA Asp Val Tyr Arg Gln Val His Ser Ile Ile Arg Arg Ser Leu Pro Ala 215 220 225	848
ACA TTA CCA CAG TGT CAG GCA GCT AAC AAG ACA TGT CCA ACA AAC TAT Thr Leu Pro Gln Cys Gln Ala Ala Asn Lys Thr Cys Pro Thr Asn Tyr 230 235 240	896
GTG TGG AAT AAC TAC ATG TGC CGA TGC CTG GCT CAG CAG GAT TTT ATC Val Trp Asn Asn Tyr Met Cys Arg Cys Leu Ala Gln Gln Asp Phe Ile 245 250 255	944
TTT TAT TCA AAT GTT GAA GAT GAC TCA ACC AAT GGA TTC CAT GAT GTC Phe Tyr Ser Asn Val Glu Asp Asp Ser Thr Asn Gly Phe His Asp Val 260 265 270 275	992
TGT GGA CCC AAC AAG GAG CTG GAT GAA GAC ACC TGT CAG TGT GTC TGC Cys Gly Pro Asn Lys Glu Leu Asp Glu Asp Thr Cys Gln Cys Val Cys 280 285 290	1040
AAG GGG GGG CTT CGG CCA TCT AGT TGT GGA CCC CAC AAA GAA CTA GAT Lys Gly Gly Leu Arg Pro Ser Ser Cys Gly Pro His Lys Glu Leu Asp 295 300 305	1088
AGA GAC TCA TGT CAG TGT GTC TGT AAA AAC AAA CTT TTC CCT AAT TCA Arg Asp Ser Cys Gln Cys Val Cys Lys Asn Lys Leu Phe Pro Asn Ser 310 315 320	1136

- 119 -

TGT GGA GCC AAC AGG GAA TTT GAT GAG AAT ACA TGT CAG TGT GTA TGT Cys Gly Ala Asn Arg Glu Phe Asp Glu Asn Thr Cys Gln Cys Val Cys 325 330 335	1184
AAA AGA ACG TGT CCA AGA AAT CAG CCC CTG AAT CCT GGG AAA TGT GCC Lys Arg Thr Cys Pro Arg Asn Gln Pro Leu Asn Pro Gly Lys Cys Ala 340 345 350 355	1232
TGT GAA TGT ACA GAA AAC ACA CAG AAG TGC TTC CTT AAA GGG AAG AAG Cys Glu Cys Thr Glu Asn Thr Gln Lys Cys Phe Leu Lys Gly Lys Lys 360 365 370	1280
TTC CAC CAT CAA ACA TGC AGT TGT TAC AGA AGA CCG TGT GCG AAT CGA Phe His His Gln Thr Cys Ser Cys Tyr Arg Arg Pro Cys Ala Asn Arg 375 380 385	1328
CTG AAG CAT TGT GAT CCA GGA CTG TCC TTT AGT GAA GAA GTA TGC CGC Leu Lys His Cys Asp Pro Gly Leu Ser Phe Ser Glu Glu Val Cys Arg 390 395 400	1376
TGT GTC CCA TCG TAT TGG AAA AGG CCA CAT CTG AAC TAAGATCATA Cys Val Pro Ser Tyr Trp Lys Arg Pro His Leu Asn 405 410 415	1422
CCAGTTTTCA GTCAGTCACA GTCATTTACT CTCTTGAAGA CTGTTGGAAC AGCACTTAGC	1482
ACTGTCTATG CACAGAAAGA CTCTGTGGGA CCACATGGTA ACAGAGGCCC AAGTCTGTGT	1542
TTATTGAACC ATGTGGATTA CTGCGGGAGA GGA CTGGCAC TCATGTGCAA AAAAAACCTC	1602
TTCAAAGACT GGTCTTCTGC CAGGGACCAG ACAGCTGAGG TTTTCTCTT GTGATTTAAA	1662
AAAAGAATGA CTATATAATT TATTTCCACT AAAAATATTG TTCCTGCATT CATTTTTATA	1722
GCAATAACAA TTGGTAAAGC TCACTGTGAT CAGTATTTTT ATAACATGCA AACTATGTT	1782
TAAAATAAAA TGAAAATTGT ATTATAAAAA AAAAAAAAAA AAAAAAAAAA GCTT	1836

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 415 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Met His Leu Leu Cys Phe Leu Ser Leu Ala Cys Ser Leu Leu Ala Ala 1 5 10 15
Ala Leu Ile Pro Ser Pro Arg Glu Ala Pro Ala Thr Val Ala Ala Phe 20 25 30

- 120 -

Glu Ser Gly Leu Gly Phe Ser Glu Ala Glu Pro Asp Gly Gly Glu Val
 35 40 45
 Lys Ala Phe Glu Gly Lys Asp Leu Glu Glu Gln Leu Arg Ser Val Ser
 50 55 60
 Ser Val Asp Glu Leu Met Ser Val Leu Tyr Pro Asp Tyr Trp Lys Met
 65 70 75 80
 Tyr Lys Cys Gln Leu Arg Lys Gly Gly Trp Gln Gln Pro Thr Leu Asn
 85 90 95
 Thr Arg Thr Gly Asp Ser Val Lys Phe Ala Ala Ala His Tyr Asn Thr
 100 105 110
 Glu Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys Thr Gln Cys Met
 115 120 125
 Pro Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe Gly Ala Ala Thr
 130 135 140
 Asn Thr Phe Phe Lys Pro Pro Cys Val Ser Val Tyr Arg Cys Gly Gly
 145 150 155 160
 Cys Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr Ser Thr Gly Tyr
 165 170 175
 Leu Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu Ser Gln Gly Pro
 180 185 190
 Lys Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser Cys Arg Cys Met
 195 200 205
 Ser Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile Ile Arg Arg Ser
 210 215 220
 Leu Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn Lys Thr Cys Pro
 225 230 235 240
 Thr Asn Tyr Val Trp Asn Asn Tyr Met Cys Arg Cys Leu Ala Gln Gln
 245 250 255
 Asp Phe Ile Phe Tyr Ser Asn Val Glu Asp Asp Ser Thr Asn Gly Phe
 260 265 270
 His Asp Val Cys Gly Pro Asn Lys Glu Leu Asp Glu Asp Thr Cys Gln
 275 280 285
 Cys Val Cys Lys Gly Gly Leu Arg Pro Ser Ser Cys Gly Pro His Lys
 290 295 300
 Glu Leu Asp Arg Asp Ser Cys Gln Cys Val Cys Lys Asn Lys Leu Phe
 305 310 315 320
 Pro Asn Ser Cys Gly Ala Asn Arg Glu Phe Asp Glu Asn Thr Cys Gln
 325 330 335

- 121 -

Cys Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro Leu Asn Pro Gly
340 345 350

Lys Cys Ala Cys Glu Cys Thr Glu Asn Thr Gln Lys Cys Phe Leu Lys
355 360 365

Gly Lys Lys Phe His His Gln Thr Cys Ser Cys Tyr Arg Arg Pro Cys
370 375 380

Ala Asn Arg Leu Lys His Cys Asp Pro Gly Leu Ser Phe Ser Glu Glu
385 390 395 400

Val	Cys	Arg	Cys	Val	Pro	Ser	Tyr	Trp	Lys	Arg	Pro	His	Leu	Asn
				405					410					415

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1741 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) **FEATURE:**

- (A) NAME/KEY: CDS
(B) LOCATION: 453..1706

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GCCCCCGCCG	AGCGCTCCGC	GCGCAGCCGC	CGGGCCGGGC	CGGCCCCGCG	AGGGCGCGCT		60
GCGAGCGGCC	ACTGGGTCCT	GCTTCCCTCC	TTCCTCTCCC	TCCTCCTCCT	CCTCCTTCTC		120
TCTGCGCTTT	CCACCGCTCC	CGAGCGAGCG	CACGCTCGGA	TGTCCGGTTT	CCTGGTGGGT		180
TTTTTACCTG	GCAAAGTCCG	GATAACTTCG	GTGAGAAATT	GCAAAGAGGC	TGGGAGCTCC		240
CCTGCAGGCG	TCTGGGAGCT	GCTGCCGCCG	TCGCATCTTC	TCCATCCCGC	GGATTTTACT		300
GCCTTGATA	TTGCGAGGGG	AGGGAGGGGG	GTGAGGACAG	CAAAAAGAAA	GGGGTGGGGG		360
GGGGGAGAGA	AAAGGAAAAG	AAGGAGCCTC	GGAATTGTGC	CCGCATTCTT	GCGCTGCCCC		420
GCGGCCCCCC	TCCGCTCTGC	CATCTCCGCA	CA ATG CAC	TTG CTG GAG	ATG CTC		473
			Met His	Leu Leu	Glu Met Leu		
			1		5		
TCC CTG GGC	TGC TGC CTC	GCT GCT GGC	GCC GTG CTC	CTG GGA CCC	CGG		521
Ser Leu Gly	Cys Cys Leu	Ala Ala Gly	Ala Val Leu	Leu Leu Gly	Pro Arg		
10		15		20			

- 122 -

CAG CCG CCC GTC GCC GCC GCC TAC GAG TCC GGG CAC GGC TAC TAC GAG Gln Pro Pro Val Ala Ala Ala Tyr Glu Ser Gly His Gly Tyr Tyr Glu 25 30 35	569
GAG GAG CCC GGT GCC GGG GAA CCC AAG GCT CAT GCA AGC AAA GAC CTG Glu Glu Pro Gly Ala Gly Glu Pro Lys Ala His Ala Ser Lys Asp Leu 40 45 50 55	617
GAA GAG CAG TTG CGA TCT GTG TCC AGT GTG GAT GAA CTC ATG ACA GTA Glu Glu Gln Leu Arg Ser Val Ser Ser Val Asp Glu Leu Met Thr Val 60 65 70	665
CTT TAC CCA GAA TAC TGG AAA ATG TTC AAA TGT CAG TTG AGG AAA GGA Leu Tyr Pro Glu Tyr Trp Lys Met Phe Lys Cys Gln Leu Arg Lys Gly 75 80 85	713
GGT TGG CAA CAC AAC AGG GAA CAC TCC AGC TCT GAT ACA AGA TCA GAT Gly Trp Gln His Asn Arg Glu His Ser Ser Ser Asp Thr Arg Ser Asp 90 95 100	761
GAT TCA TTG AAA TTT GCC GCA GCA CAT TAT AAT GCA GAG ATC CTG AAA Asp Ser Leu Lys Phe Ala Ala His Tyr Asn Ala Glu Ile Leu Lys 105 110 115	809
AGT ATT GAT ACT GAA TGG AGA AAA ACC CAG GGC ATG CCA CGT GAA GTG Ser Ile Asp Thr Glu Trp Arg Lys Thr Gln Gly Met Pro Arg Glu Val 120 125 130 135	857
TGT GTG GAT TTG GGG AAA GAG TTT GGA GCA ACT ACA AAC ACC TTC TTT Cys Val Asp Leu Gly Lys Glu Phe Gly Ala Thr Thr Asn Thr Phe Phe 140 145 150	905
AAA CCC CCG TGT GTA TCC ATC TAC AGA TGT GGA GGT TGC TGC AAT AGT Lys Pro Pro Cys Val Ser Ile Tyr Arg Cys Gly Gly Cys Cys Asn Ser 155 160 165	953
GAA GGA CTC CAG TGT ATG AAT ATC AGC ACA AAT TAC ATC AGC AAG ACA Glu Gly Leu Gln Cys Met Asn Ile Ser Thr Asn Tyr Ile Ser Lys Thr 170 175 180	1001
TTG TTT GAG ATT ACA GTG CCT CTC TCT CAT GGC CCC AAA CCT GTA ACA Leu Phe Glu Ile Thr Val Pro Leu Ser His Gly Pro Lys Pro Val Thr 185 190 195	1049
GTC AGT TTT GCC AAT CAC ACG TCC TGC CGA TGC ATG TCT AAG TTG GAT Val Ser Phe Ala Asn His Thr Ser Cys Arg Cys Met Ser Lys Leu Asp 200 205 210 215	1097
GTT TAC AGA CAA GTT CAT TCT ATC ATA AGA CGT TCC TTG CCA GCA ACA Val Tyr Arg Gln Val His Ser Ile Ile Arg Arg Ser Leu Pro Ala Thr 220 225 230	1145
CAA ACT CAG TGT CAT GTG GCA AAC AAG ACC TGT CCA AAA AAT CAT GTC Gln Thr Gln Cys His Val Ala Asn Lys Thr Cys Pro Lys Asn His Val 235 240 245	1193

- 123 -

TGG AAT AAT CAG ATT TGC AGA TGC TTA GCA CAG CAC GAT TTT GGT TTC	1241
Trp Asn Asn Gln Ile Cys Arg Cys Leu Ala Gln His Asp Phe Gly Phe	
250 255 260	
TCT TCT CAC CTT GGA GAT TCT GAC ACA TCT GAA GGA TTC CAT ATT TGT	1289
Ser Ser His Leu Gly Asp Ser Asp Thr Ser Glu Gly Phe His Ile Cys	
265 270 275	
GGG CCC AAC AAA GAG CTG GAT GAA GAA ACC TGT CAA TGC GTC TGC AAA	1337
Gly Pro Asn Lys Glu Leu Asp Glu Glu Thr Cys Gln Cys Val Cys Lys	
280 285 290 295	
GGA GGT GTG CGG CCC ATA AGC TGT GGC CCT CAC AAA GAA CTA GAC AGG	1385
Gly Gly Val Arg Pro Ile Ser Cys Gly Pro His Lys Glu Leu Asp Arg	
300 305 310	
GCA TCA TGT CAG TGC ATG TGC AAA AAC AAA CTG CTC CCC AGT TCC TGT	1433
Ala Ser Cys Gln Cys Met Cys Lys Asn Lys Leu Leu Pro Ser Ser Cys	
315 320 325	
GGG CCT AAC AAA GAA TTT GAT GAA GAA AAG TGC CAG TGT GTA TGT AAA	1481
Gly Pro Asn Lys Glu Phe Asp Glu Glu Lys Cys Gln Cys Val Cys Lys	
330 335 340	
AAG ACC TGT CCC AAA CAT CAT CCA CTA AAT CCT GCA AAA TGC ATC TGC	1529
Lys Thr Cys Pro Lys His His Pro Leu Asn Pro Ala Lys Cys Ile Cys	
345 350 355	
GAA TGT ACA GAA TCT CCC AAT AAA TGT TTC TTA AAA GGA AAA AGA TTT	1577
Glu Cys Thr Glu Ser Pro Asn Lys Cys Phe Leu Lys Gly Lys Arg Phe	
360 365 370 375	
CAT CAC CAG ACA TGC AGT TGT TAC AGA CCA CCA TGT ACA GTC CGA ACG	1625
His His Gln Thr Cys Ser Cys Tyr Arg Pro Pro Cys Thr Val Arg Thr	
380 385 390	
AAA CGC TGT GAT GCT GGA TTT CTG TTA GCT GAA GAA GTG TGC CGC TGT	1673
Lys Arg Cys Asp Ala Gly Phe Leu Leu Ala Glu Glu Val Cys Arg Cys	
395 400 405	
GTA CGC ACA TCT TGG AAA AGA CCA CTT ATG AAT TAAGCGAAGA AAGCACTACT	1726
Val Arg Thr Ser Trp Lys Arg Pro Leu Met Asn	
410 415	
CGCTATATAG TGTCG	1741

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 418 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 124 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Met	His	Leu	Leu	Glu	Met	Leu	Ser	Leu	Gly	Cys	Cys	Leu	Ala	Ala	Gly	1	5	10	15
Ala	Val	Leu	Leu	Gly	Pro	Arg	Gln	Pro	Pro	Val	Ala	Ala	Ala	Tyr	Glu	20	25	30	
Ser	Gly	His	Gly	Tyr	Tyr	Glu	Glu	Glu	Pro	Gly	Ala	Gly	Glu	Pro	Lys	35	40	45	
Ala	His	Ala	Ser	Lys	Asp	Leu	Glu	Glu	Gln	Leu	Arg	Ser	Val	Ser	Ser	50	55	60	
Val	Asp	Glu	Leu	Met	Thr	Val	Leu	Tyr	Pro	Glu	Tyr	Trp	Lys	Met	Phe	65	70	75	80
Lys	Cys	Gln	Leu	Arg	Lys	Gly	Gly	Trp	Gln	His	Asn	Arg	Glu	His	Ser	85	90	95	
Ser	Ser	Asp	Thr	Arg	Ser	Asp	Asp	Ser	Leu	Lys	Phe	Ala	Ala	Ala	His	100	105	110	
Tyr	Asn	Ala	Glu	Ile	Leu	Lys	Ser	Ile	Asp	Thr	Glu	Trp	Arg	Lys	Thr	115	120	125	
Gln	Gly	Met	Pro	Arg	Glu	Val	Cys	Val	Asp	Leu	Gly	Lys	Glu	Phe	Gly	130	135	140	
Ala	Thr	Thr	Asn	Thr	Phe	Phe	Lys	Pro	Pro	Cys	Val	Ser	Ile	Tyr	Arg	145	150	155	160
Cys	Gly	Gly	Cys	Cys	Asn	Ser	Glu	Gly	Leu	Gln	Cys	Met	Asn	Ile	Ser	165	170	175	
Thr	Asn	Tyr	Ile	Ser	Lys	Thr	Leu	Phe	Glu	Ile	Thr	Val	Pro	Leu	Ser	180	185	190	
His	Gly	Pro	Lys	Pro	Val	Thr	Val	Ser	Phe	Ala	Asn	His	Thr	Ser	Cys	195	200	205	
Arg	Cys	Met	Ser	Lys	Leu	Asp	Val	Tyr	Arg	Gln	Val	His	Ser	Ile	Ile	210	215	220	
Arg	Arg	Ser	Leu	Pro	Ala	Thr	Gln	Thr	Gln	Cys	His	Val	Ala	Asn	Lys	225	230	235	240
Thr	Cys	Pro	Lys	Asn	His	Val	Trp	Asn	Asn	Gln	Ile	Cys	Arg	Cys	Leu	245	250	255	
Ala	Gln	His	Asp	Phe	Gly	Phe	Ser	Ser	His	Leu	Gly	Asp	Ser	Asp	Thr	260	265	270	
Ser	Glu	Gly	Phe	His	Ile	Cys	Gly	Pro	Asn	Lys	Glu	Leu	Asp	Glu	Glu	275	280	285	

- 125 -

Thr Cys Gln Cys Val Cys Lys Gly Gly Val Arg Pro Ile Ser Cys Gly
 290 295 300
 Pro His Lys Glu Leu Asp Arg Ala Ser Cys Gln Cys Met Cys Lys Asn
 305 310 315 320
 Lys Leu Leu Pro Ser Ser Cys Gly Pro Asn Lys Glu Phe Asp Glu Glu
 325 330 335
 Lys Cys Gln Cys Val Cys Lys Lys Thr Cys Pro Lys His His Pro Leu
 340 345 350
 Asn Pro Ala Lys Cys Ile Cys Glu Cys Thr Glu Ser Pro Asn Lys Cys
 355 360 365
 Phe Leu Lys Gly Lys Arg Phe His His Gln Thr Cys Ser Cys Tyr Arg
 370 375 380
 Pro Pro Cys Thr Val Arg Thr Lys Arg Cys Asp Ala Gly Phe Leu Leu
 385 390 395 400
 Ala Glu Glu Val Cys Arg Cys Val Arg Thr Ser Trp Lys Arg Pro Leu
 405 410 415
 Met Asn

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Ala Val Val Met Thr Gln Thr Pro Ala Ser
 1 5 10

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- 126 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

TCTCTTCTGT GCTTGAGTTG AG

22

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

TCTCTTCTGT CCCTGAGTTG AG

22

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 65 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

TGTGCTGCAG CAAATTTTAT AGTCTCTTCT GTGGCGGCGG CGGCGGCGGG CGCCTCGCGA

60

GGACC

65

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CTGGCAGGGA ACTGCTAATA ATGGAATGAA

30

- 127 -

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 84 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GGGCTCCGCG TCCGAGAGGT CGAGTCCGGA CTCGTGATGG TGATGGTGAT GGGCGGCGGC 60
GGCGGCGGGC GCCTCGCGAG GACC 84

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GTATTATAAT GTCCTCCACC AAATTTTATA G 31

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 93 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GTTGCTGCC TGACACTGTG GTAGTGTTC TGGCGGCCGC TAGTGATGGT GATGGTGATG 60
AATAATGGAA TGAAGTTGTC TGTAACATC CAG 93

(2) INFORMATION FOR SEQ ID NO:22:

- 128 -

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CATGTACGAA CCGCCAGG

18

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

AATGACCAGA GAGAGGCGAG

20

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GCCACGGTAG GTCTGCGT

18

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- 129 -

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

TTTCTTTGAC AGGCTTAT

18

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

ATCTTGAAAA GTAAGTATGG G

21

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

ATGACTTGAC AGGTATTGAT

20

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

- 130 -

AGCAAGACGG TGGGTATTGT

20

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

CCCTTCTTTG TAGTTATTTG AA

22

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

CCACAGTGAG TATGAATTAA

20

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

TTCTTCCAAA GGTGTCAG

18

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs

- 131 -

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

GGAGATGGTA GCAGAATG

18

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 23 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

CTATTTGTCT AGACTCAACA GAT

23

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

CAAACATGCA GGTAAGAGAT CC

22

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 132 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

TGTTCTCCTA GCTGTTACAG A

21

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

GGCGAGGTCA AGGTAGGTGC AAGG

24

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

ATTGTCTTTG ACAGGCTTTT TGAAGG

26

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

GAGATCCTGA AAAGTAAGTA G

21

- 133 -

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

TGTGACTCGA CAGGTATTGA TAAT

24

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

CTCAGCAAGA CGGTAGGTAT

20

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

TTGTCCCTTG TAGTTGTTTG AAATT

25

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

- 134 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

ACATTACCAC AGTGAGTATG

20

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

GTCTCCCCAA AAGGTGTCAG GCAGCT

26

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

AATGTTGAAG ATGGTAAGTA AAA

23

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 135 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

TCTAGACTCA ACCAAT

16

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

CAAACATGCA GGTAAGGAGT GT

22

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

TTTTCCCCTA GTTGTTACAG AAGA

24

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2991 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

GTTTTAAGTA GAGACGGGGT TTCACCAACG GTTGAAAATA TTTATCATGG TCTCCCTAAG

60

ATGGACGGTG TTAGCTAGGA TGGTCTCGAT CTCCTGACCT CATGATCCAC CCGCCTCGGC

120

CTCCCAAAGT GCTGGGATTA CAGGCGTGAG CCACCGTGTC CGACCAACCT TAAGACAAAC

180

- 136 -

AACTACTGCA TGATTGTTTT TGGAGACCTT TTTTTTATTC AAATAAATTT TTGCCAGCAT	240
TTTCTGACTC AAAGTATAGC AGCAGGAAGA TAACACTTTT GTGAGAAAAA AGTTTGAATA	300
CAGCTTACTG CTGTATTTAA ATGAAACAGT AGTTAATATG ATATTAATAT ATTTTGGATA	360
TATTTTGAGT TTGTTGATTT TCCAGTCTTC ACCCGCTGCT AGGCCTGTGG GTGTTGGAAA	420
TGCCTGTGTT TCTCAATTTT GTTGCCTAT TAGAATCCTG ATGTCCAAGC CTTACTCCAG	480
TTAGACCAGT TAAGCCAGAA AGGCAGAAGG TGTACTCAAG CATCTGTTTT TTCAAAATCT	540
CCTTTTGTGA TGCCAAGTGC AATCAAAGTT TAGAATCATT GTAATAGCAA ATGGTTGAAT	600
GGAAACTCCA CCTTCTATTC AAATCCTACC CCAGTCTGCC CTTAGCTGTT CTCTTTTCAC	660
AGATCTATCA ATGTCTGAAG ATAACATATG CAGGCTGATC AAATATGCAT AGAGCAGGAA	720
GACAGCAAGA GAGTGATACA CTGACCATGT TCCAAATCAC AAAACATCTC AACAGGCTAG	780
ATCATGGACC GAGTCTGATG GGATGGAATT TCATAAAGAT ACATAAAAAA GCATCTTGGA	840
TACAGTAAAC TTAACTCCAC AAATACAGGG GAATTTAGAC GTGACTAAGT AGCAGTACAT	900
ATGAAAAATT ATTGAGGAAT TTTGTTGACT TTAAGGGTAG TGTGAGTCAA CACTGTGATT	960
TGGCTGCCAG AAAATAAACT CAATCCAAGG CTGTATCAAC AAAGGCATAC TGTCCATTCT	1020
GCATGCTCAT TACAGCACTA AGTACCGAGC CATGTTCTCA ACCGCATACT TCATGAACAT	1080
GGAAAGCTAA CAGTATGGTT AAGGGGGGAA ACTGGAAGTGC TCATCTTGGG GAATAAAAGG	1140
GATATTTAGC CAGGAGTAAA GTTAGCTTAG GGAGACCATG ATAAATATTT TCAAAATATT	1200
TGAAGGACTC AGTTGTGGAA GTGAGATTAG ATTTATTGTG TAAAACTCCA GGAGTCAAAA	1260
GCAATAGAGA GATAGAAGGA AATGCTTTTC AGCAGTGTG CTCATCAATA AAGGGAGTGA	1320
ACAGCCACAC AGAATGGAAG GTTCCCTGTC CTTTGAGATA TTTAAGCCTT CAAGTAAATT	1380
ATGGGTGAGG AGTTTCAAAT CTAGAGTTGA ACCAGATAAG AAAGTCTCTT CTTCCGGTAA	1440
GATATTATGG ACCTATAACA TCTGTGTACT TAAAAGTAGA TTGGGAGTGA AAGGCAGACT	1500
TTTGATGTTT TGTACACTGT TGAAACCCCT TAGCGTGGTC CTCTGTAACC TGCTCACCCCT	1560
GCCCCAAGGA GGCAGCTAGC CAATGCCACC AGCCCAACGG AAACCCAGT GCTTTTCCAA	1620
TGGGGAAATG CAGTCACTTT TCTTTGGATG CTACACATCC TTTCTGGAAT ATGTCTCACA	1680
CACATCTCTC TTTATCACCC CCTTTTTCAT GTAAACCAAC TTCTTGCGA AGCTGACAAAT	1740
GTGTCTCTTT ACTCTCCACG AAGATTCTGG CCCTTCTCTT CACCTGTCAG AAGTTTAGGA	1800
TTCCAAAGGG ATCATTAGCA TCCATCCCAA CAGCCTGCAC TGCATCCTGA GAACTGCGGT	1860
TCTTGGATCA TCAGGCAACT TTCAACTACA CAGACCAAGG GAGAGAGGGG ACCCCTCCGA	1920

- 137 -

GGTCCCATAG GGTTCCTCTGA CATAGTGATG ACCTTTTTTCC AAACTTTGAG CAGGGCGCTG	1980
GGGGCCAGGC GTGCGGGAGG GAGGACAAGA ACTCGGGAGT GGCCGAGGAT AAAGCGGGGG	2040
CTCCCTCCAC CCCACGGTGC CCAGTTTCTC CCCGCTGCAC GTGGTCCAGG GTGGTCGCAT	2100
CACCTCTAAA GCCGGTCCCG CCAACCGCCA GCCCCGGGAC TGAAC TTGCC CCTCCGGCCG	2160
CCCGCTCCCC GCAGGGGACA GGGGCGGGGA GGGAGAGATC CAGAGGGGGG CTGGGGGAGG	2220
TGGGGCCGCC GGGGAGGAGG CGAGGGAAAC GGGGAGCTCC AGGGAGACGG CTTCCGAGGG	2280
AGAGTGAGAG GGGAGGGCAG CCCGGGCTCG GCACGCTCCC TCCCTCGGCC GCTTTCTCTC	2340
ACATAAGCGC AGGCAGAGGG CGCGTCAGTC ATGCCCTGCC CCTGCGCCCG CCGCCGCCGC	2400
CGCCGCCGCT CAGCCCGGCG CGCTCTGGAG GATCCTGCGC CGCGGCGCTC CCGGGCCCCG	2460
CCGCCGCCAG CCGCCCCGGC GGCCCTCCTC CCGCCCCCGG CACCGCCGCC AGCGCCCCCG	2520
CCGCAGCGCC CGCGGCCCGG CTCCTCTCAC TTCGGGGAAG GGGAGGGAGG AGGGGGACGA	2580
GGGCTCTGGC GGGTTTGGAG GGGCTGAACA TCGCGGGGTG TTCTGGTGTC CCCC GCCCG	2640
CCTCTCCAAA AAGCTACACC GACGCGGACC GCGGCGGCGT CCTCCCTCGC CCTCGCTTCA	2700
CCTCGCGGGC TCCGAATGCG GGGAGCTCGG ATGTCCGGTT TCCTGTGAGG CTTTTACCTG	2760
ACACCCGCCG CTTTTCCCCG GCACTGGCTG GGAGGGCGCC CTGCAAAGTT GGGAACGCGG	2820
AGCCCCGGAC CCGCTCCCGC CGCCTCCGGC TCGCCAGGG GGGGTCGCCG GGAGGAGCCC	2880
GGGGGAGAGG GACCAGGAGG GGCCCGCGGC CTCGCAGGGG CGCCCGCGCC CCCACCCCTG	2940
CCCCCGCCAG CGGACCGGTC CCCACCCCC GGTCTTCCA CCATGCACTT G	2991

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

CACGGCTTAT GCAAGCAAAG

20

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs

- 138 -

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

AACACAGTTT TCCATAATAG

20

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

Leu Ser Lys Thr Val Ser Gly Ser Glu Gln Asp Leu Pro His Glu Leu
1 5 10 15
His Val Glu

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

GACGGACACA GATGGAGGTT TAAAG

25

(2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 196 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant

- 139 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

```

Met Arg Thr Leu Ala Cys Leu Leu Leu Leu Gly Cys Gly Tyr Leu Ala
1           5           10           15
His Val Leu Ala Glu Glu Ala Glu Ile Pro Arg Glu Val Ile Glu Arg
20           25           30
Leu Ala Arg Ser Gln Ile His Ser Ile Arg Asp Leu Gln Arg Leu Leu
35           40           45
Glu Ile Asp Ser Val Gly Ser Glu Asp Ser Leu Asp Thr Ser Leu Arg
50           55           60
Ala His Gly Val His Ala Thr Lys His Val Pro Glu Lys Arg Pro Leu
65           70           75           80
Pro Ile Arg Arg Lys Arg Ser Ile Glu Glu Ala Val Pro Ala Val Cys
85           90           95
Lys Thr Arg Thr Val Ile Tyr Glu Ile Pro Arg Ser Gln Val Asp Pro
100          105          110
Thr Ser Ala Asn Phe Leu Ile Trp Pro Pro Cys Val Glu Val Lys Arg
115          120          125
Cys Thr Gly Cys Cys Asn Thr Ser Ser Val Lys Cys Gln Pro Ser Arg
130          135          140
Val His His Arg Ser Val Lys Val Ala Lys Val Glu Tyr Val Arg Lys
145          150          155          160
Lys Pro Lys Leu Lys Glu Val Gln Val Arg Leu Glu Glu His Leu Glu
165          170          175
Cys Ala Cys Ala Thr Thr Ser Leu Asn Pro Asp Tyr Arg Glu Glu Asp
180          185          190
Thr Asp Val Arg
195

```

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 241 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 140 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

Met Asn Arg Cys Trp Ala Leu Phe Leu Ser Leu Cys Cys Tyr Leu Arg
 1 5 10 15
 Leu Val Ser Ala Glu Gly Asp Pro Ile Pro Glu Glu Leu Tyr Glu Met
 20 25 30
 Leu Ser Asp His Ser Ile Arg Ser Phe Asp Asp Leu Gln Arg Leu Leu
 35 40 45
 His Gly Asp Pro Gly Glu Glu Asp Gly Ala Glu Leu Asp Leu Asn Met
 50 55 60
 Thr Arg Ser His Ser Gly Gly Glu Leu Glu Ser Leu Ala Arg Gly Arg
 65 70 75 80
 Arg Ser Leu Gly Ser Leu Thr Ile Ala Glu Pro Ala Met Ile Ala Glu
 85 90 95
 Cys Lys Thr Arg Thr Glu Val Phe Glu Ile Ser Arg Arg Leu Ile Asp
 100 105 110
 Arg Thr Asn Ala Asn Phe Leu Val Trp Pro Pro Cys Val Glu Val Gln
 115 120 125
 Arg Cys Ser Gly Cys Cys Asn Asn Arg Asn Val Gln Cys Arg Pro Thr
 130 135 140
 Gln Val Gln Leu Arg Pro Val Gln Val Arg Lys Ile Glu Ile Val Arg
 145 150 155 160
 Lys Lys Pro Ile Phe Lys Lys Ala Thr Val Thr Leu Glu Asp His Leu
 165 170 175
 Ala Cys Lys Cys Glu Thr Val Ala Ala Ala Arg Pro Val Thr Arg Ser
 180 185 190
 Pro Gly Gly Ser Gln Glu Gln Arg Ala Lys Thr Pro Gln Thr Arg Val
 195 200 205
 Thr Ile Arg Thr Val Arg Val Arg Arg Pro Pro Lys Gly Lys His Arg
 210 215 220
 Lys Phe Lys His Thr His Asp Lys Thr Ala Leu Lys Glu Thr Leu Gly
 225 230 235 240
 Ala

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- 141 -

- (A) LENGTH: 149 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

```

Met Pro Val Met Arg Leu Phe Pro Cys Phe Leu Gln Leu Leu Ala Gly
 1             5             10             15

Leu Ala Leu Pro Ala Val Pro Pro Gln Gln Trp Ala Leu Ser Ala Gly
      20             25             30

Asn Gly Ser Ser Glu Val Glu Val Val Pro Phe Gln Glu Val Trp Gly
      35             40             45

Arg Ser Tyr Cys Arg Ala Leu Glu Arg Leu Val Asp Val Val Ser Glu
      50             55             60

Tyr Pro Ser Glu Val Glu His Met Phe Ser Pro Ser Cys Val Ser Leu
      65             70             75             80

Leu Arg Cys Thr Gly Cys Cys Gly Asp Glu Asn Leu His Cys Val Pro
      85             90             95

Val Glu Thr Ala Asn Val Thr Met Gln Leu Leu Lys Ile Arg Ser Gly
      100            105            110

Asp Arg Pro Ser Tyr Val Glu Leu Thr Phe Ser Gln His Val Arg Cys
      115            120            125

Glu Cys Arg Pro Leu Arg Glu Lys Met Lys Pro Glu Arg Cys Gly Asp
      130            135            140

Ala Val Pro Arg Arg
      145

```

(2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 191 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

- 142 -

Met Asn Phe Leu Leu Ser Trp Val His Trp Ser Leu Ala Leu Leu Leu
 1 5 10 15

Tyr Leu His His Ala Lys Trp Ser Gln Ala Ala Pro Met Ala Glu Gly
 20 25 30

Gly Gly Gln Asn His His Glu Val Val Lys Phe Met Asp Val Tyr Gln
 35 40 45

Arg Ser Tyr Cys His Pro Ile Glu Thr Leu Val Asp Ile Phe Gln Glu
 50 55 60

Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys Pro Ser Cys Val Pro Leu
 65 70 75 80

Met Arg Cys Gly Gly Cys Cys Asn Asp Glu Gly Leu Glu Cys Val Pro
 85 90 95

Thr Glu Glu Ser Asn Ile Thr Met Gln Ile Met Arg Ile Lys Pro His
 100 105 110

Gln Gly Gln His Ile Gly Glu Met Ser Phe Leu Gln His Asn Lys Cys
 115 120 125

Glu Cys Arg Pro Lys Lys Asp Arg Ala Arg Gln Glu Asn Pro Cys Gly
 130 135 140

Pro Cys Ser Glu Arg Arg Lys His Leu Phe Val Gln Asp Pro Gln Thr
 145 150 155 160

Cys Lys Cys Ser Cys Lys Asn Thr Asp Ser Arg Cys Lys Ala Arg Gln
 165 170 175

Leu Glu Leu Asn Glu Arg Thr Cys Arg Cys Asp Lys Pro Arg Arg
 180 185 190

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 188 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

Met Ser Pro Leu Leu Arg Arg Leu Leu Leu Ala Ala Leu Leu Gln Leu
 1 5 10 15

Ala Pro Ala Gln Ala Pro Val Ser Gln Pro Asp Ala Pro Gly His Gln
 20 25 30

- 143 -

Arg	Lys	Val	Val	Ser	Trp	Ile	Asp	Val	Tyr	Thr	Arg	Ala	Thr	Cys	Gln
	35						40					45			
Pro	Arg	Glu	Val	Val	Val	Pro	Leu	Thr	Val	Glu	Leu	Met	Gly	Thr	Val
	50					55					60				
Ala	Lys	Gln	Leu	Val	Pro	Ser	Cys	Val	Thr	Val	Gln	Arg	Cys	Gly	Gly
65					70					75					80
Cys	Cys	Pro	Asp	Asp	Gly	Leu	Glu	Cys	Val	Pro	Thr	Gly	Gln	His	Gln
				85					90					95	
Val	Arg	Met	Gln	Ile	Leu	Met	Ile	Arg	Tyr	Pro	Ser	Ser	Gln	Leu	Gly
			100					105					110		
Glu	Met	Ser	Leu	Glu	Glu	His	Ser	Gln	Cys	Glu	Cys	Arg	Pro	Lys	Lys
		115					120					125			
Lys	Asp	Ser	Ala	Val	Lys	Pro	Asp	Ser	Pro	Arg	Pro	Leu	Cys	Pro	Arg
	130					135					140				
Cys	Thr	Gln	His	His	Gln	Arg	Pro	Asp	Pro	Arg	Thr	Cys	Arg	Cys	Arg
145					150					155					160
Cys	Arg	Arg	Arg	Ser	Phe	Leu	Arg	Cys	Gln	Gly	Arg	Gly	Leu	Glu	Leu
				165					170					175	
Asn	Pro	Asp	Thr	Cys	Arg	Cys	Arg	Lys	Leu	Arg	Arg				
			180					185							

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 419 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) **FEATURE:**

- (A) NAME/KEY: other
(B) LOCATION: 156
(D) OTHER INFORMATION: /note= "codon 156 can be anything other than cysteine, or can be nothing"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

Met His Leu Leu Gly Phe Phe Ser Val Ala Cys Ser Leu Leu Ala Ala
1 5 10 15

- 144 -

Ala Leu Leu Pro Gly Pro Arg Glu Ala Pro Ala Ala Ala Ala Phe
 20 25 30
 Glu Ser Gly Leu Asp Leu Ser Asp Ala Glu Pro Asp Ala Gly Glu Ala
 35 40 45
 Thr Ala Tyr Ala Ser Lys Asp Leu Glu Glu Gln Leu Arg Ser Val Ser
 50 55 60
 Ser Val Asp Glu Leu Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys Met
 65 70 75 80
 Tyr Lys Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln
 85 90 95
 Ala Asn Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala
 100 105 110
 His Tyr Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys
 115 120 125
 Thr Gln Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe
 130 135 140
 Gly Val Ala Thr Asn Thr Phe Phe Lys Pro Pro Xaa Val Ser Val Tyr
 145 150 155 160
 Arg Cys Gly Gly Cys Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr
 165 170 175
 Ser Thr Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu
 180 185 190
 Ser Gln Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser
 195 200 205
 Cys Arg Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile
 210 215 220
 Ile Arg Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn
 225 230 235 240
 Lys Thr Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys
 245 250 255
 Leu Ala Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser
 260 265 270
 Thr Asp Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu
 275 280 285
 Glu Thr Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys
 290 295 300
 Gly Pro His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys
 305 310 315 320

[illegible]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 160 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

Met His Leu Leu Gly Phe Phe Ser Val Ala Cys Ser Leu Leu Ala Ala
1 5 10 15
Ala Leu Leu Pro Gly Pro Arg Glu Ala Pro Ala Ala Ala Ala Ala Thr
20 25 30
Glu Glu Thr Ile Lys Phe Ala Ala Ala His Tyr Asn Thr Glu Ile Leu
35 40 45
Lys Ser Ile Asp Asn Glu Trp Arg Lys Thr Gln Cys Met Pro Arg Glu
50 55 60
Val Cys Ile Asp Val Gly Lys Glu Phe Gly Val Ala Thr Asn Thr Phe
65 70 75 80
Phe Lys Pro Pro Cys Val Ser Val Tyr Arg Cys Gly Gly Cys Cys Asn
85 90 95
Ser Glu Gly Leu Gln Cys Met Asn Thr Ser Thr Ser Tyr Leu Ser Lys
100 105 110

- 146 -

Thr Leu Phe Glu Ile Thr Val Pro Leu Ser Gln Gly Pro Lys Pro Val
115 120 125

Thr Ile Ser Phe Ala Asn His Thr Ser Cys Arg Cys Met Ser Lys Leu
130 135 140

Asp Val Tyr Arg Gln Val His Ser Ile Ile His His His His His His
145 150 155 160

- 147 -

CLAIMS

1. A purified and isolated polypeptide selected from the group consisting of:
 - (a) a purified and isolated polypeptide capable of binding to at least one of KDR receptor tyrosine kinase (VEGFR-2) and Flt4 receptor tyrosine kinase (VEGFR-3), said polypeptide comprising a portion of the amino acid sequence in SEQ ID NO: 8 effective to permit such binding;
 - (b) a purified and isolated VEGF-C of vertebrate origin, wherein said VEGF-C has a molecular weight of about 21-23 kD or about 30-32 kD, as assessed by SDS-PAGE under reducing conditions, and wherein said VEGF-C is capable of binding to Flt4 receptor tyrosine kinase (VEGFR-3);
 - (c) a purified polypeptide analog of human VEGF-C that is capable of binding to at least one of Flt-1 receptor tyrosine kinase (VEGFR-1), KDR receptor tyrosine kinase (VEGFR-2), and Flt4 receptor tyrosine kinase (VEGFR-3); and
 - (d) a polypeptide analog of human VEGF that is capable of binding to at least one of VEGFR-1, VEGFR-2, and VEGFR-3, wherein a cysteine residue is introduced in the VEGF amino acid sequence at a position selected from residues 53 to 63 of the human VEGF165 precursor having the amino acid sequence set forth in SEQ ID NO: 56.
2. A purified and isolated polypeptide according to claim 1 that is capable of binding to at least one of KDR receptor tyrosine kinase (VEGFR-2) and Flt4 receptor tyrosine kinase (VEGFR-3), said polypeptide comprising a portion of the amino acid sequence in SEQ ID NO: 8 effective to permit such binding.
3. A polypeptide according to claim 1 or 2, wherein said polypeptide is capable of stimulating tyrosine phosphorylation of a receptor selected from the group consisting of VEGFR-2 and VEGFR-3 in a host cell expressing said receptor.
4. A purified and isolated polypeptide multimer, wherein at least one monomer thereof is a polypeptide according to any one of claims 1-3, and wherein said multimer is capable of binding to at least one of VEGFR-2 and VEGFR-3.

- 148 -

5. A multimer according to claim 4 having a VEGF-C biological activity.
6. A multimer according to claim 4 or 5 wherein at least one monomer thereof is selected from the group consisting of a vascular endothelial growth factor (VEGF) polypeptide, a vascular endothelial growth factor B (VEGF-B) polypeptide, a platelet derived growth factor A (PDGF-A) polypeptide, a platelet derived growth factor B (PDGF-B) polypeptide, a *c-fos* induced growth factor (FIGF) polypeptide, and a placenta growth factor (PlGF) polypeptide.
7. A dimer according to claim 4, 5, or 6.
8. A dimer according to claim 7 wherein each monomer thereof is capable of binding to at least one of VEGFR-2 and VEGFR-3 and has an amino acid sequence comprising a portion of SEQ ID NO: 8 effective to permit such binding.
9. A dimer according to claim 7 or 8 wherein the two monomers are free of covalent attachments to each other.
10. A purified and isolated polypeptide according to claim 1 that is a VEGF-C of vertebrate origin, wherein said VEGF-C has a molecular weight of about 21-23 kD, as assessed by SDS-PAGE under reducing conditions, and wherein said VEGF-C is capable of binding to Flt4 receptor tyrosine kinase (VEGFR-3).
11. A purified and isolated polypeptide according to claim 1 that is a VEGF-C of vertebrate origin, wherein said VEGF-C has a molecular weight of about 30-32 kD, as assessed by SDS-PAGE under reducing conditions, and wherein said VEGF-C is capable of binding to Flt4 receptor tyrosine kinase (VEGFR-3).

- 149 -

12. A polypeptide analog of the VEGF-C according to claim 10 or 11, wherein a conserved cysteine residue in said VEGF-C has been deleted or replaced, and wherein said analog is capable of binding to VEGFR-3 and has reduced VEGFR-2 binding affinity relative to said VEGF-C.
13. A polypeptide analog according to claim 10 or 11 wherein said conserved cysteine residue corresponds to the cysteine at position 156 of SEQ ID NO: 8.
14. A purified polypeptide according to claim 1 that is an analog of human VEGF-C that is capable of binding to at least one of Flt-1 receptor tyrosine kinase (VEGFR-1), KDR receptor tyrosine kinase (VEGFR-2), and Flt4 receptor tyrosine kinase (VEGFR-3).
15. A polypeptide according to claim 14 that binds VEGFR-3 and has reduced VEGFR-2 binding affinity relative to human VEGF-C having an amino acid sequence consisting essentially of amino acids 103-227 of SEQ ID NO: 8.
16. A polypeptide according to claim 14 or 15 that is a VEGF-C ΔC_{156} polypeptide.
17. A VEGF-C ΔC_{156} polypeptide according to claim 16 comprising amino acids 131 to 211 of SEQ ID NO: 8, wherein the cysteine residue at position 156 of SEQ ID NO: 8 has been deleted or replaced.

- 150 -

18. A VEGF-C ΔC_{156} polypeptide according to claim 16 or 17 comprising a continuous portion of SEQ ID NO: 8, said portion having as its amino terminal residue an amino acid between residues 102 and 114 of SEQ ID NO: 8, and having as its carboxy terminal residue an amino acid between residues 212 and 228 of SEQ ID NO: 8, wherein the cysteine residue at position 156 of SEQ ID NO: 8 has been deleted or replaced.

19. A VEGF-C ΔC_{156} polypeptide according to any one of claims 16-18 wherein the cysteine residue at position 156 of SEQ ID NO: 8 has been replaced by a serine residue.

20. A polypeptide according to claim 14 or 15 that is a VEGF-C $\Delta R_{226} \Delta R_{227}$ polypeptide.

21. A VEGF-C $\Delta R_{226} \Delta R_{227}$ polypeptide according to claim 20 having an amino acid sequence comprising amino acids 112-419 of SEQ ID NO: 8, wherein the arginine residues at positions 226 and 227 of SEQ ID NO: 8 have been deleted or replaced.

22. A polypeptide according to claim 14 that is a human VEGF-C^{basic} polypeptide.

23. A polypeptide according to claim 22 having an amino acid sequence comprising residues 131 to 211 of SEQ ID NO: 8, wherein the glutamic acid residue at position 187, the threonine residue at position 189, and the proline residue at position 191 of SEQ ID NO: 8 have been replaced by an arginine residue, a lysine residue, and a histidine residue, respectively.

24. A composition comprising a polypeptide according to any one of claims 1-4 and 10-23, and further comprising a purified myelopoietic growth factor in admixture therewith.

- 151 -

25. A kit useful for modulating myelopoiesis comprising: a first composition comprising a polypeptide according to any one of claims 1-4 and 10-23, packaged with at least one additional composition comprising a myelopoietic growth factor.

26. A composition according to claim 24 or a kit according to claim 25 wherein the myelopoietic growth factor is selected from the group consisting of granulocyte colony stimulating factor (G-CSF), macrophage-CSF (M-CSF), granulocyte-macrophage-CSF (GM-CSF), interleukin-3 (IL-3), stem cell factor (SCF), and combinations thereof.

27. A composition comprising a polypeptide according to any one of claims 1-4 and 10-23, and further comprising a purified polypeptide selected from the group consisting of vascular endothelial growth factor (VEGF) polypeptides, vascular endothelial growth factor B (VEGF-B) polypeptides, platelet derived growth factor A (PDGF-A) polypeptides, platelet derived growth factor B (PDGF-B) polypeptides, *c-fos* induced growth factor (FIGF) polypeptides, and placenta growth factor (PIGF) polypeptides.

28. A polypeptide according to claim 1 that is an analog of human VEGF, wherein a cysteine residue is introduced in the VEGF amino acid sequence at a position selected from residues 53 to 63 of the human VEGF165 precursor having the amino acid sequence set forth in SEQ ID NO: 56, and wherein the polypeptide is capable of binding to at least one of VEGFR-1, VEGFR-2, and VEGFR-3.

29. An analog according to claim 28 wherein said cysteine is introduced at position 58 of the VEGF165 precursor having the amino acid sequence set forth in SEQ ID NO: 56.

30. A purified and isolated nucleic acid comprising a nucleotide sequence encoding a polypeptide according to any one of claims 1-4, 10-23, and 28-29.

- 152 -

31. A vector comprising a nucleic acid according to claim 30.
32. A host cell transformed or transfected with a nucleic acid according to claim 30 or a vector according to claim 31.
33. A method of making a polypeptide capable of specifically binding to at least one of VEGFR-1, VEGFR-2, and VEGFR-3, said method comprising the steps of:
- (a) expressing a nucleic acid according to claim 30 or a vector according to claim 31 in a host cell; and
 - (b) purifying a polypeptide capable of specifically binding to at least one of VEGFR-1, VEGFR-2, and VEGFR-3 from said host cell or from a growth medium of said host cell.
34. An antibody which is specifically reactive with a polypeptide according to any one of claims 1-4, 10-23, and 28-29.
35. A pharmaceutical composition comprising an antibody according to claim 34 in a pharmaceutically-acceptable diluent, adjuvant, excipient, or carrier.
36. A pharmaceutical composition comprising a polypeptide according to any one of claims 1-4, 10-23, and 28-29 in a pharmaceutically-acceptable diluent, adjuvant, excipient, or carrier.
37. A method of modulating the proliferation of mammalian endothelial cells comprising the step of contacting mammalian endothelial cells with a polypeptide in an amount effective to modulate the proliferation of mammalian endothelial cells, said polypeptide selected from the group consisting of:
- (a) a polypeptide according to any one of claims 1-4, 10-23, and 28-29; and
 - (b) a polypeptide comprising an antigen binding portion of an anti-VEGF-C antibody.

- 153 -

38. A method of increasing the proliferation of mammalian endothelial cells according to claim 37, comprising contacting mammalian endothelial cells with a polypeptide in an amount effective to increase the proliferation of mammalian endothelial cells.

39. A method according to claim 37 or 38 wherein said endothelial cells are lymphatic endothelial cells.

40. An *in vivo* method according to claim 39 wherein the contacting step comprises administering to a mammalian subject in need of modulation of the proliferation of lymphatic endothelial cells a composition comprising said polypeptide, in an amount effective to modulate the proliferation of lymphatic endothelial cells *in vivo*.

41. A method according to claim 40 wherein said polypeptide has reduced effect on the permeability of mammalian blood vessels compared to a wildtype VEGF-C polypeptide with an amino acid sequence set forth in SEQ ID NO: 8 from residue 103 to residue 227.

42. A method according to any one of claims 37- 41 wherein said polypeptide is a VEGF-C ΔC_{156} polypeptide.

43. A method for modulating myelopoiesis in a mammalian subject comprising administering to a mammalian subject in need of modulation of myelopoiesis an amount of a polypeptide effective to modulate myelopoiesis, said polypeptide selected from the group consisting of:

(a) a polypeptide according to any one of claims 1-4, 10-23, and 28-29;
and

(b) a polypeptide comprising an antigen binding portion of an anti-VEGF-C antibody.

- 154 -

44. A method according to claim 43 wherein said mammalian subject suffers from granulocytopenia, and said method comprises administering to said subject an amount of a polypeptide effective to stimulate myelopoiesis.

45. A method according to claim 43 or 44 comprising administering to said subject an amount of a polypeptide effective to increase the neutrophil count in blood of said subject.

46. A method according to any one of claims 43-45 wherein said mammalian subject is human.

47. A method according to any one of claims 43-46 further comprising administering to said subject a myelopoietic growth factor selected from the group consisting of granulocyte colony stimulating factor (G-CSF), macrophage-CSF (M-CSF), granulocyte-macrophage-CSF (GM-CSF), interleukin-3 (IL-3), stem cell factor (SCF), and combinations thereof.

48. A method of modulating the proliferation of neutrophilic granulocytes in vitro comprising the step of contacting mammalian stem cells with a polypeptide in an amount effective to modulate the proliferation of neutrophilic granulocytes, said polypeptide selected from the group consisting of:

(a) a polypeptide according to any one of claims 1-4, 10-23, and 28-29;
and

(b) a polypeptide comprising an antigen binding portion of an anti-VEGF-C antibody.

49. A method of modulating the proliferation and/or differentiation of mammalian CD34⁺ progenitor cells comprising contacting mammalian CD34⁺ progenitor cells with a polypeptide according to any one of claims 1-4, 10-23, and 28-29, in an amount effective to modulate the proliferation and/or differentiation of the cells.

- 155 -

50. A method according to claim 49 further comprising contacting the mammalian CD34+ progenitor cells with a myelopoietic growth factor selected from the group consisting of granulocyte colony stimulating factor (G-CSF), macrophage-CSF (M-CSF), granulocyte-macrophage-CSF (GM-CSF), interleukin-3 (IL-3), stem cell factor (SCF), and combinations thereof, in an amounts effective to modulate the proliferation of CD34+ progenitor cells when used in combination with said polypeptide.

51. A method according to claim 49 or 50 wherein the contacting is performed *in vitro* by culturing mammalian CD34+ progenitor cells in the presence of the polypeptide and optionally the myelopoietic growth factor.

52. A method of increasing the number of neutrophils in the blood of a mammalian subject comprising the step of expressing in a cell in a subject in need of an increased number of blood neutrophils a DNA encoding a VEGF-C protein, said DNA operatively linked to a non-VEGF-C promoter or other non-VEGF-C control sequence that promotes expression of said DNA in said cell.

53. A method of increasing the number of endothelial cells in a mammalian subject comprising the step of expressing in a cell in a subject in need of an increased number of endothelial cells a DNA encoding a VEGF-C protein, said DNA operatively linked to a non-VEGF-C promoter or other non-VEGF-C control sequence that promotes expression of said DNA in said cell.

54. A cell comprising a nucleic acid having a sequence encoding human VEGF-C and further comprising a non-VEGF-C promoter sequence or other non-VEGF-C control sequence that increases RNA transcription in said cell of said sequence encoding human VEGF-C.

55. A purified nucleic acid comprising a VEGF-C promoter sequence.

- 156 -

56. A nucleic acid according to claim 55 comprising a portion of SEQ ID NO: 48, wherein said portion is capable of promoting expression of a protein encoding gene operatively linked thereto under conditions wherein VEGF-C is expressed in native host cells.

57. A chimeric nucleic acid comprising a nucleic acid according to claim 55 or 56 operatively connected to a sequence encoding a protein other than a human VEGF-C.

1/15




FIGURE 1

2/15

```

1      PDGF-A      ..... 50
      PDGF-B      ..... .MRTLACLLL
      PlGF-1      ..... MNRCWA.LFL
      VEGF165      .....
      VEGF-B167    .....
      VEGF-C      .....
      MHLGFFSVA  CSLAAALLP GPREAPAAA AFESGLDLS AEPDAGEATA

51     PDGF-A      ..... 100
      PDGF-B      ..... LGGYLAHVL AEEAIPREV IERLARSQIH SIRDQLRLL IDSVGSEDSL
      PlGF-1      ..... SLCCYLRLVS AEGDPIPEEL YEMLSHSIR SFDDLQRLH GDP.GEEDGA
      VEGF165      ..... .MPVM RLFPCL..FLQ LLAGIAL... PAVPPQQW..
      VEGF-B167    ..... .NFFLS..WVH WSLALLLYLH HAKWSQAA..
      VEGF-C      ..... .M SPLLR..RL LAALLQLAPA QAPVSQP...
      YASKDLEEQL RSVSSVDELM TVLYPEYWKM YKQLRKGGW QHNREQANLN

101    PDGF-A      ..... 150
      PDGF-B      ..... DTSRAHGVH ATKHVPEKRP LPIRRKRSI. ....EEAVP AVCKTRTVIY
      PlGF-1      ..... ELDLNMTRSH SGGELES... .LARGRRSLG SLTIAEPAMI AECKTRTEVF
      VEGF165      ..... .ALSAG NGSSEVEVVP FQE.VWGR.. ....SYCRALERLV
      VEGF-B167    ..... .PMAEG GGQNHHEVVK FMD.VYQR.. ....SYCHPIETLV
      VEGF-C      ..... .D APGHQKVVVS WID.VYTR.. ....ATCQPREVVV
      SRTEETIKFA AAHYNTEILK SIDNEWK.. ....TQCMPEVCI

151    PDGF-A      ..... 200
      PDGF-B      ..... EIPRSQVDPT SANFLIWPPC VEVKRCTGCC NTSSVKCQPS RVHHRSVKVA
      PlGF-1      ..... EISRRLLDRT NANFLVWPPC VEVQRCSCGCC NNRNVQCRPT QVQLRPVQVR
      VEGF165      ..... DVVSEYPSEV ..EHMFSPSC VSLRCTGCC GDENLHCVPV ETANVTMQLL
      VEGF-B167    ..... DIFQEYPDEI ..EYIFKPSC VPLMRCCGCC NDEGLECVPT EESNITMQIM
      VEGF-C      ..... PLTVELMGTV ..AKQLVPSC VTVQRCGGCC PDDGLECVPT GQHQVRMQIL
      DVGKEFGVAT ..NTFFKPPC VSVYRCGGCC NSEGLQCMNT STSYLSKTLF

```

FIGURE 2A

3/15

```

201
PDGF-A      KVEYVRKKPK LKEVQVRLEE HLEACAT... 250
PDGF-B      KIEIVRKKPI FKKATVTLED HLACKCETVA AARPVTRSPG GSQEQRAKTP
PIGF-1      KIRSG..DRP .SYVELTFSQ HVRCECRPLR EK.....
VEGF165     RIKPH..QGQ .HIGEMSFLQ HNKCECRPKK DR.....
VEGF-B167   MIRYP..SSQ ..LGEMSLEE HSQCECRPKK KD.....
VEGF-C      EITVPLSQGP .KPVITISFAN HTSCRCMSKL DVYRQVHSII RRSPLATLPQ

251
PDGF-A      DTDVR..... 300
PDGF-B      QTRVTIRTVR VRRPPKGKHR KFKHTHDKTA LKETLGA...
PIGF-1      .....
VEGF165     .....
VEGF-B167   .....
VEGF-C      CQAANKTCPT NYMWNHICR CLAQEDFMFS SDAGDDSTDG FHDICGPNKE

301
PDGF-A      ..... 350
PDGF-B      .....
PIGF-1      .....
VEGF165     QDPQTCCKSC KNTDS.RCKA RQLELNERTC RCDKPRR...
VEGF-B167   PDPRTCRCRC RRRSFLRCQG RGLELNPDT C RCRKLRR...
VEGF-C      LDEETCQCVC RAGLRPASC PHKELDRNSC QCVCKNKLFP SQCGANREFD

```

FIGURE 2 B

4/15

351 PDGF-A 400
PDGF-B
PlGF-1
VEGF165
VEGF-B167
VEGF-C
ENTCQCVCKR TCPRNQPLNP GKCACTES POKCLLKGGK FHHQTCSCYR

401 PDGF-A 434
PDGF-B
PlGF-1
VEGF165
VEGF-B167
VEGF-C
RPCTNRQKAC EPGFSYSEEV CRCVPSYWKR PQMS

FIGURE 2 C

5/15




FIGURE 3

6/15




FIGURE 4

7/15

VEGF-C alignment

	1				50
Hum	HMLLGFFSVA	CSLLAAALLP	GPREAPAAAA	AFESGLDLSD	AEPDAGEATA
Mou	MHLLCFLSLA	CSLLAAALIP	SPREAPATVA	AFESGLGFSE	AEPDGGEVKA
Qua	MHLLLEMLSLG	CCLAAGAVLL	GPRQPPVA.A	AYESGHGYE	EEPGAGEPKA
	51				100
Hum	YASKDLEEQL	RSVSSVDELM	TVLYPEYWKM	YKCQLRKGGW	QHNREQANLN
Mou	FEGKDLEEQL	RSVSSVDELM	SVLYPDYWKM	YKCQLRKGGW	Q....QPTLN
Qua	HASKDLEEQL	RSVSSVDELM	TVLYPEYWKM	FKCQLRKGGW	QHNREHSSSD
	101				150
Hum	SRTEETIKFA	AAHYNTEILK	SIDNEWRTKQ	CMPREVCIDV	GKEFGVATNT
Mou	TRTGDSVKFA	AAHYNTEILK	SIDNEWRTKQ	CMPREVCIDV	GKEFGAATNT
Qua	TRSDDSLKFA	AAHYNAEILK	SIDTEWRKTQ	GMPREVCVDL	GKEFGATTNT
	151				200
Hum	FFKPPCVSVY	RCGGCCNSEG	LQCMNTSTSY	LSKTLFEITV	PLSQGPKPVT
Mou	FFKPPCVSVY	RCGGCCNSEG	LQCMNTSTGY	LSKTLFEITV	PLSQGPKPVT
Qua	FFKPPCVSIY	RCGGCCNSEG	LQCMNISTNY	ISKTLFEITV	PLSHGPKPVT
	201				250
Hum	ISFANHTSCR	CMSKLDVYRQ	VHSIIRRLSP	ATLPQCOAAN	KTCPTNYMWN
Mou	ISFANHTSCR	CMSKLDVYRQ	VHSIIRRLSP	ATLPQCOAAN	KTCPTNYVWN
Qua	VSFANHTSCR	CMSKLDVYRQ	VHSIIRRLSP	ATQTQCHVAN	KTCPKNHVWN
	251				300
Hum	NHICRCCLAQE	DFMFSSDAGD	DSTDGFHDIC	GPNKELDEET	CQCVCRAGLR
Mou	NYMCRCLAQQ	DFIFYSNVED	DSTNGFHDVC	GPNKELDEDT	CQCVCCKGGLR
Qua	NQICRCCLAQH	DFGFSSHLGD	SDTSEGFHIC	GPNKELDEET	CQCVCCKGGVR
	301				350
Hum	PASCGPHKEL	DRNSCQCVCCK	NKLFPSCCGA	NREFDENTCQ	CVCKRTCPRN
Mou	PSSCGPHKEL	DRDSCQCVCCK	NKLFPNSCGA	NREFDENTCQ	CVCKRTCPRN
Qua	PISCGPHKEL	DRASCQCMCK	NKLLPSSCGP	NKEFDEEKQ	CVCKKTCPKH
	351				400
Hum	QPLNPGKCAC	ECTESPQKCL	LKGKKFHHQT	CSCYRRPCTN	RQKACEPGFS
Mou	QPLNPGKCAC	ECTENTQKCF	LKGKKFHHQT	CSCYRRPCAN	RLKHCDPGLS
Qua	HPLNPAKCIC	ECTESPNKCF	LKGKRFHHQT	CSCYRPPCTV	RTKRCDAAGFL
	401	420			
Hum	YSEEVCRCPV	SYWKRPMQMS*			
Mou	FSEEVCRCPV	SYWKRPHLN.			
Qua	LAEEVCRCPV	TSWKRPLMN*			

FIGURE 5

8/15

FIGURE 6A




FIGURE 6B




FIGURE 6C

9/15

Media




FIGURE 7A

Lysates




FIGURE 7B

10/15




FIGURE 8A




FIGURE 8B

11/15





FIGURE 9



HUMAN				
Exon length	Donor site	Intron length	Acceptor site	
.....G...E...A...T(49)A...Y...A...S.	
E1GGC.GAG.GCC.ACG.gtaggtctgcgt...>10.kb..TTTCTTTGACAG.GCT.TAT.GCA.AGC	
.....E...I...L...K(116)S...I...D...N.	
E2 .214.bp..GAG.ATC.TTG.AAA.Agtaagtatggg...1.6.kb...atgacttgacagGT.ATT.GAT.AATL...F...E...I.	
.....L...S...K...T(180)	
E3 .191.bp..CTC.AGC.AAG.ACG.gtggtattgt.....9.kb.cccttctttag.TTA.TTT.GAA.ATTC...Q...A...A.	
.....T...L...P...Q(231)D...S...T...D.	
E4 .152.bp..ACA.CTA.CCA.CAGtgagatgaattaa>10.kb..ttcttccaaagG.TGT.CAG.GCA.GCG	
.....A...G...D... (266)	
E5 .107.bp..GCT.GGA.GAT.Ggtagcagaatg.....301.bp...ctattgtcttagAC.TCA.ACA.GATC...Y...R...R.	
.....Q...T...C...S(378)	
E6 .334.bp..CAA.ACA.TGC.AGgtaagatcc.....>10.kb..tggtctccttagC.TGT.TAC.AGA.CGG	
.....Q...M...S(419) Stop	
E7 .(501).bp..CAA.ATG.AGC.TAA.GTATGTACTGTT...ATTGTATTAT	

FIGURE 11A

14/15

MOUSE			
Exon length	Donor site	Intron length	Acceptor site
.....G...E...V...K(49).....			
E1GGC.GAG.GTC.AAG.gtagtgcaagg.>10.kb.attgtctttgacag.GCT.TTT.TGA.AGG.			A...F...E...G.
.....E...I...L...K(116).....			
E2 201.bp..GAG.ATC.CTG.AAA.Agtaagtag.....4.kb....tgtgactcgacagGT.ATT.GAT.AAT			S...I...D...N.
.....L...S...K...T(180).....			
E3 191.bp..CTC.AGC.AAG.ACG.gtaggtat.....9.kb..ttgtcccctttag.TTG.TTT.GAA.ATT			L...F...E...I.
.....T...L...P...Q(231).....			
E4 152.bp..ACA.TTA.CCA.CAGtgagtatg.....10.kb.gtctccccaaaagG.TGT.CAG.GCA.GCT			C...Q...A...A.
.....N...V...E...D(266).....			
E5 107.bp..AAT.GTT.GAA.GAT.Ggtaagtaaaa...350.bp.....tctagAC.TCA.ACC.AAT			D...S...T...N.
.....Q...T...C...S(378).....			
E6 334.bp..CAA.ACA.TGC.AGgtaaggagtgt.....6.kb..ttttccccttagT.TGT.TAC.AGA.AGA			C...Y...R...R.
.....H...L...N(415)Stop.....polyA.....			
E7 506.bp..CAT.CTG.AAC.TAA.GATCATACC...ATTGTATTATAAgctgtgaag			

FIGURE 11B

INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/US 98/01973

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/52 C12N15/10 C07K16/24 C12Q1/68
C12N15/62 G01N33/50 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N C12Q A01K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOUKOV V ET AL: "A NOVEL VASCULAR ENDOTHELIAL GROWTH FACTOR, VEGF-C, IS A LIGAND FOR THE FLT4 (VEGFR-3) AND KDR (VEGFR-2) RECEPTOR TYROSINE KINASES" EMBO JOURNAL, vol. 15, no. 2, 1996, pages 290-298, XP002022272 see the whole document	1-5,7,8, 10,11, 30-34, 37-39,54
X	LEE, J., ET AL. : "vascular endothelial growth factor-related protein: a ligand and specific activator of the tyrosine kinase receptor flt4" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 93, March 1996, pages 1988-1992, XP002066360 see the whole document	1-3,10, 11, 30-34, 37,38,54
- / - -		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

29 May 1998

Date of mailing of the international search report

03.07.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Holtorf, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/01973

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 39515 A (HUMAN GENOME SCIENCES INC ;ROSEN CRAIG A (US); HU JING SHAN (US);) 12 December 1996 pages 21,34; examples 2,3,4,5,6 ---	1-3,10, 11, 30-34, 36-38, 40,54
X	LEE,J., ETAL. : "VASCULAR ENDOTHELIAL GROWTH FACTOR RELATED PROTEIN (vrp): A LIGAND AND SPECIFIC ACITVATOR OF THE TYROSINE KINASE RECEPTOR Flt4" EMBL SEQUENCE DATA LIBRARY, 10 January 1996, HEIDELBERG, GERMANY, XP002066361 ACCESSION NO. U4142 ---	55,56
X	JOUKOV,V., ET AL. : "A NOVEL VASCULAR ENDOTHELIAL GROWTH FACTOR VEGF-C IS A LIGAND FOR THE FLT4 (VEGFR-3)AND KDR (VEGFR-2) RECEPTOR TYROSINE KINASES" EMBL SEQUENCE DATA LIBRARY, 1 February 1996, HEIDELBERG, GERMANY, XP002066362 ACCESSION NO. X94216 ---	55,56
A	COHEN T ET AL: "VEGF121, A VASCULAR ENDOTHELIAL GROWTH FACTOR (VEGF) ISOFORM LACKING HEPARIN BINDING ABILITY, REQUIRES CELL-SURFACE HEPARAN SULFATES FOR EFFICIENT BINDING TO THE VEGF RECEPTORS OF HUMAN MELANOMA CELLS" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 19, 12 May 1995, pages 11322-11326, XP002061896 see the whole document ---	1-57
P,X	WO 97 05250 A (UNIV HELSINKI LICENSING ;ALITALO KARI (FI); JOUKOV VLADIMIR (FI)) 13 February 1997 cited in the application pages 10,13-17, 27; examples 8,9,11,14,15,19,22,27,28,29; claims ---	1-11, 30-34, 36-40, 43,53-56
P,X	JOUKOV,V., ET AL. : "PROTEOLYTIC PROCESSING REGULATES RECEPTOR SPECIFICITY AND ACTIVITY OF VEGF-C" THE EMBO JOURNAL, vol. 16, no. 13, June 1997, pages 3898-3911, XP002066363 see the whole document ---	1-5, 7-11,14, 15,20, 21, 30-34,54
3 P,X	WO 97 09427 A (GENENTECH INC) 13 March 1997 pages 3,5,9,10,11,24,25,34; example 5,7 ---	1-3,10, 11, 30-38,54
3	---	-/--

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/01973

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 97 17442 A (IMMUNEX CORP) 15 May 1997 pages 3,7,8,17,19,21,24; examples ---	1-3,10, 11, 30-34, 36-38, 40,54
P,X	ACHEN, M.G., ET AL.: "VASCULAR ENDOTHELIAL GROWTH FACTOR D (VEGF-D) IS A LIGAND FOR THE TYROSINE KINASES VEGF RECEPTOR 2 (Flk1) AND vegf RECEPTOR 3 (Flt4)" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 95, January 1998, pages 548-553, XP002066364 abstract, pages 548, right column, page 549, 551; figures 1 + 4; ---	1-3, 30-34
P,X	JELTSCH,M., ET AL. : "HYERPLASIA OF LYMPHATIC VESSELS IN VEGF-C TRANSGENIC MICE" SCIENCE, vol. 276, 30 May 1997, pages 1423-1425, XP002066365 see the whole document ---	1-3,10, 11, 30-32, 37-40, 53,54
T	JOUKOV,V., ET AL.: "A RECOMBINANT MUTANT VASCULAR ENDOTHELIAL GROWTH FACTOR-C THAT HAS LOST VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR-2 BINDING , ACTIVATION, AND VASCULAR PERMEABILITY ACTIVITIES" THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 12, 20 March 1998, pages 6599-6602, XP002066366 see the whole document -----	1-57

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 98/01973

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 37-47 and 49,50,52,53 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
The polypeptide analog of claim 13 was searched as referring to the analog mentioned in claim 12.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/01973

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9639515 A	12-12-1996	AU 6046796 A EP 0837934 A	24-12-1996 29-04-1998
WO 9705250 A	13-02-1997	AU 6616996 A EP 0842273 A	26-02-1997 20-05-1998
WO 9709427 A	13-03-1997	AU 7012896 A	27-03-1997
WO 9717442 A	15-05-1997	AU 1116297 A	29-05-1997

